

World Journal of *Orthopedics*

World J Orthop 2017 August 18; 8(8): 606-659





EDITORIAL

- 606 Ultrasound diagnosis of fractures in mass casualty incidents
Abu-Zidan FM

MINIREVIEWS

- 612 Postoperative deep shoulder infections following rotator cuff repair
Atesok K, MacDonald P, Leiter J, McRae S, Stranges G, Old J
- 619 Synthesis of evidence for the treatment of intersection syndrome
Balakatounis K, Angoules AG, Angoules NA, Panagiotopoulou K

ORIGINAL ARTICLE

Retrospective Study

- 624 Ponseti method treatment of neglected idiopathic clubfoot: Preliminary results of a multi-center study in Nigeria
Adegbehingbe OO, Adetiloye AJ, Adewole L, Ajodo DU, Bello N, Esan O, Hoover AC, Ior J, Lasebikan O, Ojo O, Olasinde A, Songden D, Morcuende JA

Observational Study

- 631 Functional outcomes of traumatic and non-traumatic rotator cuff tears after arthroscopic repair
Abechain JJK, Godinho GG, Matsunaga FT, Netto NA, Daou JP, Tamaoki MJS

SYSTEMATIC REVIEWS

- 638 Ipsilateral femur and tibia fractures in pediatric patients: A systematic review
Anari JB, Neuwirth AL, Horn BD, Baldwin KD

META-ANALYSIS

- 644 Femoral positioning influences ipsi-and contralateral anterior cruciate ligament rupture following its reconstruction: Systematic review and meta-analysis
de Campos GC, Teixeira PEP, Castro A, Alves Junior WM

CASE REPORT

- 651 Open wound management of esophagocutaneous fistula in unstable cervical spine after corpectomy and multilevel laminectomy: A case report and review of the literature
Elgafy H, Khan M, Azurdia J, Peters N
- 656 Bennett's fracture associated with fracture of Trapezium - A rare injury of first carpo-metacarpal joint
Goyal T

Contents

World Journal of Orthopedics
Volume 8 Number 8 August 18, 2017

ABOUT COVER

Editorial Board Member of *World Journal of Orthopedics*, Melanie Jane Franklyn, PhD, Academic Fellow, Academic Research, Department of Mechanical Engineering, The University of Melbourne, Melbourne, Victoria 3010, Australia

AIM AND SCOPE

World Journal of Orthopedics (*World J Orthop*, *WJO*, online ISSN 2218-5836, DOI: 10.5312) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJO covers topics concerning arthroscopy, evidence-based medicine, epidemiology, nursing, sports medicine, therapy of bone and spinal diseases, bone trauma, osteoarthritis, bone tumors and osteoporosis, minimally invasive therapy, diagnostic imaging. Priority publication will be given to articles concerning diagnosis and treatment of orthopedic diseases. The following aspects are covered: Clinical diagnosis, laboratory diagnosis, differential diagnosis, imaging tests, pathological diagnosis, molecular biological diagnosis, immunological diagnosis, genetic diagnosis, functional diagnostics, and physical diagnosis; and comprehensive therapy, drug therapy, surgical therapy, interventional treatment, minimally invasive therapy, and robot-assisted therapy.

We encourage authors to submit their manuscripts to *WJO*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

INDEXING/ABSTRACTING

World Journal of Orthopedics is now indexed in Emerging Sources Citation Index (Web of Science), PubMed, PubMed Central and Scopus.

FLYLEAF

I-III Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Ya-Jing Lu*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Fang-Fang Ji*
Proofing Editorial Office Director: *Jin-Lei Wang*

NAME OF JOURNAL
World Journal of Orthopedics

ISSN
ISSN 2218-5836 (online)

LAUNCH DATE
November 18, 2010

FREQUENCY
Monthly

EDITORS-IN-CHIEF
Quanjun (Trey) Cui, MD, Professor, Department of Orthopaedic Surgery, School of Medicine, University of Virginia, Charlottesville, VA 22908, United States

Bao-Gan Peng, MD, PhD, Professor, Department of Spinal Surgery, General Hospital of Armed Police Force, Beijing 100039, China

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com>

www.wjgnet.com/2218-5836/editorialboard.htm

EDITORIAL OFFICE
Xiu-Xia Song, Director
World Journal of Orthopedics
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.fjpublishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive,
Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.fjpublishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
August 18, 2017

COPYRIGHT
© 2017 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
<http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.fjpublishing.com>

Femoral positioning influences ipsi-and contralateral anterior cruciate ligament rupture following its reconstruction: Systematic review and meta-analysis

Gustavo Constantino de Campos, Paulo Eduardo Portes Teixeira, Alex Castro, Wilson de Mello Alves Junior

Gustavo Constantino de Campos, Department of Orthopaedics, Campinas's State University (UNICAMP), Campinas 13083-970, Brazil

Gustavo Constantino de Campos, Paulo Eduardo Portes Teixeira, Alex Castro, Wilson de Mello Alves Junior, Instituto Wilson Mello, Research and Study Center, Campinas 13080-650, Brazil

Author contributions: de Campos GC contributed to literature search, data extraction, data analysis and interpretation, article draft, article revision, final approval; Teixeira PEP contributed to literature search, data extraction, data analysis and interpretation, article draft, final approval; Castro A contributed to statistical analysis, data interpretation, article revision, final approval; Alves Junior WM contributed to conception and design of the study, critical revision, final approval.

Conflict-of-interest statement: The authors deny any conflict of interest.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Gustavo Constantino de Campos, MD, PhD, Instituto Wilson Mello, Research and Study Center, Rua José Rocha Bonfim, 214, ed. Chicago, 1o. andar, Campinas 13080-650, Brazil. gustavoccampos@terra.com.br
Telephone: +55-19-35212121
Fax: +55-19-35212121

Received: January 20, 2017

Peer-review started: January 20, 2017

First decision: February 15, 2017

Revised: March 9, 2017

Accepted: March 23, 2017

Article in press: March 25, 2017

Published online: August 18, 2017

Abstract

AIM

To systematically review the incidence of ipsilateral graft re-rupture and contralateral anterior cruciate ligament (ACL) rupture following its reconstruction, with special attention to the femoral drilling technique.

METHODS

Systematic review and meta-analysis of high-level prospective studies searched in MEDLINE database following PRISMA statement. The rate of ipsilateral graft re-rupture and contralateral rupture in patients submitted to either transtibial (TT) technique (isometric) or anteromedial (AM) technique (anatomic) was compared.

RESULTS

Eleven studies met the criteria and were included in final analysis. Reconstructions using the AM technique had a similar chance of contralateral ACL rupture when compared to the chance of ipsilateral graft failure (OR = 1.08, $P = 0.746$). In reconstructions using TT technique, the chance of contralateral ACL rupture was approximately 1.5 times higher than ipsilateral graft failure (OR = 1.49, $P = 0.048$). Incidence of contralateral lesions were similar among the techniques TT (7.4%) and AM (7.0%) ($P = 0.963$), but a trend could be noticed with a lower incidence of lesion in the ipsilateral limb when using the TT technique (4.9%) compared to the AM technique (6.5%) ($P = 0.081$).

CONCLUSION

ACL reconstruction by TT technique leads to lower incidence of graft re-injury than contralateral ACL lesion. There is no difference between the chance of re-injury after AM technique and the chance of contralateral ACL lesion (native ligament) with either technique.

Key words: Anterior cruciate ligament; Anterior cruciate ligament reconstruction; Arthroscopy; Graft survival

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: There is no convincing evidence that anatomic reconstruction leads to better clinical outcomes than transtibial (TT) reconstruction. Moreover, data suggests that it could lead to an increased risk of graft re-rupture. We found that anterior cruciate ligament (ACL) reconstruction by TT technique led to lower incidence of graft re-rupture than contralateral ACL lesion. The chance of graft re-rupture after anteromedial (AM) technique was the same of contralateral ACL lesion. There was no difference between contralateral lesion after both techniques and re-rupture after AM technique, what could mean that re-rupture chance after AM technique is indeed closer to normal knee, and, in fact, it is the TT technique's re-tear incidence that is lower than it should be.

de Campos GC, Teixeira PEP, Castro A, Alves Junior WM. Femoral positioning influences ipsi-and contralateral anterior cruciate ligament rupture following its reconstruction: Systematic review and meta-analysis. *World J Orthop* 2017; 8(8): 644-650 Available from: URL: <http://www.wjgnet.com/2218-5836/full/v8/i8/644.htm> DOI: <http://dx.doi.org/10.5312/wjo.v8.i8.644>

INTRODUCTION

Optimal reconstruction technique for an anterior cruciate ligament (ACL) tear still on debate. In the last fifteen years literature has shown a significant shift in trends regarding graft positioning^[1]. The "isometric" femoral graft positioning, made through the tibial tunnel (transtibial technique) and very popular during the last two decades of the twentieth century, is gradually being loathed^[2]. Since the introduction of the anterior cruciate anatomic reconstruction concept^[3], the pursuit of the ideal graft positioning has led to the so-called "anatomic" single-bundle reconstruction techniques, which can be accomplished by either by an anteromedial approach^[4,5] as well as an outside-in femoral drilling^[6].

Despite the plethora of anatomical and biomechanical studies suggesting incorrect graft positioning and inadequate knee rotational stability with transtibial (TT) femoral drilling technique, there is no convincing evidence that anatomic reconstruction leads to better clinical outcomes^[7]. Moreover, recent data suggests that the anteromedial (AM) technique for femoral drilling

could lead to an increased risk of graft re-rupture^[8].

We hypothesized that the increased risk of graft re-rupture observed in anatomical reconstructions could in fact represent an approximation to the "normal knee" ACL rupture risk, that could be represented by the risk of contralateral knee lesion. This also would explain the odd finding that there is a higher incidence of contralateral knee ACL lesion when compared to graft re-lesion following ACL reconstruction by isometric technique^[9].

Therefore, the aim of this study is to systematically review the incidence of ipsilateral graft re-rupture and contralateral ACL rupture following ACL reconstruction, with special attention to the femoral drilling technique.

MATERIALS AND METHODS

Systematic review and meta-analysis of studies including patients submitted to ACL reconstruction, registered at PROSPERO under the number CRD 42015019336. PRISMA statement guidelines were followed for conducting and reporting meta-analysis data.

Literature review

On September 13, 2015, a systematic literature search of the MEDLINE database was performed independently by two of the authors (Gustavo Constantino de Campos and Paulo Eduardo Portes Teixeira) using the following terms: "Anterior cruciate ligament" AND "contralateral" AND "reconstruction" AND "follow-up". The initial search yielded 189 results. Eligibility criteria were original studies that included adult patients submitted to ACL reconstruction, in English language. Title and abstract evaluation suggested 156 articles for full text revision. Studies were excluded if did not specifically described drilling technique for ACL femoral tunnel confection, did not report the incidence of ipsilateral graft failure and incidence of contralateral ACL lesion during follow-up, if they were experimental studies (biomechanics or animal), if patients were submitted to revision surgery, double bundle or multiple ligaments reconstruction, or graft selection comprising allograft, artificial devices or harvested from contralateral knee. Risk of bias was minimized including only grade I to III prospective studies with a minimum of 75% of patients at final follow-up.

From the 156 articles reviewed, 145 articles were excluded, resulting in 11 articles for final analysis^[10-20]. The references of the remaining eleven articles were reviewed with no additional studies for inclusion identified. These 11 studies provided the data for the present analysis. Literature review is summarized in Figure 1.

Data extraction

Extracted data included type of study, evidence level, patient demographics, follow-up duration, loss of follow-up, surgical technique, graft choice, ACL graft re-lesion




Figure 1 Prisma flow diagram. ACL: Anterior cruciate ligament.

incidence and contralateral ACL rupture incidence. Data were extracted by two authors independently (GCC and PEPT). Discrepancies were resolved by consensus.

Statistical analysis

A meta-analysis of the incidence of contralateral vs ipsilateral lesions to each of surgical techniques (AM and TT) was performed by "Comprehensive Meta-Analysis software version 3.3.070". The heterogeneity of variances between studies was examined by χ^2 test for heterogeneity and I^2 statistics (proportion of the total variance due to heterogeneity) and χ^2 (effect size variance between studies)^[21,22]. When a significant heterogeneity was observed between studies, the combined effects across studies were analyzed by random effects model. When there was no significant heterogeneity between studies, these effects were analyzed using fixed effects model^[22].

The estimate of the pooled effect (pooled odds ratio) between studies was calculated using the Mantel-Haenszel method (MH OR). Publication biases were analyzed by the asymmetry in the Begg and Mazumdar's funnel plot^[23]. Additionally, an association analysis between the surgical technique used (AM or TT) and the incidence of total ipsilateral and contralateral lesions of all publications was performed using χ^2 2 × 2 test statistic in SPSS 18.0 software (SPSS Inc., Chicago, United States).

The chance of injury occurrence was calculated by OR

using simple logistic regression in which was considered the sum of the number of events (contralateral or ipsilateral lesions) and the sum of the number of cases of all publications that have used the technique AM or TT separately. The significance level (α) used for all analyzes was 5% ($P < 0.05$).

RESULTS

Follow-up


All studies had a minimum of 2 years of follow-up after ACL reconstruction. The duration of follow-up's ranged from 2 to 15 years. Data concerning the incidence of lesions were based on the total amount of individuals who have completed 100% of follow-up for each study. The percentage of individuals who completed the follow-up in each study ranged from 75% to 98% (Table 1).

Quality evaluation and potential biases

Study quality was assessed as recommended by previous studies^[23,24]. Of the eleven studies included in this meta-analysis, 6 (55%) were classified as level of evidence 1, 3 (27%) with level of evidence 2 and 2 (18%) with evidence of Level 3 (Table 1). The potential bias across studies were evaluated by Begg and Mazumdar's funnel plot for the incidence of contralateral and ipsilateral lesions in each of the AM and TT techniques, which were predominantly symmetrical, suggesting lack of significant biases in the publications

Table 1 Data of the studies *n* (%)

Author	Year	Evidence level	No. of initial patients	No. of patients included in follow-up	Follow-up (yr)	No. of contralateral injuries	No. of Ipsilateral injuries
Anteromedial drilling technique							
Webb	2013	3	200	181 (91)	15	19 (10.5)	35 (19.3)
Shelbourne	2009	2	1820	1415 (78)	14	75 (5.3)	61 (4.3)
Pinczewsky	2007	2	180	178 (99)	10	29 (16.3)	19 (10.7)
Sajovic	2006	1	64	61 (95)	5	5 (8.2)	4 (6.6)
Shaieb	2002	1	82	70 (85)	2	6 (8.6)	4 (5.7)
Total	-	-	2346	1905 (81)	-	134 (7.0)	123 (6.5)
Transtibial drilling technique							
Barenius	2014	1	164	134 (82)	14.1	6 (4.5)	6 (4.5)
Webster	2014	3	750	561 (75)	4.8	42 (7.5)	25 (4.5)
Holm	2010	1	72	57 (79)	10	7 (12.3)	6 (10.5)
Keays	2007	2	62	62 (100)	6	5 (8.1)	1 (1.6)
Drogset	2005	1	41	38 (93)	2	1 (2.6)	3 (7.9)
Aune	2001	1	72	64 (89)	2	3 (4.7)	3 (4.7)
Total	-	-	1161	916 (79)	-	64 (7.4)	44 (4.9)

**Figure 2** Funnel plot for the incidence of contralateral and ipsilateral lesions in each of the anteromedial (A) and transtibial (B) techniques included in the meta-analysis. MH: Mantel-Haenszel.

included in the meta-analysis (Figure 2).

Incidence of ipsilateral vs contralateral rupture

After the meta-analysis and combining the data, it was observed that ACL reconstructions using the AM technique had a similar chance of contralateral ACL rupture when compared to the chance of ipsilateral graft failure (MH OR = 1.08, 95%CI: 0.67 to 1.75, $P = 0.746$; Figure 3). However, after reconstructions using the TT technique, the chance of contralateral ACL rupture was approximately 1.5 times higher than the chance of ipsilateral graft failure (MH OR = 1.49, 95%CI: 1.00 to 2.21, $P = 0.048$, Figure 3). There was no heterogeneity of variances between studies using the TT technique ($I^2 = 0.0\%$, $\chi^2 = 0.0$, $P = 0.517$, Figure 3).

Nevertheless, those who used AM technique showed significant heterogeneity of variances ($I^2 = 56.9\%$, $\chi^2 = 0.151$, $P = 0.054$, Figure 3). Finally, from the publications included in the present meta-analysis (Table 1), no differences in the incidence of contralateral lesions were identified among the techniques TT (7.4%)

and AM (7.0%) (OR = 0.99, 95%CI: 0.73 to 1.35, $P = 0.963$). But a trend could be noticed with a lower incidence of lesion in the ipsilateral limb when using the TT technique (4.9%) compared to the AM technique (6.5%) (OR = 0.73, 95%CI: 0.51 to 1.04, $P = 0.081$).

DISCUSSION

The present study found no difference between the risk of an ipsilateral graft re-rupture and a contralateral ACL rupture in individuals operated with the AM technique. There was also no difference in the risk of a contralateral ACL rupture when comparing both techniques (7%). However, the present analysis found a lower rate of ipsilateral graft re-rupture in patients operated with TT technique (4.9%; $P = 0.048$). This is the first analysis focusing the influence of femoral drilling technique over the graft failure and contralateral ACL lesion after ACL reconstruction.

Ipsilateral graft re-lesion is, for obvious reasons, an undesirable event, occurring in 1.8%^[14] to 10.4%^[25]




Figure 3 Results of individual studies along with a Forest plot that graphically displays the point estimates on a common scale surrounded by the 95%CI (indicated by the horizontal lines). The diamond represents the pooled effect between studies. M-H: Mantel-Haenszel; df: Degrees of freedom.

of ACL reconstructed patients during the follow-up. Literature shows higher rates of contralateral ACL lesions than ipsilateral graft failure after ACL reconstruction^[9]. An explanation to this finding could be inadequate rehabilitation^[16] or even the assumption that the neoligament would be stronger than the original ACL^[16]. Some authors attribute such finding to the protective role of the uninjured limb that renders it more suitable to lesion^[26].

However, recent studies found a decrease in graft in situ forces when positioned in more vertical locations (isometric position), indicating that the femoral tunnel drilled through transtibial approach will result in a more vertical graft with in situ force lower than the in situ force found in the native ACL^[27-29]. Therefore, the lower graft failure rates found in TT reconstruction could be because the neoligament simply is not submitted to the physiological forces that occur in the "normal" knee. With that in mind, one could speculate that it is not the AM technique that presents a higher failure rate, but the TT graft failure rate that is lower than it should be.

Anatomic ACL reconstruction results in graft inclination angle closer to the native ACL inclination angle and graft force and knee anterior laxity closer to those of the native knee^[29]. It is our understanding that this could be the explanation to the findings of the present analysis, since we found same risk of contralateral ACL rupture ("normal" knee) for both techniques, with no difference for graft failure in the AM group (7%), and lower incidence of ipsilateral graft failure in TT group (4.9%, $P = 0.048$).

It is fundamental, thus, to provide information regarding femoral drilling technique when discussing ipsi or contralateral lesions rates during follow-up. Most of studies just focus on the graft choice and patient demographics. The vast majority of studies initially included in our analysis did not adequately describe the ACL reconstruction technique used. This is in accordance to van Eck *et al.*^[30], who examined 74 studies that claimed to use "anatomic technique" for ACL reconstruction and found a gross under-reporting of specific operative technique data.

Rahr-Wagner *et al.*^[8] found increased risk of revision in patients undergone to ACL reconstruction with femoral drilling through AM portal when compared to TT technique in the Danish Knee Ligament Reconstruction Register. They explained this finding as a result of a learning curve due to the introduction of a new and more complex technique. The higher force over the more anatomically placed grafts could be a more accurate explanation.

Our results indicate that the increased risk is in fact an approximation to the expected failure rate of the normal knee, that could be represented by the rate of contralateral ACL lesion. van Eck *et al.*^[31] found that the majority of graft failures following anatomic ACL reconstruction occurred between six and nine months postoperatively, precisely the commonly recommended period for return to sports. Although there has been a transition toward the "anatomic" reconstruction over the last decades, rehabilitation protocols still the same. Maybe the higher forces over the graft could alter the

time frame for complete graft healing and maturation. More studies are needed on that matter.

In another point of view, one can argue that, since there is no convincing data on the superiority of anatomic technique regarding clinical outcomes, it would be better to choose a technique with a lower rate of graft failure (TT technique). This question will only be answered as longer follow-up of anatomic reconstruction series successfully demonstrates better clinical outcomes and lower evolution to osteoarthritis.

We acknowledge that this meta-analysis has strengths and limitations. One limitation is the lack of demographic data, such as age, gender, body mass index and level of activity. Post-operative rehabilitation protocols were not analyzed or even considered. We acknowledge the importance of rehabilitation data. However, meta-analysis was performed comparing two groups operated by the same authors, therefore subjected to same rehabilitation protocols. Also, we did not include anatomic reconstruction by out-in femoral drilling technique. We chose to use only AM technique to standardize our analysis.

In addition, we were not able to show the data on direct comparison between AM and TT techniques on graft failure and contralateral ACL rupture. To perform this comparison and present it in a meta-analysis format, the comparison effect between techniques would have to be presented individually in each study selected for this meta-analysis. All selected studies for this review were independent and presented only intra-technical comparisons. Although we performed an exploratory analysis to test the differences between the incidences of graft failure and contralateral ACL rupture among AM and TT techniques, it could be biased. This issue is still a gap in the literature and more clinical studies are needed to conduct future meta-analyses to clarify the subject. The major strength is the rigorous criteria used. We only included studies that reported both ipsi- and contralateral failure information, thus ensuring proper comparison when performing the meta-analysis. Moreover, we only included high quality prospective studies with high level of evidence and loss of follow-up lower than 25%.

The ACL reconstruction by transtibial technique leads to lower incidence of graft re-injury than contralateral ACL lesion. There is no difference between the chance of re-injury after anteromedial technique and the chance of contralateral ACL lesion (native ligament).

COMMENTS

Background

Optimal reconstruction technique for an anterior cruciate ligament (ACL) tear still on debate. Evidence regarding the influence of graft femoral positioning over ipsilateral graft re-lesion and contralateral rupture following ACL reconstruction is conflicting.

Research frontiers

Since the introduction of the anterior cruciate anatomic reconstruction concept, the pursuit of the ideal graft positioning has led to the so-called "anatomic"

single-bundle reconstruction techniques. However, despite the plethora of anatomical and biomechanical studies suggesting incorrect graft positioning and inadequate knee rotational stability with transtibial (TT) femoral drilling technique, there is no convincing evidence that anatomic reconstruction leads to better clinical outcomes. Moreover, recent data suggests that the anteromedial (AM) technique for femoral drilling could lead to an increased risk of graft re-rupture.

Innovations and breakthroughs

This is the first analysis focusing the influence of femoral drilling technique over the graft failure and contralateral ACL lesion after ACL reconstruction. The authors found that ACL reconstruction by TT technique led to lower incidence of graft re-injury than contralateral ACL lesion. There was no difference between the chance of re-injury after AM technique and the chance of contralateral ACL lesion (native ligament) with any technique.

Applications

The increased risk of graft re-rupture observed in anatomical reconstructions could in fact represent an approximation to the "normal knee" ACL rupture risk, that could be represented by the risk of contralateral knee lesion. The lower graft failure rates found in TT reconstruction could be because the neoligament simply is not submitted to the physiological forces that occur in the "normal" knee. With that in mind, one could speculate that it is not the AM technique that presents a higher failure rate, but it is the TT graft failure rate that is lower than it should be, probably due to incorrect femoral positioning.

Terminology

Transtibial technique refers to an anterior cruciate ligament reconstruction technique in which the femoral tunnel is created with a drill inserted through a tibial tunnel previously drilled. It leads to a non-anatomic positioning of the femoral tunnel, searching for an isometric position of the femoral tunnel. In the anteromedial portal technique, the femoral tunnel is created with a drill inserted through the arthroscopic anteromedial portal, which makes it possible to create the femoral tunnel in its anatomic position.

Peer-review

This is a very nice paper, it is well written with very interesting results and conclusions.

REFERENCES

- 1 **Rayan F**, Nanjayan SK, Quah C, Ramoutar D, Konan S, Haddad FS. Review of evolution of tunnel position in anterior cruciate ligament reconstruction. *World J Orthop* 2015; **6**: 252-262 [PMID: 25793165 DOI: 10.5312/wjo.v6.i2.252]
- 2 **Robin BN**, Jani SS, Marvil SC, Reid JB, Schillhammer CK, Lubowitz JH. Advantages and Disadvantages of Transtibial, Anteromedial Portal, and Outside-In Femoral Tunnel Drilling in Single-Bundle Anterior Cruciate Ligament Reconstruction: A Systematic Review. *Arthroscopy* 2015; **31**: 1412-1417 [PMID: 25749530 DOI: 10.1016/j.arthro.2015.01.018]
- 3 **Ferretti M**, Ekdahl M, Shen W, Fu FH. Osseous landmarks of the femoral attachment of the anterior cruciate ligament: an anatomic study. *Arthroscopy* 2007; **23**: 1218-1225 [PMID: 17986410 DOI: 10.1016/j.arthro.2007.09.008]
- 4 **Steiner M**. Anatomic single-bundle ACL reconstruction. *Sports Med Arthrosc* 2009; **17**: 247-251 [PMID: 19910783 DOI: 10.1097/JSA.0b013e3181c0ccf8]
- 5 **Chalmers PN**, Mall NA, Cole BJ, Verma NN, Bush-Joseph CA, Bach BR Jr. Anteromedial versus transtibial tunnel drilling in anterior cruciate ligament reconstructions: a systematic review. *Arthroscopy* 2013; **29**: 1235-1242 [PMID: 23591382 DOI: 10.1016/j.arthro.2013.02.015]
- 6 **Arnold MP**, Duthon V, Neyret P, Hirschmann MT. Double incision iso-anatomical ACL reconstruction: the freedom to place the femoral tunnel within the anatomical attachment site without exception. *Int Orthop* 2013; **37**: 247-251 [PMID: 23093291 DOI: 10.1007/s00264-012-1681-8]

- 7 **Riboh JC**, Hasselblad V, Godin JA, Mather RC 3rd. Transtibial versus independent drilling techniques for anterior cruciate ligament reconstruction: a systematic review, meta-analysis, and meta-regression. *Am J Sports Med* 2013; **41**: 2693-2702 [PMID: 24130247 DOI: 10.1177/0363546513506979]
- 8 **Rahr-Wagner L**, Thillemann TM, Pedersen AB, Lind MC. Increased risk of revision after anteromedial compared with transtibial drilling of the femoral tunnel during primary anterior cruciate ligament reconstruction: results from the Danish Knee Ligament Reconstruction Register. *Arthroscopy* 2013; **29**: 98-105 [PMID: 23276417 DOI: 10.1016/j.arthro.2012.09.009]
- 9 **Wright RW**, Magnussen RA, Dunn WR, Spindler KP. Ipsilateral graft and contralateral ACL rupture at five years or more following ACL reconstruction: a systematic review. *J Bone Joint Surg Am* 2011; **93**: 1159-1165 [PMID: 21776554 DOI: 10.2106/JBJS.J.00898]
- 10 **Aune AK**, Holm I, Risberg MA, Jensen HK, Steen H. Four-strand hamstring tendon autograft compared with patellar tendon-bone autograft for anterior cruciate ligament reconstruction. A randomized study with two-year follow-up. *Am J Sports Med* 2001; **29**: 722-728 [PMID: 11734484 DOI: 10.1177/03635465010290060901]
- 11 **Shaieb MD**, Kan DM, Chang SK, Marumoto JM, Richardson AB. A prospective randomized comparison of patellar tendon versus semitendinosus and gracilis tendon autografts for anterior cruciate ligament reconstruction. *Am J Sports Med* 2002; **30**: 214-220 [PMID: 11912091 DOI: 10.1177/03635465020300021201]
- 12 **Drogset JO**, Grøntvedt T, Tegnander A. Endoscopic reconstruction of the anterior cruciate ligament using bone-patellar tendon-bone grafts fixed with bioabsorbable or metal interference screws: a prospective randomized study of the clinical outcome. *Am J Sports Med* 2005; **33**: 1160-1165 [PMID: 16000666 DOI: 10.1177/0363546504272264]
- 13 **Sajovic M**, Vengust V, Komadina R, Tavcar R, Skaza K. A prospective, randomized comparison of semitendinosus and gracilis tendon versus patellar tendon autografts for anterior cruciate ligament reconstruction: five-year follow-up. *Am J Sports Med* 2006; **34**: 1933-1940 [PMID: 16923826 DOI: 10.1177/0363546506290726]
- 14 **Keays SL**, Bullock-Saxton JE, Keays AC, Newcombe PA, Bullock MI. A 6-year follow-up of the effect of graft site on strength, stability, range of motion, function, and joint degeneration after anterior cruciate ligament reconstruction: patellar tendon versus semitendinosus and Gracilis tendon graft. *Am J Sports Med* 2007; **35**: 729-739 [PMID: 17322130 DOI: 10.1177/0363546506298277]
- 15 **Pinczewski LA**, Lyman J, Salmon LJ, Russell VJ, Roe J, Linklater J. A 10-year comparison of anterior cruciate ligament reconstructions with hamstring tendon and patellar tendon autograft: a controlled, prospective trial. *Am J Sports Med* 2007; **35**: 564-574 [PMID: 17261567 DOI: 10.1177/0363546506296042]
- 16 **Shelbourne KD**, Gray T, Haro M. Incidence of subsequent injury to either knee within 5 years after anterior cruciate ligament reconstruction with patellar tendon autograft. *Am J Sports Med* 2009; **37**: 246-251 [PMID: 19109531 DOI: 10.1177/0363546508325665]
- 17 **Holm I**, Oiestad BE, Risberg MA, Aune AK. No difference in knee function or prevalence of osteoarthritis after reconstruction of the anterior cruciate ligament with 4-strand hamstring autograft versus patellar tendon-bone autograft: a randomized study with 10-year follow-up. *Am J Sports Med* 2010; **38**: 448-454 [PMID: 20097928 DOI: 10.1177/0363546509350301]
- 18 **Webb JM**, Salmon LJ, Leclerc E, Pinczewski LA, Roe JP. Posterior tibial slope and further anterior cruciate ligament injuries in the anterior cruciate ligament-reconstructed patient. *Am J Sports Med* 2013; **41**: 2800-2804 [PMID: 24036571 DOI: 10.1177/0363546513503288]
- 19 **Barenus B**, Ponzer S, Shalabi A, Bujak R, Norlén L, Eriksson K. Increased risk of osteoarthritis after anterior cruciate ligament reconstruction: a 14-year follow-up study of a randomized controlled trial. *Am J Sports Med* 2014; **42**: 1049-1057 [PMID: 24644301 DOI: 10.1177/0363546514526139]
- 20 **Webster KE**, Feller JA, Leigh WB, Richmond AK. Younger patients are at increased risk for graft rupture and contralateral injury after anterior cruciate ligament reconstruction. *Am J Sports Med* 2014; **42**: 641-647 [PMID: 24451111 DOI: 10.1177/0363546513517540]
- 21 **Pettiti DB**. Meta-analysis, decision analysis, and cost-effectiveness analysis: methods for quantitative synthesis in medicine. New York: Oxford University Press, 2000
- 22 **Higgins JPT**, Green S, editors. Cochrane handbook for systematic reviews of interventions. West Sussex: John Wiley & Sons Ltd., 2008
- 23 **Sterne JA**, Sutton AJ, Ioannidis JP, Terrin N, Jones DR, Lau J, Carpenter J, Rücker G, Harbord RM, Schmid CH, Tetzlaff J, Deeks JJ, Peters J, Macaskill P, Schwarzer G, Duval S, Altman DG, Moher D, Higgins JP. Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ* 2011; **343**: d4002 [PMID: 21784880]
- 24 **Jackson D**, White IR, Riley RD. Quantifying the impact of between-study heterogeneity in multivariate meta-analyses. *Stat Med* 2012; **31**: 3805-3820 [PMID: 22763950 DOI: 10.1002/sim.5453]
- 25 **Drogset JO**, Grøntvedt T. Anterior cruciate ligament reconstruction with and without a ligament augmentation device : results at 8-Year follow-up. *Am J Sports Med* 2002; **30**: 851-856 [PMID: 12435652 DOI: 10.1177/03635465020300061601]
- 26 **Bourke HE**, Salmon LJ, Waller A, Patterson V, Pinczewski LA. Survival of the anterior cruciate ligament graft and the contralateral ACL at a minimum of 15 years. *Am J Sports Med* 2012; **40**: 1985-1992 [PMID: 22869626 DOI: 10.1177/0363546512454414]
- 27 **Kato Y**, Ingham SJ, Kramer S, Smolinski P, Saito A, Fu FH. Effect of tunnel position for anatomic single-bundle ACL reconstruction on knee biomechanics in a porcine model. *Knee Surg Sports Traumatol Arthrosc* 2010; **18**: 2-10 [PMID: 19784631 DOI: 10.1007/s00167-009-0916-8]
- 28 **Kato Y**, Maeyama A, Lertwanich P, Wang JH, Ingham SJ, Kramer S, Martins CQ, Smolinski P, Fu FH. Biomechanical comparison of different graft positions for single-bundle anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2013; **21**: 816-823 [PMID: 22419266 DOI: 10.1007/s00167-012-1951-4]
- 29 **Araujo PH**, Asai S, Pinto M, Protta T, Middleton K, Linde-Rosen M, Irrgang J, Smolinski P, Fu FH. ACL Graft Position Affects in Situ Graft Force Following ACL Reconstruction. *J Bone Joint Surg Am* 2015; **97**: 1767-1773 [PMID: 26537164 DOI: 10.2106/JBJS.N.00539]
- 30 **van Eck CF**, Schreiber VM, Mejia HA, Samuelsson K, van Dijk CN, Karlsson J, Fu FH. "Anatomic" anterior cruciate ligament reconstruction: a systematic review of surgical techniques and reporting of surgical data. *Arthroscopy* 2010; **26**: S2-12 [PMID: 20810090 DOI: 10.1016/j.arthro.2010.03.005]
- 31 **van Eck CF**, Schkrohowsky JG, Working ZM, Irrgang JJ, Fu FH. Prospective analysis of failure rate and predictors of failure after anatomic anterior cruciate ligament reconstruction with allograft. *Am J Sports Med* 2012; **40**: 800-807 [PMID: 22238055 DOI: 10.1177/0363546511432545]

P- Reviewer: Fenichel I, Ohishi T **S- Editor:** Song XX **L- Editor:** A
E- Editor: Lu YJ





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

