

1 **Name of Journal:** *World Journal of Orthopedics*

2 **Manuscript NO:** 32876

3 **Manuscript Type:** Original Article

4

5 *Retrospective Study*

6 **Two-stage surgical treatment for septic non-union of the forearm: Retrospective review**

7 **of 18 patients with 6 years follow-up**

8

9 **Fabrizio Perna, Federico Pilla, Matteo Nanni, Lisa Berti, Giada Lullini, Francesco Traina,**

10 **Cesare Faldini**

11

12 **Abstract**

13 **Aim:**

14 To investigate the effectiveness of a two-stage surgical procedure for the treatment of

15 septic forearm non-union.

16 **Introduction:**

17 Septic non-unions are rare complications of forearm fractures. When they occur, they

Match Overview

1

Crossref 95 wordsFaldini, Cesare, Francesco Traina, Fabrizio Perna, Raffaele
Borghi, Matteo Nanni, and Mohammadreza Chehrassan. "Sur

2%

[All](#)[Images](#)[Videos](#)[News](#)[Shopping](#)[More](#)[Settings](#)[Tools](#)

About 12,200 results (1.05 seconds)

Did you mean: Two-stage surgical treatment for septic **nonunion** of the forearm: retrospective review of 18 patients with 6 years follow-up.

[Application of the Induced Membrane Technique for Forearm Bo ...](#)
[journals.lww.com](#) › [Home](#) › [March 2016 - Volume 31 - Issue 1](#)

The surgical treatment of forearm fracture nonunions remains a therapeutic challenge ... In patients with infected nonunion or osteomyelitis, this 2-stage technique Soucacos et al retrospectively reviewed 18 patients affected by large skeletal ... graft and he showed bone union in 15/16 patients with an average follow-up of ...

[The use of recombinant morphogenic protein-2 \(rhBMP-2\) in children ...](#)
<https://www.ncbi.nlm.nih.gov> › [NCBI](#) › [Literature](#) › [PubMed Central \(PMC\)](#)

by MC Papanna - 2016 - [Related articles](#)

Mar 16, 2016 - We undertook a retrospective review of all the patients who received ... Nineteen patients (21 surgical procedures) received rhBMP-2 as a part ... 18 years, had autologous bone graft in addition to rhBMP-2 or had a spinal fusion procedure. application of rhBMP-2 to tibial non-union sites at different stages.

[Proximal ulna comminuted fractures: Fixation using a double-plating ...](#)
www.sciencedirect.com/science/article/pii/S1877056810001532 ▼

by S Rochet - 2010 - [Cited by 11](#) - [Related articles](#)

Aug 30, 2010 - Retrospective study. The objective of surgery is first to obtain bone union, but also to restore a ... Only 18 patients were seen again (six were lost to follow-up) by a ... lateral X-ray of the operated elbow and an AP image of the forearm. fractures required non-surgical orthopaedic treatment (two cases), ...

[\[PDF\] Nonunion of Fractures of the Ulna and Radius Diaphyses: Clinical a...](#)
www.morthoj.org/2016/v10n2/ulna-and-radius-diaphyses.pdf ▼

complication of forearm fractures, accounting for 2% to 10% of all forearm ... the functional and radiological results of surgical treatment of diaphyseal aseptic nonunion of the radius and ulna, with ... retrospectively reviewed, the average age was 35 years with The patients were

找到约 23,600 条结果 (用时 0.83 秒)

[PDF] Progarimme Book - HKSSH

www.hkssh.org/2017/hkssh_30th_annual_congress_programme_book.pdf ▾ 翻译此页

2017年3月26日 - This year, five surgeons from the Russia Hand Surgery Society 17:00 - 18:00

Outcome for Arthroscopic Treatment of Septic Arthritis of the Wrist ... Reconstruction- a Novel Treatment with Long Term Follow up of 7.9 years defect (2), perilunate dislocation (5) forearm malunion, nonunion, bone defect.

[PDF] Nonunion of Fractures of the Ulna and Radius Diaphyses: Clinical an...

www.morthoj.org/2016/v10n2/ulna-and-radius-diaphyses.pdf ▾ 翻译此页

complication of forearm fractures, accounting for 2% to 10% of all forearm ... the functional and radiological results of surgical treatment of diaphyseal aseptic nonunion of the radius and ulna, with ... retrospectively reviewed, the average age was 35 years with The patients were followed-up to assess pain on analogue.

Application of the Induced Membrane Technique for Forearm Bo ...

journals.lww.com > Home > March 2016 - Volume 31 - Issue 1 - 翻译此页

The surgical treatment of forearm fracture nonunions remains a therapeutic challenge ... In patients with infected nonunion or osteomyelitis, this 2-stage technique Soucacos et al retrospectively reviewed 18 patients affected by large skeletal ... graft and he showed bone union in 15/16 patients with an average follow-up of ...

Treatment of Infected Forearm Nonunions With Large Complete ...

[https://www.jhandsurg.org/article/S0363-5023\(16\)30197-6/fulltext](https://www.jhandsurg.org/article/S0363-5023(16)30197-6/fulltext) - 翻译此页

2016年7月9日 - At a median of 42 months follow-up, 7 patients who had an average In the second stage, 6 to 8 weeks later, the spacer is removed, but the Two patients underwent 3 additional surgical procedures unrelated to ... a retrospective review of 35 patients with forearm nonunion, 11 of which were infected.

[全部](#)[图片](#)[视频](#)[新闻](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 117,000 条结果 (用时 0.80 秒)

您是不是要找: **Two-stage surgical treatment for septic *nonunion* of the forearm**

Surgical treatment options for septic non-union of the tibia: two staged ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5283947/> - 翻译此页

作者: R Kawakami - 2016 - 相关文章

For the **treatment of** extensive bone defects resulting from **septic non-union** of the and bone/soft tissue reconstruction in one **stage** or with a **two-staged operation**. ... Soutar et al. first reported a flow-through flap using a radial **forearm** flap, ...

Infection and Local Treatment in Orthopedic Surgery

<https://books.google.com.hk/books?isbn=3540479996> - 翻译此页

E. Meani, C. Romanò, L. Crosby - 2007 - Medical

Six patients had a **septic non-union** of the **forearm**, five a **septic non-union** of ... A **two-stage technique** was always performed and AMB was essentially used to ...

[PDF] **Nonunion of Fractures of the Ulna and Radius Diaphyses: Clinical an...**

www.morthoj.org/2016/v10n2/ulna-and-radius-diaphyses.pdf ▼ 翻译此页

complication of **forearm** fractures, accounting for **2%** to **10%** of all **forearm** ... the functional and radiological results of **surgical treatment** ... aseptic **nonunion** of the **forearm** fractures remains significant ... Exclusion criteria were **septic** nonunions and ... and three cases of open fracture (**stage I** in one case and **stage. II** in **two** ...

[PDF] **Download Scientific Program Accepted e-posters - 3rd World Ortho ...**

www.illrs-asami-br2017.org/docs/accepted-e-posters-final-24032017.pdf ▼ 翻译此页

#205 **Two-stage** reconstruction with external fixation of a bone defect after open distal shaft ... #67

Unusual **management of septic** pseudarthrosis of the humerus with length ... **Treatment of** radius infected