

Name of Journal: *World Journal of Clinical Oncology*

Manuscript NO: 32964

Manuscript Type: Review

Magnetic resonance imaging for prostate cancer before radical and salvage radiotherapy: What radiation oncologists need to know

Felipe Couñago, Gemma Sancho, Violeta Catalá, Diana Hernández, Manuel Recio, Sara Montemuiño, Jhonathan Alejandro Hernández, Antonio Maldonado, Elia del Cerro

### Abstract

External beam radiotherapy (EBRT) is one of the principal curative treatments for patients with prostate cancer (PCa). Risk group classification is based on prostate-specific antigen

### Match Overview

11	Crossref 13 words Functional Imaging in Oncology, 2014.	<1%
12	Crossref 13 words S D Heenan. "Magnetic resonance imaging in prostate cancer", Prostate Cancer and Prostatic Diseases	<1%
13	Internet 12 words crawled on 29-Jan-2016 <a href="http://aura.abdn.ac.uk">aura.abdn.ac.uk</a>	<1%
14	Internet 11 words crawled on 10-Oct-2012 <a href="http://www.tumoronline.it">www.tumoronline.it</a>	<1%
15	Crossref 10 words Medical Radiology, 2014.	<1%
16	Crossref 10 words Abdominal Imaging, 2013.	<1%
17	Internet 10 words crawled on 09-Oct-2015 <a href="http://uroweb.org">uroweb.org</a>	<1%
18	Crossref 10 words Interstitial Prostate Brachytherapy, 2013.	<1%

[All](#)[News](#)[Images](#)[Videos](#)[Shopping](#)[More](#)[Settings](#)[Tools](#)

About 150,000 results (0.45 seconds)

### Salvage radiotherapy following biochemical relapse after radical ...

<https://www.ncbi.nlm.nih.gov> › [NCBI](#) › [Literature](#) › [PubMed Central \(PMC\)](#)

by M Sia - 2008 - [Cited by 14](#) - [Related articles](#)

See commentary "Clarifying the role of **salvage radiotherapy**" on page 508. ... prostate cancer after radical prostatectomy undergo salvage radio-therapy — the ... after radical prostatectomy, the Genito-Urinary **Radiation Oncologists** of Canada .... in 41 patients with biochemical recurrence after surgery using endorectal MRI.

### Salvage Radiotherapy for Patients with PSA Relapse Following ... - NCBI

<https://www.ncbi.nlm.nih.gov> › [NCBI](#) › [Literature](#) › [PubMed Central \(PMC\)](#)

by R Choo - 2010 - [Cited by 18](#) - [Related articles](#)

31 Mar 2010 - Keywords: **Salvage radiotherapy**, PSA relapse, Post radical prostatectomy ... The recurrence of prostate cancer following RP is invariably .... **Magnetic resonance imaging (MRI)** has gained increasing favor in ..... respectively, using the **Radiation Therapy Oncology Group (RTOG)** toxicity scoring system (87).

### Predicting the Outcome of Salvage Radiation Therapy for Recurrent ...

<https://www.ncbi.nlm.nih.gov> › [NCBI](#) › [Literature](#) › [PubMed Central \(PMC\)](#)

by AJ Stephenson - 2007 - [Cited by 620](#) - [Related articles](#)

**Salvage radiation therapy (SRT)** may eradicate locally recurrent cancer, but studies to ... Significant variables in the model were PSA level before SRT ( $P < .001$ ), ... Nearly half of patients with recurrent prostate cancer after radical ..... such as endorectal coil **magnetic resonance imaging** may improve the efficacy of SRT.

### Optimal timing of salvage radiotherapy for ... - Radiation Oncology

<https://ro-journal.biomedcentral.com/articles/10.1186/s13014-016-0671-1> ▼

by S Taguchi - 2016 - [Related articles](#)



Magnetic resonance imaging for prostate cancer before radical and salve



All

News

Images

Videos

Shopping

More

Settings

Tools

About 197,000 results (0.89 seconds)

### Evaluation of tumor recurrences after radical prostatectomy using 18F ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5142428/>

by F Couñago - 2016 - [Related articles](#)

7 Dec 2016 - Multiparametric **MRI** and 18F-CH PET/CT yield similar detection rates for LR, ... After **radical prostatectomy** (RP), 20–50% of patients with **prostate cancer** (PCa) will ... to **determine** potential predictors of positive findings on the imaging studies. .... 11C-Choline PET/CT scan **before salvage radiation therapy**?

### Pelvic MRI findings in relapsed prostate cancer after radical ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4690270/>

by D Hernandez - 2015 - [Cited by 3](#) - [Related articles](#)

24 Dec 2015 - Pelvic **MRI** findings in relapsed **prostate cancer** after **radical prostatectomy** ... patients **have** recurrence in the ten years after **salvage radiotherapy** [3]. ... an mp-MRI to investigate the site of recurrence **before** planning the radiation treatment. ... of **radiation oncology** at the Hospital de la Santa Creu i Sant Pau.

### Predicting the Outcome of Salvage Radiation Therapy for Recurrent ...

[https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMed Central \(PMC\)](https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral(PMC))

by AJ Stephenson - 2007 - [Cited by 644](#) - [Related articles](#)

**Salvage radiation therapy** (SRT) may eradicate locally recurrent cancer, but ... Significant variables in the model were PSA level **before** SRT ( $P < .001$ ), ... Nearly half of patients with recurrent **prostate cancer** after **radical prostatectomy** **have** a ..... as endorectal coil **magnetic resonance imaging** may improve the efficacy of ...

### What Medical, Urologic, and Radiation Oncologists Want from ...

[jnm.snmjournals.org/content/57/Supplement\\_3/6S.full](http://jnm.snmjournals.org/content/57/Supplement_3/6S.full)

by LK Ballas - 2016 - [Cited by 1](#) - [Related articles](#)

1 Oct 2016 - As molecular imaging better delineates the state of **prostate cancer**, clinical .... **MRI** alone and PET alone in 53 patients undergoing **radical prostatectomy**. .... with the original treatment

Magnetic resonance imaging for prostate cancer before radical and salvage radi

全部 图片 视频 新闻 地图 图书

找到约 196,000 条结果

时间不限

过去 1 小时内  
过去 24 小时内  
过去 1 周内  
过去 1 个月内  
过去 1 年内

所有结果

精确匹配

[Evaluation of tumor recurrences after radical prostatectomy using ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5142428/>

7 Dec 2016 ... The median prostate-specific antigen (PSA) value pre-MRI/PET-CT was 0.9 ... After radical prostatectomy (RP), 20–50% of patients with prostate cancer (PCa) will ... to determine potential predictors of positive findings on the imaging studies . . . . . <sup>11</sup>C-Choline PET/CT scan before salvage radiation therapy?

[Pelvic MRI findings in relapsed prostate cancer after radical ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4690270/>

24 Dec 2015 ... Pelvic MRI findings in relapsed prostate cancer after radical prostatectomy ... patients have recurrence in the ten years after salvage radiotherapy [3]. ... an mp- MRI to investigate the site of recurrence before planning the radiation treatment. ... of radiation oncology at the Hospital de la Santa Creu i Sant Pau.

[Treatment of recurrent prostate cancer following radical ... - NCBI](#)

[www.ncbi.nlm.nih.gov/pubmed/23069920](http://www.ncbi.nlm.nih.gov/pubmed/23069920)

Recurrence of prostate cancer after radical prostatectomy is a common event. Salvage radiation therapy (RT) is the mainstay of treatment in cases with ... of MRI and choline-PET/CT in the salvage setting from the radiation-oncologist point of ...

[What Medical, Urologic, and Radiation Oncologists Want from ...](#)

[jnm.snmjournals.org/content/57/Supplement\\_3/6S.full](http://jnm.snmjournals.org/content/57/Supplement_3/6S.full)

1 Oct 2016 ... As molecular imaging better delineates the state of prostate cancer, clinical ... MRI alone and PET alone in 53 patients undergoing radical prostatectomy. .... with the original treatment to determine any disease control or survival advantage . . . . . <sup>11</sup>C-choline PET/CT scan before salvage radiation therapy?

[Clinical Radiation Oncology - Google 图书结果](#)

<https://books.google.co.uk/books?isbn=0323240984>