

World Journal of *Gastroenterology*

World J Gastroenterol 2017 July 7; 23(25): 4473-4660



**EDITORIAL**

- 4473** Challenges of modern day transition care in inflammatory bowel disease: From inflammatory bowel disease to biosimilars

Hakizimana A, Ahmed I, Russell R, Wright M, Afzal NA

REVIEW

- 4480** Expression and role of nuclear receptor coregulators in colorectal cancer

Triki M, Lapierre M, Cavailles V, Mokdad-Gargouri R

- 4491** Eubiotic properties of rifaximin: Disruption of the traditional concepts in gut microbiota modulation

Ponziani FR, Zocco MA, D'Aversa F, Pompili M, Gasbarrini A

MINIREVIEWS

- 4500** Localization and role of metabotropic glutamate receptors subtype 5 in the gastrointestinal tract

Ferrigno A, Berardo C, Di Pasqua LG, Siciliano V, Richelmi P, Vairetti M

ORIGINAL ARTICLE**Basic Study**

- 4508** Treatment with dimethyl fumarate ameliorates liver ischemia/reperfusion injury

Takasu C, Vaziri ND, Li S, Robles L, Vo K, Takasu M, Pham C, Farzaneh SH, Shimada M, Stamos MJ, Ichii H

- 4517** Enhanced electrogastrography: A realistic way to salvage a promise that was never kept?

Poscente MD, Mintchev MP

- 4529** Glutamine prevents oxidative stress in a model of portal hypertension

Zabot GP, Carvalhal GF, Marroni NP, Licks F, Hartmann RM, da Silva VD, Fillmann HS

- 4538** Hepatitis C virus NS5A region mutation in chronic hepatitis C genotype 1 patients who are non-responders to two or more treatments and its relationship with response to a new treatment

Muñoz de Rueda P, Fuentes Rodríguez JM, Quiles Pérez R, Gila Medina A, Martín Álvarez AB, Casado Ruiz J, Ruiz Extremera Á, Salmerón J

- 4548** Distinct gut microbiota profiles in patients with primary sclerosing cholangitis and ulcerative colitis

Bajer L, Kverka M, Kostovcik M, Macinga P, Dvorak J, Stehlikova Z, Brezina J, Wohl P, Spicak J, Drastich P

- 4559 Anti-inflammatory and anti-apoptotic effects of rosuvastatin by regulation of oxidative stress in a dextran sulfate sodium-induced colitis model

Shin SK, Cho JH, Kim EJ, Kim EK, Park DK, Kwon KA, Chung JW, Kim KO, Kim YJ

- 4569 miR-29a promotes hepatitis B virus replication and expression by targeting SMARCE1 in hepatoma carcinoma

Wu HJ, Zhuo Y, Zhou YC, Wang XW, Wang YP, Si CY, Wang XH

Retrospective Cohort Study

- 4579 Management and outcome of hepatocellular adenoma with massive bleeding at presentation

Klomphehouwer AJ, de Man RA, Thomeer MGJ, Ijzermans JNM

Retrospective Study

- 4587 Chronic hepatitis B, nonalcoholic steatohepatitis and physical fitness of military males: CHIEF study

Chen YJ, Chen KW, Shih YL, Su FY, Lin YP, Meng FC, Lin F, Yu YS, Han CL, Wang CH, Lin JW, Hsieh TY, Li YH, Lin GM

- 4595 Comparison of short- and long-term outcomes of laparoscopic vs open resection for gastric gastrointestinal stromal tumors

Ye X, Kang WM, Yu JC, Ma ZQ, Xue ZG

Observational Study

- 4604 Functional lipidomics in patients on home parenteral nutrition: Effect of lipid emulsions

Pironi L, Guidetti M, Verrastro O, Iacona C, Agostini F, Pazzeschi C, Sasdelli AS, Melchiorre M, Ferreri C

- 4615 Cryptogenic multifocal ulcerous stenosing enteritis: Radiologic features and clinical behavior

Hwang J, Kim JS, Kim AY, Lim JS, Kim SH, Kim MJ, Kim MS, Song KD, Woo JY

- 4624 Partners of patients with ulcerative colitis exhibit a biologically relevant dysbiosis in fecal microbial metacommunities

Chen GL, Zhang Y, Wang WY, Ji XL, Meng F, Xu PS, Yang NM, Bo XC

Randomized Controlled Trial

- 4632 Long-term irritable bowel syndrome symptom control with reintroduction of selected FODMAPs

Harvie RM, Chisholm AW, Bisanz JE, Burton JP, Herbison P, Schultz K, Schultz M

EVIDENCE-BASED MEDICINE

- 4644 Anti-apoptotic effect of banhasasim-tang on chronic acid reflux esophagitis

Shin MR, An HJ, Seo BI, Roh SS

SYSTEMATIC REVIEWS

- 4654** Nosocomial spontaneous bacterial peritonitis antibiotic treatment in the era of multi-drug resistance pathogens: A systematic review

Fiore M, Maraolo AE, Gentile I, Borgia G, Leone S, Sansone P, Passavanti MB, Aurilio C, Pace MC

Contents

World Journal of Gastroenterology
Volume 23 Number 25 July 7, 2017

ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Ludovico Abenavoli, MD, MSc, PhD, Associate Professor, Health Sciences, University Magna Graecia, Campus Germaneto, Viale Europa, 88100 Catanzaro, Italy

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7th, 14th, 21st, and 28th each month. The *WJG* Editorial Board consists of 1375 experts in gastroenterology and hepatology from 68 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

World Journal of Gastroenterology (*WJG*) is now indexed in Current Contents[®]/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch[®]), Journal Citation Reports[®], Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2017 edition of Journal Citation Reports[®] cites the 2016 impact factor for *WJG* as 3.365 (5-year impact factor: 3.176), ranking *WJG* as 29th among 79 journals in gastroenterology and hepatology (quartile in category Q2).

FLYLEAF

I-IX Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Fen-Fen Zhang*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Ze-Mao Gong*
Proofing Editorial Office Director: *Jin-Lei Wang*

NAME OF JOURNAL
World Journal of Gastroenterology

ISSN
ISSN 1007-9327 (print)
ISSN 2219-2840 (online)

LAUNCH DATE
October 1, 1995

FREQUENCY
Weekly

EDITORS-IN-CHIEF
Damian Garcia-Olmo, MD, PhD, Doctor, Professor, Surgeon, Department of Surgery, Universidad Autonoma de Madrid; Department of General Surgery, Fundacion Jimenez Diaz University Hospital, Madrid 28040, Spain

Stephen C Strom, PhD, Professor, Department of Laboratory Medicine, Division of Pathology, Karolinska Institutet, Stockholm 141-86, Sweden

Andrzej S Tarnawski, MD, PhD, DSc (Med), Professor of Medicine, Chief Gastroenterology, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach,

CA 90822, United States

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE
Jin-Lei Wang, Director
Yuan Qi, Vice Director
Ze-Mao Gong, Vice Director
World Journal of Gastroenterology
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>

<http://www.wjgnet.com>

PUBLICATION DATE
July 7, 2017

COPYRIGHT
© 2017 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.f6publishing.com>

Observational Study

Partners of patients with ulcerative colitis exhibit a biologically relevant dysbiosis in fecal microbial metacommunities

Guang-Lan Chen, Ye Zhang, Wang-Yue Wang, Xue-Liang Ji, Fei Meng, Pei-Song Xu, Ning-Min Yang, Fu-Qiang Ye, Xiao-Chen Bo

Guang-Lan Chen, Ye Zhang, Wang-Yue Wang, Xue-Liang Ji, Department of Gastroenterology, Lishui People's Hospital, Lishui 323000, Zhejiang Province, China

Fei Meng, Pei-Song Xu, Ning-Min Yang, Department of Research Service, Zhiyuan Medical Inspection Institute Co., LTD, Hangzhou 310030, Zhejiang Province, China

Fu-Qiang Ye, Xiao-Chen Bo, Department of Radiation Toxicology and Oncology, Beijing Institute of Radiation Medicine, People's Republic of China, Beijing 100089, China

Author contributions: Chen GL, Yang NM and Bo XC conceived the study; Zhang Y and Wang WY collected samples; Meng F and Xu PS performed the 16S rRNA sequencing; Ye FQ analyzed the data; Chen GL, Meng F and Xu PS wrote the paper; all authors contributed to this manuscript.

Supported by Lishui Science and Technology Bureau Research Fund, No. 2013JYZB43; and Medical and Health Science and Technology Plan Project of Zhejiang Province, No. 2015KYB371.

Institutional review board statement: This study was approved by the Ethics Committee of Lishui People's Hospital.

Informed consent statement: All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

Conflict-of-interest statement: To the best of our knowledge, no conflicts of interest exist.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on

different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Correspondence to: Dr. Xiao-Chen Bo, Department of Radiation Toxicology and Oncology, Beijing Institute of Radiation Medicine, People's Republic of China, No. 27, Taiping Road, Haidian District, Beijing 100089, China. boxc@bmi.ac.cn
Telephone: +86-10-66931207
Fax: +86-10-66931207

Received: February 15, 2017

Peer-review started: February 16, 2017

First decision: April 7, 2017

Revised: April 10, 2017

Accepted: May 19, 2017

Article in press: May 19, 2017

Published online: July 7, 2017

Abstract

AIM

To investigate alterations in the fecal microbiome using 16S rRNA amplicon sequencing in couples in the same cohabitation environment.

METHODS

Fecal samples were collected from eight ulcerative colitis (UC) patients and their healthy partners at Lishui People's Hospital, Zhejiang Province, China. DNA was extracted and the variable regions V3 and V4 of the 16S rRNA genes were PCR amplified using a two-step protocol. Clear reads were clustered into operational taxonomic units (OTUs) at the 97% sequence similarity level using UCLUST v1.2.22. The Wilcoxon rank-sum

test (R v3.1.2) was used to compare inter-individual differences. Differences with a P value < 0.05 were considered statistically significant.

RESULTS

Fecal microbial communities were more similar among UC patients than their healthy partners ($P = 0.024$). UC individuals had a lower relative abundance of bacteria belonging to the *Firmicutes*, especially *Blautia*, *Clostridium*, *Coprococcus* and *Roseburia* ($P < 0.05$). Microbiota dysbiosis was detected in UC patients and their healthy partners. Relevant genera included *Akkermansiam*, *Bacteroides*, *Escherichia*, *Lactobacillales*, *Klebsiella* and *Parabacteroides*. The enriched pathways in fecal samples of UC patients were related to lipid and nucleotide metabolism. Additionally, the pathways involved in membrane transport and metabolism of cofactors and vitamins were more abundant in the healthy partners.

CONCLUSION

Our results suggested that the microbial composition might be affected in healthy partners cohabiting with UC patients, especially in terms of microbiota dysbiosis.

Key words: Ulcerative colitis; Patients; Healthy partner; Fecal microbial communities; Microbiota dysbiosis

© The Author(s) 2017. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: To identify the influence on the gut microbial community between ulcerative colitis (UC) patients and their healthy partners, we investigated the gut bacterial community using 16S rRNA amplicon sequencing. The results showed that fecal microbial communities were more similar in UC patients, which had a lower relative abundance of Firmicutes bacteria. Microbiota dysbiosis was also founded in healthy partners. The pathways involved in lipid and nucleotide metabolism were more abundant in the UC patients. The membrane transport and metabolism of cofactors and vitamins pathways were significantly enriched in the healthy partners. Microbial composition might be affected in healthy partners cohabiting with UC patients.

Chen GL, Zhang Y, Wang WY, Ji XL, Meng F, Xu PS, Yang NM, Bo XC. Partners of patients with ulcerative colitis exhibit a biologically relevant dysbiosis in fecal microbial metacommunities. *World J Gastroenterol* 2017; 23(25): 4624-4631 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v23/i25/4624.htm> DOI: <http://dx.doi.org/10.3748/wjg.v23.i25.4624>

INTRODUCTION

Ulcerative colitis (UC), one of the main types of inflammatory bowel disease (IBD), has become

increasingly prevalent in developed countries over the past two decades^[1]. However, according to the latest Asian epidemiological investigation, the incidence of the IBD has expanded dramatically into developing countries with the increased westernization of lifestyles^[1,2].

UC is a multi-faceted disorder associated with a germline genetic background, an aberrant immune system response, and environmental factors^[3]. Recently, the gut microbial community has attracted substantial research attention, especially in terms of its influence in healthy and IBD patients^[4-6]. Some alterations in the microbial community are shared in Crohn's disease (CD) and UC patients relative to healthy people, including reduced gut microbiota diversity (particularly *Firmicutes*), the presence of non-commensals, and increased abundance of pathogenic Proteobacteria strains^[7,8]. However, some alterations in the bacterial community are specific to UC. For example, an increased presence of *Escherichia coli* and *Fusobacterium spp* group^[6], as well as a reduction in *Clostridium coccoides* group, has been reported in UC patients^[9].

Previous studies have demonstrated that environmental factors, including dietary age^[9], habits^[10], and obesity^[11], impact the composition of the gut microbiota. Consequently, we wondered whether the cohabitation environment can influence the microbial community. In 1994, an investigation of 10 couples showed that individuals with IBD symptoms before marriage influenced their partners, resulting in similar symptoms in couples^[12]. However, these results mainly focused on clinical symptoms and did not involve the gut microbial community.

Profiling the fecal microbiome using methods based on analysis of the 16S ribosomal RNA gene is less biased than cultivation-based approaches. In recent years, bacterial 16S rRNA amplicon sequencing, referred to as "16S rRNA gene sequencing", has been used widely for the metagenomic analysis of the environment, including analysis of the composition of human and animal gut and fecal microbiota^[13,14]. In particular, compared with bacterial culture, 16S rRNA gene sequencing has a huge advantage in identifying new pathogens or difficult to culture bacteria compared with culture-independent methods. Compared with other high-throughput sequencing platforms (pyrosequencing, Life Technologies platform), the MiSeq platform generates the highest base sequence accuracy with little limitation on the DNA input^[15,16]. Although the cohabitation environment has been studied using 16S rRNA gene sequencing, many studies on the gut microbial community of IBD patients focused on comparisons between twins and siblings^[11,17].

This study aimed to explore deeply the composition of the UC gut bacterial community using 16S rRNA sequencing to identify the influence on the gut microbial

community between couples. Additionally, the current study investigates the predominant fecal microbiota of UC patients compared with their healthy cohabiting partners.

MATERIALS AND METHODS

Patient selection

The diagnosis of UC was determined by endoscopic and pathological findings, and patients diagnosed with UC for longer than 3 mo were recruited from Lishui People's Hospital at Zhejiang Province. Healthy couples composing the control group, were recruited from a common living environment cohabiting with the UC patients. All participants in this study were divided into two groups ($n = 8$ each) according to their disease status. Group A consisted of UC patients, identified as UC1-UC8. Group B consisted of their healthy partners, termed HF1-HF8. The study was approved by the Ethics Committee of the hospital and all participants provided written informed consent upon enrolment.

Sample collection and DNA extraction

Fecal samples were collected from all participants and were immediately stored at -80°C until further processing. The total DNA was extracted from 200 mg of fecal sample using the QIAamp DNA Stool Mini Kit (Qiagen, Inc., Valencia, CA, United States), according to the manufacturer's instructions. The DNA concentration was quantified using a Qubit 2.0 Fluorometer and diluted appropriately based on the total DNA concentration.

16S amplicon-sequencing

The variable regions V3 and V4 of the bacterial 16S rRNA gene were amplified using a two-step PCR protocol. In brief, the first PCR was performed with universal primers (356F 5'-CCTACGGGNGGCWGCAG-3' and 803R 5'-GACTACHVGGGTATCTAATCC-3') and an attached overhang adapter (forward primer overhang adapter 5'-TCGTCCGCGAGCGTCAGATGTGTATAAGAGACAG-3' and reverse primer overhang adapter 5'-GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG-3'). The first PCR reaction mixture (20 μL) contained 10 ng of DNA template, 10 μL of 2 \times High-Fidelity PCR Master Mix with HF Buffer (New England Biolabs, Ipswich, MA, United States), each primer at 5 $\mu\text{mol/L}$, and reagent-grade water (Sigma Aldrich, St. Louis, MO, United States). The first PCR program for V3 and V4 comprised an initial denaturation of 95°C for 3 min; followed by 25 cycles of denaturation at 95°C for 30 s, annealing at 60°C for 30 s, extension at 72°C for 30 s; and a final extension at 72°C for 5 min. Then, the PCR product was cleaned using AMPure XP DNA purification beads (Beckman Coulter, Danvers, MA, United States), according to the manufacturer's instructions. The second PCR reaction mixture (25 μL) contained 2 μL of cleaned PCR products; 12.5 μL of

2 \times High-Fidelity PCR Master Mix and HF Buffer; P5/P7 primers, including adapters and sample barcodes (Illumina); and reagent-grade water. The second PCR program was similar to the first PCR program except the annealing temperature was at 50°C and the program comprised 10 cycles. Finally, the PCR product was purified using AMPure XP beads to remove primer dimers and was then quantified on an Agilent Bioanalyzer 2100 using High Sensitivity DNA chips (Agilent Technologies, Santa Clara, CA, United States). The library of each sample was pooled at an equimolar concentration, and sequencing was performed on the Illumina MiSeq platform to generate 2 \times 250 bp paired-end reads.

Taxonomic classification and pathway profiles of 16S rRNA gene sequencing data

We merged the paired-end reads using FLASH v1.2.11 and obtained approximately 460 bp V3-V4 16S sequences. Merged reads were processed with QIIME v1.8.0, which removed reads with N bases and trimmed reads with more than three consecutive low-quality bases ($Q < 20$). The reads that passed the high quality filters were aligned to the Greengenes Database (Aug, 2013 version), and USEARCH v6.1 was applied for chimera checking. Then, UCLUST v1.2.22 was used for operational taxonomic unit (OTU) clustering at the 97% sequence similarity level. Each OTU was classified according to an assignment of the taxonomic rank using the reference dataset from the Ribosomal Database Project (version 2.2). Phylogenetic Investigation of Communities by Reconstruction of Unobserved States (PICRUST) analysis was performed to generate Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway profiles as previously described^[18,19].

Statistical analysis

Correlations between the overlapping genera of the 16S rRNA gene sequencing were identified using Pearson's correlation in the R language. The Wilcoxon rank-sum test (R v3.1.2) was employed to detect interindividual differences. Under the condition of multiple comparisons, P values were corrected to control for the false-discovery rate. Differences with P value < 0.05 were considered statistically significant.

RESULTS

Study subjects and 16S rRNA sequencing

To measure the compositional and functional differences in the common living environment between the gut microbiota of the UC patients and healthy individuals, 16 fecal samples were collected from eight families, including eight UC patients and eight healthy control partners. All patients in this study were cohabiting with their partners. The demographics and characteristics of 8 UC patients and their partners are shown in Table 1.




Figure 1 Distributions of the unweighted unifrac distance within the ulcerative colitis patients and healthy partners groups. Orange and red boxplots denote the distributions of 16S rRNA gene sequencing of fecal samples in ulcerative colitis patients and healthy partners, respectively.

UC patients vs healthy couples		
Phylum	Genus	P value
Actinobacteria	Coriobacteriaceae (unclassified)↓	0.021
Bacteroidetes	Alistipes↓	0.016
Firmicutes	Anaerostipes↓	0.009
	Blautia↓	0.041
	Clostridium↓	0.007
	Coprococcus↓	0.020
	Lachnospira↓	0.033
	Megasphaera↓	0.042
	Roseburia↓	0.007
	Turicibacter↓	0.026
	Lactobacillales (unclassified)↑	0.032

Figure 2 Significant differences in the fecal microbiota composition between ulcerative colitis patients and their healthy partners. Comparisons for each sample were calculated using the Wilcoxon test. The down arrows indicate less abundant microbiota in ulcerative colitis (UC) patients, and the up arrow represents the more abundant microbiota in UC patients compared with their healthy partners.

16S rRNA amplicon-sequencing of fecal DNA samples was performed using next-generation sequencing (NGS) technology. Low quality reads and chimera sequences were filtered from the raw data, eventually producing an average of 47469 reads per sample. These reads corresponded to 1137 operational taxonomic units (OTUs) at the 97% sequence similarity level and identified 167 genera. There was no significant difference between the number of OTUs in UC patients and their partners (947 ± 298.4 for UC patients and 1327.1 ± 419.9 for their partners, $P = 0.071$).

Microbiota diversity and similarity analyses

We used three metrics to characterize the fecal microbiota diversity between UC patients and their partners. Microbial alpha diversity was not significantly different by Chao1 analysis ($P = 0.573$). Additionally, there were no significant differences observed by Shannon-Wiener ($P = 0.505$) or Simpson's indices

Table 1 Information and clinical characteristics of ulcerative colitis patients and their healthy partners n (%)

	UC patients ($n = 8$)	Healthy partners ($n = 8$)
mean age (SD)	53 (9.57)	42.25 (9.41)
Male	6 (75.0)	3 (37.5)
Age at diagnosis		
16-40 yr	3 (37.5)	
Above 40 yr	5 (62.5)	
Disease location		
Colonic	5 (62.5)	
Rectal	3 (37.5)	
Concomitant upper GI disease	4 (50.0)	
CRP		
Normal (0-10 mg/L)	5 (62.5)	
Abnormal	3 (37.5)	

GI: Gastrointestinal; CRP: C-reactive protein; UC: Ulcerative colitis.

($P = 0.574$). The distribution of fecal microbiota was determined using Bray Curtis similarities analysis. The entire microbiota of UC patients was not significantly divergent from those of their cohabiting partners ($P = 0.449$). However, using un-weighted analysis, the microbiota composition was more similar among UC patients than it was to that of their partners (Figure 1, $P = 0.024$). These results indicated that cohabitation with UC patients might influence the fecal microbiota composition of healthy people; however, higher dissimilarity of the microbiota was observed among the healthy partners.

Microbial composition

Although there was no significant difference in overall microbial diversity, the Wilcoxon test indicated that some bacteria were significantly different between UC patients and their partners. As shown in Figure 2, at the genus level, 10 genera were less abundant in the fecal microbiota of UC patients compared with that in their healthy partners, and only one bacterium was more abundant in the UC patients ($P < 0.05$). At the phylum level, nine significantly different bacteria species belonged to the *Firmicutes*, and the other bacteria were classified as *Actinobacteria* and *Bacteroidetes*.

In this study, the fecal microbiota in the UC patients showed a decrease in the *Firmicutes* bacteria, especially *Blautia*, *Clostridium*, *Coprococcus*, and *Roseburia*. *Blautia* was detected at 5.81% in the healthy partners, and this percentage was lower in the UC patients. *Clostridium* and *Coprococcus* accounted for 6.49% and 3.4% of the fecal microbiota of the healthy partners, whereas these genera were less abundant in the UC patients (0.96% and 0.45%, respectively). A similar trend was observed for *Roseburia*, which was present at 1.48% in the healthy partners and 0.15% in the UC patients. Other *Firmicutes* genera, such as *Anaerostipes*, *Lachnospira*, *Megasphaera*, and *Turicibacter*, were present at a




Figure 3 Relative abundance at the genus level, as shown by 16S rRNA gene sequencing, from ulcerative colitis patients and their partners showing a microbiota imbalance. Each column represents one fecal sample and different colors indicate different genera in the microbiota composition. Microbiota with a substantial proportional imbalance are listed. UC: UC patient; HF: Healthy partners.

very low abundance in the UC patients. Although they were not detected in all healthy partners, these rare species were detected at significantly different levels in the healthy partners compared with the UC patients. For example, *Anaerostipes* was detected in UC patients at a percentage of 0.006%, but at 0.68% in the healthy partners. Additionally, *Alistipes*, belonging to the *Bacteroidetes*, and an unclassified bacterium of *Actinobacteria*, did not appear in UC patients, but were detected in the healthy partners at low abundances of 0.14% and 0.23%, respectively.

Microbiota dysbiosis

An unclassified bacterium belonging to the *Lactobacillales* was detected in a UC patient (UC1) with a higher relative abundance than in their partner (Figure 2, $P = 0.032$). The proportion of *Lactobacillus* genus showed large dysbiosis in patient UC1, accounting for 81.8% (Figure 3). Microbiota dysbiosis was also observed in other UC patients and their partners. In patient UC2, the first and second most abundant genera were *Bacteroides* and *Parabacteroides*, which were present in percentages of 58.63% and 28.93%, respectively (Figure 3). *Bacteroides* was also the dominant genus, accounting for 70.24% in patient UC6. In addition, the *Escherichia* genus accounted for 44.39% in patient UC5. For the healthy partners, microbiota dysbiosis was also found in a few individuals. For example, in HF6, the relative abundance of the *Klebsiella* genus reached 63.3%. *Akkermansia*, belonging to the *Verrucomicrobia* phylum, was only detected in patient UC4 and his partner, accounting for 18.46% and 68.74%, respectively (Figure 3). Microbiota dysbiosis destroys the gut microbiota composition and influences its normal function, which might accelerate the occurrence of intestinal disease.

Microbial metabolic pathways

Combining the microbial composition with a genome database from KEGG (Kyoto Encyclopedia of Genes and Genomes), 328 pathways were identified. We continued our analysis using the Wilcoxon test, and 20 of 328 (6.09%) total metabolic pathways were differentially abundant at $Q < 0.05$ between the UC patients and their healthy partners. In the UC patients, we observed that 10 pathways were increased significantly, especially those concerning lipid metabolism and nucleotide metabolism. Another eight pathways, such as those involving membrane transport and metabolism of cofactors and vitamins, were more abundant in the healthy partners (Table 2).

The lipid metabolic pathway, including primary bile acid biosynthesis, secondary bile acid biosynthesis ($P = 0.019$), and linoleic acid ($P = 0.025$), was enriched significantly in the UC patients. The pathway involved in purine metabolism was also enriched in the UC patients ($P = 0.019$). In contrast, there was a decrease in porphyrin and chlorophyll metabolism in the UC patients ($P = 0.032$). In addition, the microbiota of the UC patients had fewer ABC transporters for membrane transport ($P = 0.041$).

DISCUSSION

Alteration in the composition of the gut microbiota and decreases in community diversity are associated with the pathogenesis of UC^[20]. However, our understanding of the interaction between couples cohabiting a shared environment in terms of their microbiota is poor. In this study, we analyzed the differences in the bacterial profiles and metabolic pathways between UC patients and their cohabiting partners. Marked microbiota dysbiosis and reduction of the diversity of *Firmicutes* were observed. Furthermore, to understand

Table 2 Microbial metabolism differentially abundant in ulcerative colitis patients and their healthy partners

Pathway	UC patients		Healthy partners		P value
	Mean abundance	Standard error	Mean abundance	Standard error	
Membrane transport					
ABC transporters ¹	3.0271	0.5695	3.7117	0.7130	0.04149
Signal transduction					
Phosphatidylinositol signaling system	0.1022	0.0155	0.0845	0.0182	0.018959
Signaling molecules and interaction					
G protein-coupled receptors ¹	1.76×10^{-5}	3.63×10^{-5}	6.54×10^{-5}	1.02×10^{-5}	0.02515
Infectious diseases					
Staphylococcus aureus infection	0.295	0.582	0.00545	0.00442	0.041492
Biosynthesis of other secondary					
Metabolites; isoflavonoid biosynthesis	8.97×10^{-6}	8.03×10^{-6}	9.75×10^{-7}	1.39×10^{-7}	0.003511
Novobiocin biosynthesis ¹	0.1161	0.010	0.1277	0.0122	0.02494
Carbohydrate metabolism					
C5-Branched dibasic acid metabolism ¹	0.287	0.0536	0.3214	0.0218	0.04149
Glycan biosynthesis and metabolism					
Glycosphingolipid biosynthesis	3.74×10^{-4}	3.95×10^{-4}	1.37×10^{-4}	1.91×10^{-4}	0.032476
Lipid metabolism					
Linoleic acid metabolism	0.0851	0.0505	0.0518	0.0197	0.024942
Primary bile acid biosynthesis	0.0416	0.0186	0.0257	0.0120	0.018959
Secondary bile acid biosynthesis	0.0414	0.0187	0.0255	0.0120	0.018959
Metabolism of cofactors and vitamins					
Porphyrin and chlorophyll II metabolism ¹	0.7225	0.1723	0.8785	0.1079	0.03248
Nucleotide metabolism					
Purine metabolism	2.2171	0.2176	2.0283	0.1561	0.018959
Xenobiotics biodegradation and metabolism					
Atrazine degradation ¹	0.0227	0.0252	0.0441	0.0329	0.03248
Ethylbenzene degradation	0.0769	0.0643	0.0429	0.00795	0.018959
Naphthalene degradation	0.1827	0.0823	0.1300	0.0261	0.018959
Nitrotoluene degradation ¹	0.0651	0.0346	0.1022	0.0198	0.02494
Styrene degradation ¹	0.0155	0.0131	0.0329	0.0231	0.02494

¹Indicated higher abundance in the healthy partners ($P < 0.05$). UC: Ulcerative colitis.

the functions of the different bacteria, we compared the differences in microbial metabolic pathways. Lipid metabolism and the biosynthesis of bile acids were significantly upregulated in the microbiota of the UC patients.

Unlike previous reports, we did not observe a significant difference in the OTU distribution associated with the disease state^[21]. Additionally, we did not identify significant differences between UC patients and their partners in terms of the microbial alpha diversity using the Chao1, Shannon, and Simpson's indices^[20,22]. In contrast, we confirmed a compositional similarity between UC patients and their partners. Although there was no direct evidence that gut microbiota could spread between IBD patients and normal individuals, the results of this study suggested that fecal microbiota likely influence each other during long-term cohabitation with UC patients. Remarkably, our results indicated that the fecal microbiota composition was more similar among UC patients than among healthy individuals (Figure 1). These findings further demonstrated that the gut microbiota composition, and alterations to it, plays a crucial role in the occurrence of UC.

Although the cause of UC has many uncertain factors, gut microbiota dysbiosis has been considered a major trigger of inflammation^[23]. Consistent with

previous studies^[22,24], we also found that the fecal microbiota in the UC patients in the present study had a decrease in *Firmicutes* bacteria, especially in *Blautia*, *Clostridium*, *Coprococcus*, and *Roseburia* bacteria (Figure 2). Although we realized that the gut microbial composition and its interaction with the host likely play an important role in IBD, the relationship between these has remained a mystery. Our research further confirmed that *Firmicutes* plays a crucial role in UC patients. The reduction in *Clostridium* likely decreased the utilization of short chain fatty acids and butyric acid salt in intestinal epithelial cells and induced inflammation. The depletion of *Actinobacteria* and *Bacteroidetes* is controversial, and we found that these bacteria were greatly depleted^[21,25,26].

Interestingly, comparison of the relative abundance at the genus level using 16S rRNA gene sequencing from UC patients and their partners demonstrated that the composition of the fecal microbiota was dominated by *Bacteroides* in patients UC2 and UC6. By contrast, other samples, including those from healthy partners, did not show this situation. Additionally, among the healthy partners, HF4 and HF6 were dominated by *Akkermansia* and *Klebsiella*, respectively. We speculated that this microbiota dysbiosis is likely a consequence of interaction in the same environment.

In agreement with previous studies, we observed

that lipid metabolism was remarkably increased^[27]. Previous studies have suggested that commensal bacteria might increase or decrease certain specific metabolic pathways to participate in competition for limited energy resources while living in the host intestinal environment. Davenport *et al.*^[27] posited that because of a lack of carbohydrates in the inflamed regions, such as in the case of mucin production dysfunction, gut commensal bacteria start metabolizing lipids and amino acids for necessary nutrients. This hypothesis was supported by the decrease in *Firmicutes*, which are unable to utilize amino acids for energy^[27,28].

Furthermore, one study found that patients with IBD may also suffer from the co-occurrence of primary biliary cirrhosis^[29]. In the present study, we observed a significant increase in both primary bile acid biosynthesis and secondary bile acid biosynthesis of lipids ($P = 0.019$) in the UC patients (Table 2). In healthy individuals, it is difficult to detect bile acid in the blood. However, when liver cells are damaged, bile acid biosynthesis is abnormal and its concentration increases. We speculated that the increase in bile acid biosynthesis in UC patients might be associated with frequent liver disease, such as chronic liver disease or autoimmune liver disease. Of course, liver cell damage was also related to drug toxicity produced by long term medication and immune deficiency in UC patients. This would lead to oxidative stress, and some bacteria need to maintain homeostasis under oxidative stress. Interestingly, we observed an increase in linoleic acid metabolism in the UC patients ($P = 0.025$). Linoleic acid is a type of unsaturated fatty acid that is associated with prostaglandin biosynthesis, which could participate in liver injury protection.

In conclusion, this study presents a comprehensive evaluation of the bacterial composition and the differences in the pathways of UC fecal microbiota. Although our results were similar to the results of previous studies, we also identified an increased prevalence of lipid metabolic pathways and bile acid biosynthesis. The gut microbiota of the UC patients and their partners likely influenced each other. Furthermore, we verified that microbiota dysbiosis is more likely a consequence, rather than a cause, of inflammation.

COMMENTS

Background

Ulcerative colitis (UC) is one of the main types of inflammatory bowel disease. Compared with healthy people, some alterations in the microbial community are shared in Crohn's disease and UC patients. However, some alterations in the bacterial community are specific to UC patients. The authors wondered whether cohabitation could influence the microbial community. Previous studies mainly focused on clinical symptoms and did not involve gut microbial community. Profiling the fecal microbiome using methods based on analysis of the 16S ribosomal RNA gene is less biased than cultivation-based approaches. Therefore, we investigated the gut bacterial community between UC patients and their healthy partners using 16S rRNA amplicon sequencing.

Research frontiers

Recently, the gut microbial community has attracted substantial attention, especially the influence in healthy and inflammatory bowel disease (IBD) patients. Some studies have demonstrated that environmental factors, including dietary age, habits, and obesity, impact the composition of the gut microbiota.

Innovations and breakthroughs

This study is the first to investigate the influence on the gut microbial community between UC patients and their cohabiting healthy partners. This study showed that microbial composition might affect healthy partners cohabiting with UC patients, especially in terms of microbiota dysbiosis.

Applications

In this study, the authors analyzed the differences in the bacterial profiles and metabolic pathways between UC patients and their cohabiting partners. The findings further demonstrated that the gut microbiota composition and alterations play a crucial role in the occurrence of UC. Furthermore, the gut microbiota of the UC patients and their partners likely influence each other.

Terminology

16S rRNA gene sequencing is a non-culture method based on high-throughput sequencing. At present, 16S rRNA gene sequencing has been utilized widely for metagenomic analysis of the environment, including analysis of the composition of the human and animal guts and fecal microbiota.

Peer-review

Gut dysbiosis is considered one of the factors inducing inflammation in chronic IBD; however, its role in the etiology of ulcerative colitis is controversial. There have been many studies on fecal microbiota in recent years. The present study used a very sensitive method to assess bacterial strains and compared the microbiota in patients with inflammatory bowel disease with that of their healthy partners. The study assessed some type of bacteria only found in patients with UC. These bacteria are difficult to detect using less sensitive methods.

REFERENCES

- 1 **Molodecky NA**, Soon IS, Rabi DM, Ghali WA, Ferris M, Chernoff G, Benchimol EI, Panaccione R, Ghosh S, Barkema HW, Kaplan GG. Increasing incidence and prevalence of the inflammatory bowel diseases with time, based on systematic review. *Gastroenterology* 2012; **142**: 46-54.e42; quiz e30 [PMID: 22001864 DOI: 10.1053/j.gastro.2011.10.001]
- 2 **Ng SC**. Epidemiology of inflammatory bowel disease: focus on Asia. *Best Pract Res Clin Gastroenterol* 2014; **28**: 363-372 [PMID: 24913377 DOI: 10.1016/j.bpg.2014.04.003]
- 3 **Flint HJ**, Scott KP, Louis P, Duncan SH. The role of the gut microbiota in nutrition and health. *Nat Rev Gastroenterol Hepatol* 2012; **9**: 577-589 [PMID: 22945443 DOI: 10.1038/nrgastro.2012.156]
- 4 **Sokol H**, Seksik P, Rigottier-Gois L, Lay C, Lepage P, Podglajen I, Marteau P, Doré J. Specificities of the fecal microbiota in inflammatory bowel disease. *Inflamm Bowel Dis* 2006; **12**: 106-111 [PMID: 16432374 DOI: 10.1097/01.MIB.0000200323.38139.c6]
- 5 **Sokol H**, Lepage P, Seksik P, Doré J, Marteau P. Temperature gradient gel electrophoresis of fecal 16S rRNA reveals active *Escherichia coli* in the microbiota of patients with ulcerative colitis. *J Clin Microbiol* 2006; **44**: 3172-3177 [PMID: 16954244 DOI: 10.1128/JCM.02600-05]
- 6 **Sokol H**, Lay C, Seksik P, Tannock GW. Analysis of bacterial bowel communities of IBD patients: what has it revealed? *Inflamm Bowel Dis* 2008; **14**: 858-867 [PMID: 18275077 DOI: 10.1002/ibd.20392]
- 7 **Kostic AD**, Xavier RJ, Gevers D. The microbiome in inflammatory bowel disease: current status and the future ahead. *Gastroenterology* 2014; **146**: 1489-1499 [PMID: 24560869 DOI: 10.1053/j.gastro.2014.02.009]
- 8 **Ohkusa T**, Sato N, Ogihara T, Morita K, Ogawa M, Okayasu I.

- Fusobacterium varium localized in the colonic mucosa of patients with ulcerative colitis stimulates species-specific antibody. *J Gastroenterol Hepatol* 2002; **17**: 849-853 [PMID: 12164960]
- 9 **Martín-de-Carpi J**, Rodríguez A, Ramos E, Jiménez S, Martínez-Gómez MJ, Medina E; SPIRIT-IBD Working Group of Sociedad Española de Gastroenterología, Hepatología y Nutrición Pediátrica. Increasing incidence of pediatric inflammatory bowel disease in Spain (1996-2009): the SPIRIT Registry. *Inflamm Bowel Dis* 2013; **19**: 73-80 [PMID: 22535573 DOI: 10.1002/ibd.22980]
 - 10 **Wu GD**, Chen J, Hoffmann C, Bittinger K, Chen YY, Keilbaugh SA, Bewtra M, Knights D, Walters WA, Knight R, Sinha R, Gilroy E, Gupta K, Baldassano R, Nessel L, Li H, Bushman FD, Lewis JD. Linking long-term dietary patterns with gut microbial enterotypes. *Science* 2011; **334**: 105-108 [PMID: 21885731 DOI: 10.1126/science.1208344]
 - 11 **Turnbaugh PJ**, Hamady M, Yatsunenko T, Cantarel BL, Duncan A, Ley RE, Sogin ML, Jones WJ, Roe BA, Affourtit JP, Egholm M, Henrissat B, Heath AC, Knight R, Gordon JI. A core gut microbiome in obese and lean twins. *Nature* 2009; **457**: 480-484 [PMID: 19043404 DOI: 10.1038/nature07540]
 - 12 **Comes MC**, Gower-Rousseau C, Colombel JF, Belaïche J, Van Kruiningen HJ, Nuttens MC, Cortot A. Inflammatory bowel disease in married couples: 10 cases in Nord Pas de Calais region of France and Liège county of Belgium. *Gut* 1994; **35**: 1316-1318 [PMID: 7959244]
 - 13 **Shen H**, Ye F, Xie L, Yang J, Li Z, Xu P, Meng F, Li L, Chen Y, Bo X, Ni M, Zhang X. Metagenomic sequencing of bile from gallstone patients to identify different microbial community patterns and novel biliary bacteria. *Sci Rep* 2015; **5**: 17450 [PMID: 26625708 DOI: 10.1038/srep17450]
 - 14 **Jiang W**, Wu N, Wang X, Chi Y, Zhang Y, Qiu X, Hu Y, Li J, Liu Y. Dysbiosis gut microbiota associated with inflammation and impaired mucosal immune function in intestine of humans with non-alcoholic fatty liver disease. *Sci Rep* 2015; **5**: 8096 [PMID: 25644696 DOI: 10.1038/srep08096]
 - 15 **Loman NJ**, Misra RV, Dallman TJ, Constantinidou C, Gharbia SE, Wain J, Pallen MJ. Performance comparison of benchtop high-throughput sequencing platforms. *Nat Biotechnol* 2012; **30**: 434-439 [PMID: 22522955 DOI: 10.1038/nbt.2198]
 - 16 **Quail MA**, Smith M, Coupland P, Otto TD, Harris SR, Connor TR, Bertoni A, Swerdlow HP, Gu Y. A tale of three next generation sequencing platforms: comparison of Ion Torrent, Pacific Biosciences and Illumina MiSeq sequencers. *BMC Genomics* 2012; **13**: 341 [PMID: 22827831 DOI: 10.1186/1471-2164-13-341]
 - 17 **Hedin C**, van der Gast CJ, Rogers GB, Cuthbertson L, McCartney S, Stagg AJ, Lindsay JO, Whelan K. Siblings of patients with Crohn's disease exhibit a biologically relevant dysbiosis in mucosal microbial metacommunities. *Gut* 2016; **65**: 944-953 [PMID: 25856344 DOI: 10.1136/gutjnl-2014-308896]
 - 18 **Langille MG**, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA, Clemente JC, Burkpile DE, Vega Thurber RL, Knight R, Beiko RG, Huttenhower C. Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences. *Nat Biotechnol* 2013; **31**: 814-821 [PMID: 23975157 DOI: 10.1038/nbt.2676]
 - 19 **Ye F**, Shen H, Li Z, Meng F, Li L, Yang J, Chen Y, Bo X, Zhang X, Ni M. Influence of the Biliary System on Biliary Bacteria Revealed by Bacterial Communities of the Human Biliary and Upper Digestive Tracts. *PLoS One* 2016; **11**: e0150519 [PMID: 26930491 DOI: 10.1371/journal.pone.0150519]
 - 20 **Forbes JD**, Van Domselaar G, Bernstein CN. Microbiome Survey of the Inflamed and Noninflamed Gut at Different Compartments Within the Gastrointestinal Tract of Inflammatory Bowel Disease Patients. *Inflamm Bowel Dis* 2016; **22**: 817-825 [PMID: 26937623 DOI: 10.1097/MIB.0000000000000684]
 - 21 **Frank DN**, St Amand AL, Feldman RA, Boedeker EC, Harpaz N, Pace NR. Molecular-phylogenetic characterization of microbial community imbalances in human inflammatory bowel diseases. *Proc Natl Acad Sci USA* 2007; **104**: 13780-13785 [PMID: 17699621 DOI: 10.1073/pnas.0706625104]
 - 22 **Manichanh C**, Rigottier-Gois L, Bonnaud E, Gloux K, Pelletier E, Frangeul L, Nalin R, Jarrin C, Chardon P, Marteau P, Roca J, Dore J. Reduced diversity of faecal microbiota in Crohn's disease revealed by a metagenomic approach. *Gut* 2006; **55**: 205-211 [PMID: 16188921 DOI: 10.1136/gut.2005.073817]
 - 23 **Hijova E**, Soltesova A. Effects of probiotics and prebiotics in ulcerative colitis. *Bratisl Lek Listy* 2013; **114**: 540-543 [PMID: 24020713]
 - 24 **Gophna U**, Sommerfeld K, Gophna S, Doolittle WF, Veldhuyzen van Zanten SJ. Differences between tissue-associated intestinal microfloras of patients with Crohn's disease and ulcerative colitis. *J Clin Microbiol* 2006; **44**: 4136-4141 [PMID: 16988016 DOI: 10.1128/JCM.01004-06]
 - 25 **Wang W**, Chen L, Zhou R, Wang X, Song L, Huang S, Wang G, Xia B. Increased proportions of Bifidobacterium and the Lactobacillus group and loss of butyrate-producing bacteria in inflammatory bowel disease. *J Clin Microbiol* 2014; **52**: 398-406 [PMID: 24478468 DOI: 10.1128/JCM.01500-13]
 - 26 **Gevers D**, Kugathasan S, Denson LA, Vázquez-Baeza Y, Van Treuren W, Ren B, Schwager E, Knights D, Song SJ, Yassour M, Morgan XC, Kostic AD, Luo C, González A, McDonald D, Haberman Y, Walters T, Baker S, Rosh J, Stephens M, Heyman M, Markowitz J, Baldassano R, Griffiths A, Sylvester F, Mack D, Kim S, Crandall W, Hyams J, Huttenhower C, Knight R, Xavier RJ. The treatment-naïve microbiome in new-onset Crohn's disease. *Cell Host Microbe* 2014; **15**: 382-392 [PMID: 24629344 DOI: 10.1016/j.chom.2014.02.005]
 - 27 **Davenport M**, Poles J, Leung JM, Wolff MJ, Abidi WM, Ullman T, Mayer L, Cho I, Loke P. Metabolic alterations to the mucosal microbiota in inflammatory bowel disease. *Inflamm Bowel Dis* 2014; **20**: 723-731 [PMID: 24583479 DOI: 10.1097/MIB.0000000000000111]
 - 28 **Kamada N**, Chen GY, Inohara N, Núñez G. Control of pathogens and pathobionts by the gut microbiota. *Nat Immunol* 2013; **14**: 685-690 [PMID: 23778796 DOI: 10.1038/ni.2608]
 - 29 **Raszeja-Wyszomirska J**, Wunsch E, Krawczyk M, Rigopoulou EI, Kostrzewa K, Norman GL, Bogdanos DP, Milkiewicz P. Assessment of health related quality of life in polish patients with primary biliary cirrhosis. *Clin Res Hepatol Gastroenterol* 2016; **40**: 471-479 [PMID: 26621536 DOI: 10.1016/j.clinre.2015.10.006]

P- Reviewer: Bordas JM **S- Editor:** Qi Y **L- Editor:** Stewart G
E- Editor: Zhang FF





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>



ISSN 1007-9327

