

Name of Journal: World Journal of Gastroenterology

Manuscript NO: 33771

Manuscript Type: ORIGINAL ARTICLE

Prospective Study

**Fecal markers levels as predictors of the need for endoscopic balloon
dilation in Crohn's disease patients with anastomotic strictures**

Susana Lopes, Patrícia Andrade, Eduardo Rodrigues-Pinto, Joana Afonso,
Guilherme Macedo, Fernando Magro

Abstract

AIM

To evaluate the accuracy and the best cut-off value of fecal calprotectin (FC)

Match Overview

1	Crossref 119 words Andrade, Patrícia, João Santos-Antunes, Susana Rodrigues, Susana Lopes, and Guilherme Macedo. "Treatment with I	3%
2	Internet 39 words crawled on 24-Feb-2017 www.thieme-connect.com	1%
3	Internet 18 words crawled on 13-Jan-2017 pdfs.semanticscholar.org	1%
4	Crossref 14 words Nanda, Kavinderjit, William Courtney, Denise Keegan, Kathryn Byrne, Blathnaid Nolan, Diarmuid O'Donoghue, Hugh	<1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 97,200 条结果 (用时 0.86 秒)

显示的是以下查询字词的结果: **Fecal markers levels as predictors of the need for endoscopic balloon *dilatation* in Crohn's disease patients with anastomotic strictures**

仍然搜索: Fecal markers levels as predictors of the need for endoscopic balloon dilation in Crohn's disease patients with anastomotic strictures

Intestinal Stricture in Crohn's Disease

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: CW Chang - 2015 - 被引用次数: 13 - [相关文章](#)

2015年1月29日 - Crohn's disease (CD), an IBD, is the chronic inflammation of unknown ... 25% of CD patients have had at least one small bowel stricture and 10% have had at least ... RISK FACTORS AND PREDICTORS OF INTESTINAL STRICTURE Dilation of Crohn's strictures with endoscopy is widely used, but repeat ...

Crohn's disease complicated by strictures: a systematic review - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: F Rieder - 2013 - 被引用次数: 99 - [相关文章](#)

2013年4月26日 - The occurrence of strictures as a complication of Crohn's disease is in patients with CD have been reported to range from 48% to 52% at ... as predictive of a patient's progression to stricturing disease and a Through-the-scope endoscopic pneumatic balloon dilation of an anastomotic ileocecal stricture ...

[PDF] Current role of endoscopy in inflammatory bowel disease ... - NGHD

www.ncbi.nlm.nih.gov/pmc/articles/PMC4060000/ - [翻译此页](#)

Fecal markers levels as predictors of the need for endoscopic balloon dilation i

全部 图片 视频 新闻 购物 地图 图书

找到约 80,600 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

显示的是以下查询字词的结果: [Fecal markers levels as predictors of the need for endoscopic balloon dilatation in Crohn's disease patients with anastomotic strictures](#)

仍然搜索: [Fecal markers levels as predictors of the need for endoscopic balloon dilation in Crohn's disease patients with anastomotic strictures](#)

[Intestinal Stricture in Crohn's Disease - NCBI - NIH](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4316217/>

29 Jan 2015 ... [Crohn's disease](#) (CD), an IBD, is the chronic inflammation of unknown ... 25% of CD [patients have](#) had at least one small bowel [stricture](#) and 10% [have](#) had at least ... RISK FACTORS AND PREDICTORS OF INTESTINAL STRICTURE [Dilation](#) of Crohn's [strictures](#) with [endoscopy](#) is widely used, but repeat ...

[Poster presentations: Clinical: Therapy and observation \(2017\)](#)

<https://www.ecco-ibd.eu/index.php/publications/.../2.html>

P588 Ustekinumab for the treatment of [Crohn's disease patients](#) with TNF antagonist ... Is [fecal](#) calprotectin a [marker](#) of therapeutic response? ... P600 Infliximab trough [levels](#) for remission induction and long term therapy ... P631 Results and [predictors](#) of outcome of [endoscopic balloon dilation](#) of colonic [strictures](#) in ...

[\[PDF\] Multidisciplinary Management of Gastrointestinal Fibrotic Stenosis in ...](#)

<link.springer.com/content/pdf/10.1007/s10620-014-3421-y.pdf>

8 Nov 2014 ... the [need](#) for surgery in [patients](#) with CD. Unfortunately ... choice between [endoscopic balloon dilation](#), stricturo- plasty ... [Crohn's disease](#) (CD) is a chronic inflammatory bowel disease ... intestinal surgery are [predictors](#) of postoperative recurrence treatment could be estimated using this [fecal marker](#) [62].

[\[PDF\] Endoscopic Balloon Dilation of Intestinal Strictures in Crohn's Disease](#)

[www.gastrojournal.org/article/S0016-5085\(11\)60569-6/pdf](http://www.gastrojournal.org/article/S0016-5085(11)60569-6/pdf)

Validation of the Ulcerative Colitis [Endoscopic](#) Index of Severity (UCEIS) and ... original Thia1 study [values](#) (R2=0.81, 0.85, respectively). The current ... Bowel Follow Through (SBFT) Prior to IC in [Patients](#) With Suspected [Crohn's](#) after 2.4 (0.07-18.0) years and 46/88 (52.3%) [patients with anastomotic strictures](#) after 3.6

Fecal markers levels as predictors of the need for endoscopic balloon dilatation i

全部

图片

视频

新闻

地图

图书

找到约 76,700 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[Intestinal Stricture in Crohn's Disease - NCBI - NIH](#)<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4316217/>

29 Jan 2015 ... **Crohn's disease** (CD), an IBD, is the chronic inflammation of unknown ... 25% of CD **patients have** had at least one small bowel **stricture** and 10% **have** had at least ... RISK FACTORS AND **PREDICTORS** OF INTESTINAL **STRICTURE** **Dilation** of Crohn's **strictures** with **endoscopy** is widely used, but repeat ...

[\[PDF\] Endoscopic Balloon Dilation of Intestinal Strictures in Crohn's Disease](#)[www.gastrojournal.org/article/S0016-5085\(11\)60569-6/pdf](http://www.gastrojournal.org/article/S0016-5085(11)60569-6/pdf)

Validation of the Ulcerative Colitis **Endoscopic** Index of Severity (UCEIS) and ... original Thia1 study **values** ($R^2=0.81, 0.85$, respectively). The current ... Bowel Follow Through (SBFT) Prior to IC in **Patients** With Suspected **Crohn's** after 2.4 (0.07-18.0) years and 46/88 (52.3%) **patients with anastomotic strictures** after 3.6.

[\[PDF\] Multidisciplinary Management of Gastrointestinal Fibrotic Stenosis in ...](#)link.springer.com/content/pdf/10.1007/s10620-014-3421-y.pdf

8 Nov 2014 ... the **need** for surgery in **patients** with CD. Unfortunately ... choice between **endoscopic balloon dilation**, stricturo- plasty ... **Crohn's disease** (CD) is a chronic inflammatory bowel disease ... intestinal surgery are **predictors** of postoperative recurrence treatment could be estimated using this **fecal marker** [62].

[Poster presentations: Clinical: Therapy and observation \(2017\)](#)<https://www.ecco-ibd.eu/index.php/publications/.../2.html>

P588 Ustekinumab for the treatment of **Crohn's disease** **patients** with TNF antagonist ... Is **fecal calprotectin** a **marker** of therapeutic response? ... P600 Infliximab trough **levels** for remission induction and long term therapy ... P631 Results and **predictors** of outcome of **endoscopic balloon dilation** of colonic **strictures** in ...

[\[PDF\] P409 Smoking is associated with extra-intestinal manifestations in ...](#)