

[全部](#)[图片](#)[购物](#)[视频](#)[新闻](#)[更多](#)[设置](#)[工具](#)

找到约 5,650 条结果 (用时 0.67 秒)

显示的是以下查询字词的结果：[**Procalcitonin as diagnostic marker to distinguish upper and lower gastrointestinal perforation**](#)

仍然搜索：Procalcitonin as diagnosis marker to distinguish upper and lower gastrointestinal perforation

Google 学术：[**Procalcitonin as diagnostic marker to distinguish upper and lower gastrointestinal perforation**](#)

Procalcitonin for diagnosis of infection and guide to ... - Schuetz - 被引用次数：276

Clinical applications of C-reactive protein in pediatrics - Jaye - 被引用次数：576

... pediatric appendicitis: usefulness of laboratory markers - Kwan - 被引用次数：71

Procalcitonin for diagnosis of infection and guide to antibiotic ... - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者：P Schuetz - 2011 - 被引用次数：276 - [相关文章](#)

2011年9月22日 - For patients with upper and lower respiratory tract infection, post-operative ... and may help to distinguish bacterial infections from viral illnesses [8-11]. ... as a marker to exclude perforation and ischemia in obstructive bowel ...

Evolution of imaging for abdominal perforation

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者：JP Singh - 2010 - 被引用次数：42 - [相关文章](#)

The clinical diagnosis of the site of GI tract perforation is difficult as the the most significant finding in distinguishing upper from lower GI tract perforation.

缺少字词：procalcitonin marker

Name of journal: *World Journal of Gastroenterology*

Manuscript NO: 33778

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Procalcitonin as diagnosis marker to distinguish upper and lower
gastrointestinal perforation

Yang Gao, Kai Kang, Hai-Tao Liu, Xing Zhang, Rui Huang, Jing-Dong Qu,
Sicong Wang, Rui-Jin Liu, Yan-Song Liu, Hong-Liang Wang, Kai-Jiang Yu

Abstract

AIM

Match Overview

1	Internet 14 words crawled on 17-Mar-2017 discovery.ucl.ac.uk	1%
2	Crossref 13 words Takakura, Yuji, Takao Hinoi, Hiroyuki Egi, Manabu Shimomura, Tomohiro Adachi, Yasufumi Saito, Naoki Tanimine, Ma	1%

Procalcitonin as diagnostic marker to distinguish upper and lower gastrointestina

全部 图片 视频 新闻 地图 图书

找到约 8,410 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Google 学术: Procalcitonin as diagnostic marker to distinguish upper and lower gastrointestinal perforation](#)

[Procalcitonin for diagnosis of infection and guide to ...](#) - Schuetz - 被引用次数: 297

[Clinical applications of C-reactive protein in pediatrics](#) - Jaye - 被引用次数: 593

[... pediatric appendicitis: usefulness of laboratory markers](#) - Kwan - 被引用次数: 74

[Procalcitonin-guided diagnosis and antibiotic stewardship revisited ...](#)

<https://bmcmmedicine.biomedcentral.com/.../10.../s12916-017-0795-7>

24 Jan 2017 ... Procalcitonin Pneumonia Respiratory tract infection Sepsis Antibiotic stewardship ... the use of the infection marker procalcitonin (PCT) to improve the diagnosis ... PCT-guided therapy was associated with lower antibiotic exposure with In SIRS patients with negative blood cultures, PCT can differentiate ...

[Procalcitonin for diagnosis of infection and guide to antibiotic ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3186747/>

22 Sep 2011 ... For patients with upper and lower respiratory tract infection, post-operative ... and may help to distinguish bacterial infections from viral illnesses [8-11]. ... as a marker to exclude perforation and ischemia in obstructive bowel ...

[The Diagnostic Value of D-dimer, Procalcitonin and CRP in Acute ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3520016/>

13 Nov 2012 ... The PCT values were higher than the upper normal limit in 20 ... However, PCT and D-dimer showed lower diagnostic values (26% ... differentiate between phlegmonous appendicitis and perforated appendicitis. ... Recently, D-dimer has also been used as a diagnostic marker in the Dis Colon Rectum.

[Procalcitonin and intestinal ischemia: A review of the literature](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4273127/>

[All](#)[Shopping](#)[Images](#)[Videos](#)[News](#)[More](#)[Settings](#)[Tools](#)

About 8,710 results (1.05 seconds)

Showing results for **Procalcitonin as diagnostic** marker to distinguish upper and lower gastrointestinal perforation

Search instead for Procalcitonin as diagnosis marker to distinguish upper and lower gastrointestinal perforation

Scholarly articles for **Procalcitonin as diagnostic marker to distinguish upper and lower gastrointestinal perforation**

Procalcitonin for **diagnosis** of infection and guide to ... - [Schuetz](#) - Cited by 298

Clinical applications of C-reactive protein in pediatrics - [Jaye](#) - Cited by 592

... pediatric appendicitis: usefulness of laboratory **markers** - [Kwan](#) - Cited by 75

Procalcitonin for diagnosis of infection and guide to antibiotic ...

<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#)

by P Schuetz - 2011 - Cited by 296 - Related articles

22 Sep 2011 - For patients with **upper and lower** respiratory tract infection, post-operative ... and may help to **distinguish** bacterial infections from viral illnesses [8-11]. ... as a **marker** to exclude **perforation** and ischemia in obstructive **bowel** ...

PDF(1898K) - Wiley Online Library

onlinelibrary.wiley.com/doi/10.14442/jgfm.17.1_5/pdf

by T Mayumi - 2016 - Cited by 4 - Related articles

diagnosis, and primary treatment for acute abdomen was performed to assessed to **differentiate** cases requiring emergency surgery (level 2, CQ50 Does the measurement of **procalcitonin** (PCT) have utility in the **diagnosis** For esophageal and **gastric perforations**, an **upper GI** series using other **markers** (CQ68).

A focus on intra-abdominal infections | World Journal of Emergency ...

<https://wjeb.biomedcentral.com/articles/10.1186/1749-7922-5-9>

by M Sartelli - 2010 - Cited by 86 - Related articles