

World Journal of *Gastroenterology*

World J Gastroenterol 2017 August 7; 23(29): 5257-5450





EDITORIAL

- 5257 Liver and the defects of cholesterol and bile acids biosynthesis: Rare disorders many diagnostic pitfalls
Corso G, Dello Russo A, Gelzo M

REVIEW

- 5266 Contribution of galectin-1, a glycan-binding protein, to gastrointestinal tumor progression
Bacigalupo ML, Carabias P, Troncoso MF
- 5282 From diagnosis to treatment of hepatocellular carcinoma: An epidemic problem for both developed and developing world
Dimitroulis D, Damaskos C, Valsami S, Davakis S, Garmpis N, Spartalis E, Athanasiou A, Moris D, Sakellariou S, Kykalos S, Tsourouflis G, Garmpi A, Delladetsima I, Kontzoglou K, Kouraklis G

ORIGINAL ARTICLE

Basic Study

- 5295 Partial external biliary diversion in bile salt export pump deficiency: Association between outcome and mutation
Ellinger P, Stindt J, Dröge C, Sattler K, Stross C, Kluge S, Herebian D, Smits SHJ, Burdelski M, Schulz-Jürgensen S, Ballauff A, Schulte am Esch J, Mayatepek E, Häussinger D, Kubitz R, Schmitt L
- 5304 Celecoxib-induced gastrointestinal, liver and brain lesions in rats, counteraction by BPC 157 or L-arginine, aggravation by L-NAME
Drmic D, Kolenc D, Ilic S, Bauk L, Sever M, Zenko Sever A, Luetic K, Suran J, Seiwert S, Sikiric P
- 5313 Hwangryunhaedok-tang induces the depolarization of pacemaker potentials through 5-HT₃ and 5-HT₄ receptors in cultured murine small intestine interstitial cells of Cajal
Kim HJ, Lee GS, Kim H, Kim BJ
- 5324 MicroRNA exhibit altered expression in the inflamed colonic mucosa of ulcerative colitis patients
Valmiki S, Ahuja V, Paul J
- 5333 Salvianolic acid B protects hepatocytes from H₂O₂ injury by stabilizing the lysosomal membrane
Yan XF, Zhao P, Ma DY, Jiang YL, Luo JJ, Liu L, Wang XL
- 5345 PBX1 attributes as a determinant of connexin 32 downregulation in *Helicobacter pylori*-related gastric carcinogenesis
Liu XM, Xu CX, Zhang LF, Huang LH, Hu TZ, Li R, Xia XJ, Xu LY, Luo L, Jiang XX, Li M

Case Control Study

- 5356 Influence of dietary isoflavone intake on gastrointestinal symptoms in ulcerative colitis individuals in remission

Gląbska D, Guzek D, Grudzińska D, Lech G

- 5364 Genetic polymorphisms of *MAFK*, encoding a small Maf protein, are associated with susceptibility to ulcerative colitis in Japan

Arisawa T, Nakamura M, Otsuka T, Jing W, Sakurai N, Takano H, Hayashi T, Ota M, Nomura T, Hayashi R, Shimasaki T, Tahara T, Shibata T

Retrospective Study

- 5371 Elaboration and validation of Crohn's disease anoperineal lesions consensual definitions

Horaist C, de Parades V, Abramowitz L, Benfredj P, Bonnaud G, Boucharde D, Fathallah N, Sénéjoux A, Siproudhis L, Staumont G, Viguier M, Marteau P

- 5379 Efficacy of tolvaptan for the patients with advanced hepatocellular carcinoma

Miyazaki M, Yada M, Tanaka K, Senjyu T, Goya T, Motomura K, Kohjima M, Kato M, Masumoto A, Kotoh K

- 5386 Outcomes of preoperative endoscopic nasobiliary drainage and endoscopic retrograde biliary drainage for malignant distal biliary obstruction prior to pancreaticoduodenectomy

Zhang GQ, Li Y, Ren YP, Fu NT, Chen HB, Yang JW, Xiao WD

Clinical Trials Study

- 5395 Phase I clinical study of personalized peptide vaccination combined with radiotherapy for advanced hepatocellular carcinoma

Shen J, Wang LF, Zou ZY, Kong WW, Yan J, Meng FY, Chen FJ, Du J, Shao J, Xu QP, Ren HZ, Li RT, Wei J, Qian XP, Liu BR

Observational Study

- 5405 Transition clinic attendance is associated with improved beliefs and attitudes toward medicine in patients with inflammatory bowel disease

Fu N, Jacobson K, Round A, Evans K, Qian H, Bressler B

- 5412 Gut barrier failure biomarkers are associated with poor disease outcome in patients with primary sclerosing cholangitis

Tornai T, Palyu E, Vitalis Z, Tornai I, Tornai D, Antal-Szalmas P, Norman GL, Shums Z, Veres G, Dezsofi A, Par G, Par A, Orosz P, Szalay F, Lakatos PL, Papp M

- 5422 Efficacy of forced coagulation with low high-frequency power setting during endoscopic submucosal dissection

Ishida T, Toyonaga T, Ohara Y, Nakashige T, Kitamura Y, Ariyoshi R, Takihara H, Baba S, Yoshizaki T, Kawara F, Tanaka S, Morita Y, Umegaki E, Hoshi N, Azuma T

Prospective Study

- 5431** Clinical outcomes of isolated renal failure compared to other forms of organ failure in patients with severe acute pancreatitis

Gougol A, Dugum M, Dudekula A, Greer P, Slivka A, Whitcomb DC, Yadav D, Papachristou GI

SYSTEMATIC REVIEW

- 5438** Laparoscopic ultrasonography as an alternative to intraoperative cholangiography during laparoscopic cholecystectomy

Dili A, Bertrand C

ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Shinji Tanaka, MD, PhD, Professor, Department of Endoscopy, Hiroshima University Hospital, Hiroshima 734-8551, Japan

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7th, 14th, 21st, and 28th each month. The *WJG* Editorial Board consists of 1375 experts in gastroenterology and hepatology from 68 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

World Journal of Gastroenterology (*WJG*) is now indexed in Current Contents[®]/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch[®]), Journal Citation Reports[®], Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2017 edition of Journal Citation Reports[®] cites the 2016 impact factor for *WJG* as 3.365 (5-year impact factor: 3.176), ranking *WJG* as 29th among 79 journals in gastroenterology and hepatology (quartile in category Q2).

FLYLEAF

I-IX Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Dan Li*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Ze-Mao Gong*
Proofing Editorial Office Director: *Jin-Lei Wang*

NAME OF JOURNAL
World Journal of Gastroenterology

ISSN
ISSN 1007-9327 (print)
ISSN 2219-2840 (online)

LAUNCH DATE
October 1, 1995

FREQUENCY
Weekly

EDITORS-IN-CHIEF
Damian Garcia-Olmo, MD, PhD, Doctor, Professor, Surgeon, Department of Surgery, Universidad Autonoma de Madrid; Department of General Surgery, Fundacion Jimenez Diaz University Hospital, Madrid 28040, Spain

Stephen C Strom, PhD, Professor, Department of Laboratory Medicine, Division of Pathology, Karolinska Institutet, Stockholm 141-86, Sweden

Andrzej S Tarnawski, MD, PhD, DSc (Med), Professor of Medicine, Chief Gastroenterology, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach,

CA 90822, United States

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE
Jin-Lei Wang, Director
Yuan Qi, Vice Director
Ze-Mao Gong, Vice Director
World Journal of Gastroenterology
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.fpublishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.fpublishing.com/helpdesk>

<http://www.wjgnet.com>

PUBLICATION DATE
August 7, 2017

COPYRIGHT
© 2017 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.fpublishing.com>

Basic Study

Hwangryunhaedok-tang induces the depolarization of pacemaker potentials through 5-HT₃ and 5-HT₄ receptors in cultured murine small intestine interstitial cells of Cajal

Hyun Jung Kim, Guem San Lee, Hyungwoo Kim, Byung Joo Kim

Hyun Jung Kim, Byung Joo Kim, Division of Longevity and Biofunctional Medicine, Healthy Aging Korean Medical Research Center, School of Korean Medicine, Pusan National University, Yangsan 50612, South Korea

Guem San Lee, Wonkwang University College of Korean Medicine, Iksan 54538, South Korea

Hyungwoo Kim, Division of Pharmacology, School of Korean Medicine, Pusan National University, Yangsan 50612, South Korea

Byung Joo Kim, Division of Longevity and Biofunctional Medicine, School of Korean Medicine, Pusan National University, Gyeongsangnamdo 50612, South Korea

Author contributions: Kim HJ and Kim BJ designed the research; Kim HJ, Lee GS and Kim H performed the experiments; Kim HJ and Kim BJ analyzed the data; and Kim HJ and Kim BJ wrote the paper.

Supported by the National Research Foundation of Korea Grant funded by the Korea Government (MSIP), No. 2014R1A5A2009936.

Institutional animal care and use committee statement: The animal care and experiments in accordance with the guidelines issued by the Institutional Animal Care and Use Committee at Pusan National University (Busan, Republic of Korea; Approval No. PNU-2015-1036).

Conflict-of-interest statement: The authors have no disclosures to report.

Data sharing statement: No additional are available.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and

the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Correspondence to: Byung Joo Kim, PhD, Associate Professor, Division of Longevity and Biofunctional Medicine, School of Korean Medicine, Pusan National University, Beomeori, Mulgeum-eup, Yangsan, Gyeongsangnamdo 50612, South Korea. vision@pusan.ac.kr
Telephone: +82-51-5108469
Fax: +82-51-5108420

Received: March 2, 2017

Peer-review started: March 4, 2017

First decision: April 6, 2017

Revised: April 18, 2017

Accepted: June 9, 2017

Article in press: June 12, 2017

Published online: August 7, 2017

Abstract**AIM**

To investigate the effects of a water extract of Hwangryunhaedok-tang (HHTE) on the pacemaker potentials of mouse interstitial cells of Cajal (ICCs).

METHODS

We dissociated ICCs from small intestines and cultured. ICCs were immunologically identified using an anti-c-kit antibody. We used the whole-cell patch-clamp configuration to record the pacemaker potentials generated by cultured ICCs under the current clamp mode ($I = 0$). All experiments were performed at 30 °C-32 °C

RESULTS

HHTE dose-dependently depolarized ICC pacemaker potentials. Pretreatment with a 5-HT₃ receptor anta-

gonist (Y25130) or a 5-HT₄ receptor antagonist (RS39604) blocked HHTE-induced pacemaker potential depolarizations, whereas pretreatment with a 5-HT₇ receptor antagonist (SB269970) did not. Intracellular GDP β S inhibited HHTE-induced pacemaker potential depolarization and pretreatment with a Ca²⁺-free solution or thapsigargin abolished the pacemaker potentials. In the presence of a Ca²⁺-free solution or thapsigargin, HHTE did not depolarize ICC pacemaker potentials. In addition, HHTE-induced pacemaker potential depolarization was unaffected by a PKC inhibitor (calphostin C) or a Rho kinase inhibitor (Y27632). Of the four ingredients of HHT, Coptidis Rhizoma and Gardeniae Fructus more effectively inhibited pacemaker potential depolarization.

CONCLUSION

These results suggest that HHTE dose-dependently depolarizes ICC pacemaker potentials through 5-HT₃ and 5-HT₄ receptors *via* external and internal Ca²⁺ regulation and *via* G protein-, PKC- and Rho kinase-independent pathways.

Key words: Hwangryunhaedok-tang; Interstitial cells of Cajal; Pacemaker potentials; Gastrointestinal tract; Motility

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Hwangryunhaedok-tang, a traditional herbal medicine, has been used to treat gastrointestinal diseases. Our data suggest that HHT can depolarize the pacemaker potentials of cultured interstitial cells of Cajal from mouse small intestine and may be a novel prokinetic agent.

Kim HJ, Lee GS, Kim H, Kim BJ. Hwangryunhaedok-tang induces the depolarization of pacemaker potentials through 5-HT₃ and 5-HT₄ receptors in cultured murine small intestine interstitial cells of Cajal. *World J Gastroenterol* 2017; 23(29): 5313-5323 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v23/i29/5313.htm> DOI: <http://dx.doi.org/10.3748/wjg.v23.i29.5313>

INTRODUCTION

Herbal medicines are becoming increasingly popular in modern societies and substantial research has been conducted to establish their pharmacological properties according to scientific principles^[1]. Since ancient times, people have used herbal medicines to treat and prevent disease. Due to their natural origins, herbal medicines have mild activities and few side effects; for this reason, many people are becoming increasingly interested in traditional herbal medicines^[2-5].

Hwangryunhaedok-tang (HHT; Huang-Lian-Jie-Tang in Chinese or Oren-gedoku-to in Japanese) is a

preparation that consists of Rhizoma Coptidis, Radix Scutellariae, Cortex Phellodendri and Fructus Gardenia^[6]. HHT has been used to treat various symptoms, including inflammatory diseases^[7,8], cardiovascular diseases^[9], diabetes mellitus^[10], liver diseases^[11,12], brain injury^[13,14] and obesity^[15]. HHT is also used to prevent or ameliorate gastrointestinal (GI) diseases^[16-19]. HHT has been shown to have a preventive effect on the development of water immersion restraint (WIR) stress-induced acute gastric lesions in rats^[16] and it suppresses the symptoms of inflammatory bowel disease by reducing levels of interleukin-8 (IL-8), leukotriene B₄ (LTB₄) and prostaglandin E₂ (PGE₂)^[17]. In addition, HHT has been shown to prevent lethal indomethacin (IND)-induced enteropathy by decreasing adenosine deaminase (ADA) and, thus, increasing the adenosine levels in female BALB/c mice^[18].

Interstitial cells of Cajal (ICCs) contribute to normal GI function, such as by generating electrical slow waves and mediating neuromuscular signaling^[19-21]. Damage to ICCs or reductions in ICC numbers have been described in many GI motility disorders^[22-24]; therefore, it is very important to understand the mechanisms of ICC pacemaker activity. However, no scientific evidence of a relationship between HHT and ICCs has been presented until now. In the present study, we investigated the effects of water extracts of HHT and its components on the pacemaker potentials of cultured ICCs from murine small intestine.

MATERIALS AND METHODS

Preparation of HHT extract

HHT was purchased from I-WORLD Pharmaceuticals (Incheon, South Korea). HHT (3.67 g) consists of Coptidis Rhizoma (0.67 g), Scutellariae Radix (1.00 g), Phellodendri Cortex (1.00 g) and Gardeniae Fructus (1.00 g). The materials were authenticated by Prof. Hyungwoo Kim (Division of Pharmacology, Pusan National University, School of Korean Medicine, Yangsan, Korea). HHT extract (HHTE) was prepared by boiling HHT in water for three hours. The standard adult dose of HHTE is 10-15 g (crude material) per day. More information about HHTE can be found at the I-WORLD Pharmaceuticals Homepage (<http://i-pharm.koreasme.com>). HHTE was dissolved in distilled water at a concentration of 0.5 g (crude drug)/mL and stored in a refrigerator until required.

Reagents

Geniposide, berberine chloride, baicalin and wogonin were purchased from Wako (Japan). High-performance liquid chromatography (HPLC)-grade water and methanol were obtained from JT Baker Inc. (Phillipsburg, NJ, United States). Trifluoroacetic acid was purchased from Sigma-Aldrich (St. Louis, MO, United States).

Preparation of standard solutions

Each standard compound (geniposide, berberine chloride,




Figure 1 Effects of water extract of Hwangryunhaedok-tang on the pacemaker potentials of cultured interstitial cells of Cajal from murine small intestine. A-D: HHTE (0-50 mg/mL) induced pacemaker potential depolarizations and decreased amplitudes in a concentration-dependent manner. Responses to HHTE are summarized (E-G). Bars indicate the means \pm SEMs. ^e $P < 0.001$: significantly different from the control. CTRL: Control; HHTE: Water extract of Hwangryunhaedok-tang.

baicaline and wogonin) was accurately weighed and dissolved in methanol at a concentration of 1000 g/mL and the solution was 10-fold diluted before injection.

Chromatographic conditions

An Agilent 1200 (Agilent Technologies, Santa Clara, CA, United States) equipped with a quaternary pump, autosampler, column oven and diode-array detector was used. Acquired data were processed using Chemstation software (Ver. B. 03. 02). Chromatographic separation was performed on an XDB C₁₈ column (4.6 mm \times 150 mm, 5 μ m; Agilent, United States) at 35 $^{\circ}$ C. The mobile phase consisted of water with 0.1% trifluoroacetic acid (A) and methanol (B). The following program was used: 30% (B) for 1 min, 30%-80% (B) over 1 min-13 min and 80% (B) for 1 min, which was followed by a re-equilibration to 30% (B). The flow rate was set at 0.8 mL/min and the injection volume was 10 μ L. The detection wavelengths were 240 nm for geniposide, 260 nm for berberine chloride and 275 nm for baicalin and wogonin.

Identification of standard compounds in HHTE

Standard compounds were detected in the chromatogram of HHTE at the following retention times: 5.2 min for geniposide, 9.1 min for berberine chloride, 10.1 min for baicalin and 14.7 min for wogonin (Figure 1).

Ethics

We conducted the animal care and experiments in accordance with the guidelines issued by the Institutional Animal Care and Use Committee (IACUC) at Pusan National University (Busan, Republic of Korea; Approval no. PNU-2015-1036) and those issued by the National Institute of Health Guide for the Care and Use of Laboratory Animals.

Preparation of ICC and ICC clusters

ICR mice (Samtako Bio Korea Co., Ltd., Osan, South Korea; 3-7 day-old; weighing 1.9 g-2.2 g) of either sex were used and kept under controlled conditions (20 $^{\circ}$ C \pm 3 $^{\circ}$ C, RH 49% \pm 6% and light on 7 a.m.-7 p.m.). The small intestines were removed and opened; then, the luminal contents were cleaned using Krebs-Ringer bicarbonate solution and the obtained tissues were pinned to the bases of Sylgard dishes. Mucosae were removed and small tissue strips of intestine muscle were equilibrated for 25 min in Ca²⁺-free Hank's solution with the following solutions (in mmol/L): KCl 5.36, NaCl 125, NaOH 0.34, Na₂HCO₃ 0.44, glucose 10, sucrose 2.9 and HEPES 11 (pH 7.4). Cells were then dispersed in an enzyme solution [1.3 mg/mL collagenase (Worthington Biochemical, Lakewood, NJ, United States), 2.1 mg/mL bovine serum albumin (Sigma-Aldrich, St. Louis, MO, United States), 2.6

mg/mL trypsin inhibitor (Sigma-Aldrich, St. Louis, MO, United States) and 0.50 mg/mL ATP (Sigma-Aldrich, St. Louis, MO, United States)]. Cells were then cultured at 37 °C in a 95% O₂-5% CO₂ incubator in smooth muscle growth medium (SMGM) (Clonetics, San Diego, CA, United States) supplemented with 2% antibiotics/antimycotics (Gibco, Grand Island, NY, United States) and 5 ng/mL of murine stem cell factor (SCF) (Sigma-Aldrich, St. Louis, MO, United States). All experiments were performed on ICC cells that were cultured for < 12 h. ICCs were identified immunologically using an anti-c-kit antibody (phycoerythrin (PE)-conjugated rat anti-mouse c-kit monoclonal antibody) (eBioscience, San Diego, CA, United States) at a dilution of 1:50 for 20 min^[25]. Because ICCs differed morphologically from other cell types in culture, they were identified by phase contrast microscopy after incubation with anti c-kit antibody.

Patch-clamp experiments

The cells were transferred onto a solution chamber on the stage of an inverted microscope (IX70, Olympus, Japan). The whole-cell patch-clamp technique was used to record the membrane potentials (current clamp) of cultured ICCs as described previously^[21,22] and an Axopatch 200B amplifier (Axon Instruments, Foster City, CA, United States) was used to amplify membrane potentials. The bath solution [Normal Tyrode (NT)] contained 135 mmol/L NaCl, 5 mmol/L KCl, 2 mmol/L CaCl₂, 1 mmol/L MgCl₂, 10 mmol/L glucose and 10 mmol/L HEPES, with a pH adjusted to 7.4 using NaOH. The internal solution contained 145 mmol/L Cs-glutamate, 8 mmol/L NaCl, 10 mmol/L BAPTA and 10 mmol/L HEPES-CsOH, with a pH adjusted to 7.2 with CsOH. All experiments were performed at 30-32 °C.

Drugs

Y25130, RS39604 and SB269970 were purchased from Tocris Bioscience (Bristol, United Kingdom) and all other drugs were obtained from Sigma-Aldrich (St. Louis, MO, United States). For stock solutions, all drugs were dissolved in distilled water or dimethylsulfoxide (DMSO) and stored at -20 °C. The final concentration of DMSO in the bath solution was always < 0.1%. Furthermore, the addition of these chemicals did not alter the bath solution pH.

Statistical analysis

The results are expressed as the means ± SEMs. N values refer to the numbers of cells used in the experiments. For multiple comparison analysis, we used one-way ANOVA with Bonferroni's *post hoc* comparison. For statistical analyses, we used Prism 6.0 (GraphPad Software Inc., La Jolla, CA, United States) and Origin version 8.0 (OriginLab Corporation, Northampton, MA, United States). *P* values < 0.05 were considered statistically significant.

RESULTS

Functional components of HHTE

The presence of geniposide, berberine chloride, baicalin and wogonin in HHTE was established by HPLC and their levels were quantified using calibration curves obtained with purchased standards^[26]. Validation of the method confirmed its reliability and stability. The contents of the four marker components of HHTE by HPLC were 2.684% ± 0.014% in geniposide, 0.993% ± 0.006% in berberine chloride, 5.081% ± 0.027% in baicalin and 0.040% ± 0.003% in wogonin.

Effects of HHTE on the pacemaker potentials of ICCs

Using the whole-cell patch clamp technique, we investigated the characteristics of cell clusters. Cell clusters formed network-like structures after culture for < 12 h and were viable as demonstrated by the generation of spontaneous rhythmic contractions. ICCs were immunologically identified using an anti-c-kit antibody^[25]. All experiments were performed at 30 °C-32 °C. Under the current clamp mode (*I* = 0), ICCs generated pacemaker potentials (Figure 1A) with a mean resting membrane potential of -58.3 ± 1.5 mV and a mean amplitude of 24.9 ± 1.1 mV. Initially, we examined the effects of HHTE on the pacemaker potentials of ICCs. HHTE (10-50 mg/mL) was found to depolarize pacemaker potentials and decrease their amplitudes in a concentration-dependent manner (Figure 1B-D). In the presence of HHTE, the mean degrees of depolarization were 0.1 ± 0.1 mV at 10 mg/mL, 11.1 ± 0.9 mV at 20 mg/mL and 19.6 ± 1.1 mV at 30 mg/mL (Figure 1E, *n* = 17), the mean amplitudes were 24.8 ± 1.0 mV at 10 mg/mL, 12.1 ± 0.3 mV at 20 mg/mL and 1.7 ± 0.6 mV at 30 mg/mL (Figure 1F, *n* = 17) and the mean frequencies were 17.5 ± 1.0 cycles/min at 10 mg/mL, 4.5 ± 1.3 cycles/min at 20 mg/mL and 1.8 ± 0.9 cycles/min at 30 mg/mL (Figure 1G, *n* = 17). The summarized values showing the effects of HHTE on pacemaker potentials are provided in Figure 1E, 1F and 1G. These results show HHTE dose-dependently depolarizes the ICC pacemaker potentials.

HHTE mechanisms of action on ICCs

To identify the connection between HHTE and its receptors in cultured ICCs, we studied the correlation of 5-HT receptors because these receptors are involved in GI tract motility, which is strongly associated with prokinetic activity^[27]. In previous studies, 5-HT has been shown to interact with seven 5-HT receptor subtypes in the GI tract and only murine small intestine ICCs express three of these receptors (5-HT₃, 4 and 7)^[28,29]. To confirm the 5-HT receptor subtypes involved in the effects of HHTE, we used the various 5-HT receptor antagonists and then treated samples with HHTE. A 5-HT₃ receptor antagonist (Y25130), a 5-HT₄ receptor antagonist (RS39604), or a 5-HT₇




Figure 2 Effects of 5-HT receptor antagonists on water extract of Hwangryunhaedok-tang-induced pacemaker potential depolarizations in the cultured interstitial cells of Cajal of murine small intestine. A: Y25130 (a 5-HT₃ receptor antagonist) inhibited HHTE-induced responses. B: A 5-HT₄ receptor antagonist, RS39604, blocked HHTE-induced responses. C: A 5-HT₇ receptor antagonist, SB269970, did not block HHTE-induced responses. D and E responses to 5-HT receptor antagonists are summarized. Bars indicate the means \pm SEMs. ^a $P < 0.05$ and ^e $P < 0.001$: Significantly different from nontreated controls. CTRL: Control; HHTE: Water extract of Hwangryunhaedok-tang.




Figure 3 Effects of GDP- β -S on water extract of Hwangryunhaedok-tang-induced pacemaker potential depolarizations in cultured interstitial cells of Cajal. A: When GDP- β -S (1 mmol/L) was applied in the pipette, HHTE did not depolarize the pacemaker potential. B and C: HHTE-induced responses are summarized. Bars indicate the means \pm SEMs. ^e $P < 0.001$: Significantly different from nontreated controls. CTRL: Control; HHTE: Water extract of Hwangryunhaedok-tang.

receptor antagonist (SB269970) was used to pretreat the samples at 10 μ mol/L for 5 min; then, HHTE was added. In the case of pretreatment with Y25130, HHTE-induced pacemaker potential depolarization was blocked (Figure 2A) and HHTE-induced pacemaker potential depolarization in the presence of Y25130 was 0.8 ± 0.5 mV ($n = 5$; Figure 2D and E). RS39604 also blocked HHTE-induced pacemaker potential depolarization. HHTE-induced pacemaker potential depolarization in the presence of RS39604 was 0.5 ± 0.6 mV ($n = 5$; Figure 2B, D and E). However, pretreatment with SB269970 did not block the HHTE-induced pacemaker potential depolarization. HHTE-

induced pacemaker potential depolarization in the presence of SB269970 was 19.4 ± 1.8 mV ($n = 5$; Figure 2C-E). These results indicate that HHTE affects ICCs in the murine small intestine through the 5-HT₃ and 5-HT₄ receptors.

Connection of G protein in HHTE-induced pacemaker potential depolarization in ICCs

To determine whether G proteins are involved in HHTE-induced pacemaker potential depolarization, we used GDP- β -S (a non-hydrolysable guanosine 5' diphosphate analogue), which inhibits G-protein activation^[30,31]. HHTE (50 mg/mL) induced pacemaker potential depolarization in ICCs (Figure 1D); however, when GDP- β -S (1 mmol/L) was applied in the pipette solution, HHTE (50 mg/mL) only caused slight pacemaker potential depolarization (5.2 ± 0.3 mV, $n = 5$; Figure 3A and B). These results indicate that G protein is involved in HHTE-induced ICC pacemaker potential depolarization.

Effects of external and internal Ca²⁺ on HHTE-induced pacemaker potential depolarization in ICCs

Regulation of external and internal Ca²⁺ has an important role in mediating GI smooth muscle contraction and the generation of pacemaker potentials in ICCs^[32]. To identify the roles of Ca²⁺, HHTE was applied in external Ca²⁺-free conditions or in the presence of an inhibitor of Ca²⁺-ATPase in the endoplasmic reticulum (thapsigargin). In the external Ca²⁺-free conditions, pacemaker potentials were abolished; under this condition, HHTE did not induce pacemaker potential depolarization ($n = 6$; Figure 4A and C). In addition, pretreatment with thapsigargin abolished pacemaker potentials and under this condition, HHTE also did not induce pacemaker potential depolarization ($n =$




Figure 4 Effects of external and internal Ca^{2+} on water extract of Hwangryunhaedok-tang-induced pacemaker potential depolarization in cultured interstitial cells of Cajal from murine small intestine. A: In external Ca^{2+} -free conditions, the pacemaker potentials were abolished and HHTE had no effect. B: Thapsigargin abolished the pacemaker potentials and HHTE had no effects; C-E: HHTE-induced pacemaker potential depolarizations are summarized. Bars represent the means \pm SEMs. $^{\circ}P < 0.001$: Significantly different from untreated controls. CTRL: Control; HHTE: Water extract of Hwangryunhaedok-tang.




Figure 5 Effects of protein kinase C inhibitor and Rho kinase inhibitor on water extract of Hwangryunhaedok-tang-induced pacemaker potential depolarizations in cultured interstitial cells of Cajal from murine small intestine. Both (A) calphostin C (a PKC inhibitor) and (B) Y27631 (a Rho kinase inhibitor) had no effects on the pacemaker potentials; under these conditions, HHTE induced pacemaker potential depolarizations. C and D responses to HHTE in the presence of calphostin C or Y27631 are summarized. Bars represent the means \pm SEMs. $^{\circ}P < 0.001$: significantly different from the nontreated control. CTRL: Control; HHTE: Water extract of Hwangryunhaedok-tang; PKC: Protein kinase C.

5; Figure 4B and D). These results show that HHTE-induced pacemaker potential depolarization depends

on internal and external Ca^{2+} .

Pathways mediating HHTE-induced pacemaker potential depolarization in ICCs

To determine whether HHTE-induced pacemaker potential depolarization is associated with protein kinase C (PKC) or Rho kinase, we pretreated ICCs with a PKC inhibitor (calphostin C, 10 μ mol/L, Figure 5A) or a Rho kinase inhibitor (Y27631, 1 μ mol/L, Figure 5B) for 5 min. HHTE-induced pacemaker potential depolarizations were not inhibited by calphostin C or Y27631 (Figure 5C). We confirmed that treatment with these inhibitors alone did not affect pacemaker potential depolarization. These results suggest that HHTE-induced pacemaker potential depolarization is not mediated by PKC or Rho kinase in the cultured ICCs of murine small intestine.

Effects of Coptidis Rhizoma, Scutellariae Radix, Phellodendri Cortex and Gardeniae Fructus extracts in ICCs

We investigated the effects of the four components of HHT, Coptidis Rhizoma, Scutellariae Radix, Phellodendri Cortex and Gardeniae Fructus^[6], in cultured ICCs from murine small intestine. All four components depolarized the pacemaker potentials and decreased the amplitudes of the pacemaker potentials (Figures 6 and 7) in a concentration-dependent manner. At a concentration of 30 μ g/mL, Coptidis Rhizoma had a mean degree of depolarization of 13.1 ± 2.2 mV (Figure 6A, $n = 7$), Scutellariae Radix had a mean




Figure 6 Effects of *Coptidis Rhizoma* and *Scutellariae Radix* on pacemaker potentials in cultured interstitial cells of Cajal from murine small intestine. **A:** Pacemaking activities of ICCs exposed to *Coptidis Rhizoma* (30 µg/mL) in current-clamp mode ($I = 0$); **B:** *Coptidis Rhizoma* (10–50 µg/mL) concentration-dependently induced pacemaker potential depolarization. Responses to *Coptidis Rhizoma* are summarized; **C:** Pacemaking activities of ICCs exposed to *Scutellariae Radix* (30 µg/mL) in the current-clamp mode ($I = 0$); **D:** *Scutellariae Radix* (10–100 µg/mL) concentration-dependently induced pacemaker potential depolarization. Responses to *Scutellariae Radix* are summarized; **E:** Frequency changes to *Coptidis Rhizoma* and *Scutellariae Radix* are summarized. Bars represent the means \pm SEMs. ^b $P < 0.01$ and ^e $P < 0.001$: Significantly different from the control. CTRL: Control; CR: *Coptidis Rhizoma*; SR: *Scutellariae Radix*; ICCs: Interstitial cells of Cajal.

degree of depolarization of 0.6 ± 0.4 mV (Figure 6C, $n = 7$), *Phellodendri Cortex* had a mean degree of depolarization of 0.7 ± 0.6 mV (Figure 7A, $n = 6$) and *Gardeniae Fructus* had a mean degree of depolarization of 9.7 ± 1.5 mV (Figure 7C, $n = 7$). The summarized values and bar graph showing the effects of these components on pacemaker potentials are provided in Figures 6B, 6D, 6E, 7B, 7D and 7E. These results suggest that *Coptidis Rhizoma* and *Gardeniae Fructus* are the main components responsible for HHT-induced pacemaker potential depolarizations in the cultured ICCs of murine small intestine.

DISCUSSION

ICCs are located within and around smooth muscle layers of the GI tract and they control GI motility by acting as pacemaker cells^[20–22,33,34]. The pacemaker functions of these cells result from the electrical slow waves generated and propagated in smooth muscles^[20,35]. Additionally, ICCs are connected to both nerve and smooth muscle cells and they are involved in mechanosensation and neurotransmission^[32,34,36]. Many GI motility disorders, such as achalasia and inflammatory bowel disease, have been associated with ICC dysfunctions^[37]. Therefore, understanding the mechanisms of ICC and pacemaker activity is very

important. In the present study, we show that HHT depolarized ICC pacemaker potentials through 5-HT₃ and 5-HT₄ receptors *via* external and internal Ca²⁺ regulation and *via* G protein-, PKC- and Rho kinase-independent pathways. Additionally, we tested the effects of HHT on GI motility in mice *in vivo* and found that HHT increased the GI motility in mice^[26].

HHT has long been used as an herbal medicine to treat GI disorders in Asia^[16–18]. Ohta *et al.*^[16] suggested that oral HHT prevents stress-induced acute gastric lesions in rats. Additionally, Zhou and Mineshita^[17] reported that HHT showed anti-inflammatory and anti-ulcerative effects on trinitrobenzene-sulfonic acid (TNB)-induced colitis in rats. In that study, HHT was found to inhibit myeloperoxidase activity and to significantly decrease the IL-8, PGE2 and LTB4 levels. Therefore, the authors suggested that HHT may be useful for treating inflammatory bowel disease. In addition, Watanabe-Fukuda *et al.*^[18] observed that HHT prevented lethal indomethacin (IND)-induced enteropathy by downregulating adenosine deaminase (ADA) and thus upregulating adenosine (an anti-inflammatory nucleoside) levels. Furthermore, these authors reported that berberine, which is a major ingredient of HHT, suppressed ADA expression and reduced the incidence of lethality.

In the present study, we found that 5-HT₃ or 5-HT₄




Figure 7 Effects of Phellodendri Cortex and Gardeniae Fructus on pacemaker potentials in cultured interstitial cells of Cajal from murine small intestine. A: Pacemaking activities of ICCs exposed to Phellodendri Cortex (30 µg/mL) in the current-clamp mode ($I = 0$); B: Phellodendri Cortex (10-100 µg/mL) concentration-dependently induced pacemaker potential depolarization. Responses to Phellodendri Cortex are summarized; C: Pacemaking activities of ICCs exposed to Gardeniae Fructus (30 µg/mL) in the current-clamp mode ($I = 0$); D: Gardeniae Fructus (10-100 µg/mL) concentration-dependently induced pacemaker potential depolarizations. Responses to Gardeniae Fructus are summarized; E: Frequency changes to Phellodendri Cortex and Gardeniae Fructus are summarized. Bars represent the means \pm SEMs. ^a $P < 0.05$ and ^e $P < 0.001$: Significantly different from the control. CTRL: Control; PC: Phellodendri Cortex. GF: Gardeniae Fructus.

receptors, *via* external and internal Ca^{2+} regulation and *via* G protein, were involved in HHTE-induced responses in ICCs. Of the seven 5-HT receptor subtypes, the 5-HT₃ and 5-HT₄ receptors are the most highly expressed and best understood 5-HT receptor subtypes in the GI tract^[38,39]. 5-HT₃ receptors are ligand-gated cation channels that can initiate peristalsis and secretion in the GI tract^[38]. Additionally, 5-HT₄ receptors are G-protein-coupled metabotropic receptors and their activation regulates GI motility^[39]. Furthermore, these receptors are coupled to G proteins, which have Ca^{2+} /calmodulin-dependent activation^[40,41]. Additionally, in the present study, HHT-induced pacemaker potential depolarizations *via* 5-HT₃ and 5-HT₄ receptors were not inhibited by the Rho kinase inhibitor, Y27632, or the PKC inhibitor, calphostin (Figure 5). These results suggest that HHT-induced pacemaker potential depolarizations are independent of PKC and Rho kinase activation.

5-HT has played a major role in gut function regulation^[42]. Among 5-HT receptors, 5-HT₃ and 5-HT₄ receptors have been the most widely investigated for gut motility^[27]. 5-HT₃ receptors are mainly targeted to treat diarrhea and abdominal pain^[27,43,44] and 5-HT₄ receptors are targeted to treat constipation^[44,45]. Although each agonist and antagonist has been

used effectively, pharmacological agents of 5-HT receptors have problems in their clinical use because of deleterious side effects. In the present study, HHTE affects ICC pacemaker potentials through 5-HT₃ and 5-HT₄ receptors; therefore, HHTE may be used to treat diarrhea, abdominal pain and constipation, among other symptoms. Of note, future studies will have to be designed to determine the side effects of HHT.

The 5-HT₃ receptor channels are the only ionotropic receptors within the 5-HT receptor family^[46-48]. 5-HT₃ receptors are involved in many pathophysiological processes, such as GI motility disorders and the development of nausea and vomiting^[48-50]. Therefore, 5-HT₃ receptors play an important role in GI motility functions. In this paper, we found that HHTE may activate 5-HT₃ and 5-HT₄ receptors; among them, 5-HT₄ receptors may have the main role in regulating the ICC, whereas 5-HT₃ receptors may have a supportive role. Additionally, when GDP- β -S (1 mmol/L) was applied in the pipette solution, we found that HHTE caused slight pacemaker potential depolarization (Figure 3), which might be due to the 5-HT₃ receptor signaling pathway.

GI disorders are observed in Western and Eastern countries^[51]. Because GI disorders are not lethal, the medications used to treat these disorders have a higher safety threshold than those for medications

used to treat non-GI diseases. However, recent studies have highlighted the substantial health impacts of GI disorders^[52]. Furthermore, several GI treatments have significant adverse effects, which has led to the withdrawal of cisapride and tegaserod from the market^[53,54]. Of note, herbal products are often used to treat GI disorders like irritable bowel syndrome^[55,56] and herbal medications have a desirable role in the treatment of GI disorders.

HHT consists of *Coptidis Rhizoma*, *Scutellariae Radix*, *Phellodendri Cortex* and *Gardeniae Fructus*^[6]. We investigated the effects of these four constituents on the pacemaker potentials of ICCs to identify the components that are primarily responsible for the effect of HHT on ICC pacemaker potentials. All four components depolarized pacemaker potentials and decreased the amplitudes of pacemaker potentials in a concentration-dependent manner (Figures 6 and 7). However, *Coptidis Rhizoma* and *Gardeniae Fructus* were observed to have greater depolarizing effects. We will perform future experiments to determine the detailed effects of the HHT active elements (*Coptidis Rhizoma* and *Gardeniae Fructus*) and identified compounds (geniposide, berberine chloride, baicalin and wogonin) on ICC pacemaking potential.

ICCs are pacemaker cells that appear to play important roles in the determination and regulation of GI motility^[19,20]. HHT depolarized the pacemaker potentials of ICCs in a G protein- and Ca²⁺-dependent manner. Based on the findings described in this study, we propose the following model of the effects of HHT in ICCs. HHT binds 5-HT₃ and 5-HT₄ receptors and HHT-induced pacemaker potential depolarizations are G protein- and Ca²⁺-dependent in ICCs. An increase in intracellular Ca²⁺ induces membrane depolarizations on ICCs^[57,58]. In addition, these pacemaker potential depolarizations might be regulated by transient receptor potential melastatin 7 (TRPM7) and Ca²⁺-activated Cl⁻ (ANO1) channels^[20,59].

In the present study, HHT was found to regulate ICC pacemaker potential depolarization in a G protein- and Ca²⁺-dependent manner. Accordingly, HHT may play an excitatory role in GI motility regulation. Additionally, because GI motility regulations are highly integrated behaviors between smooth muscle cells, ICCs and enteric nervous systems, further studies are needed to investigate the effects of HHT on other cells *in vitro* and *in vivo*.

In summary, our findings suggest that HHT can depolarize the pacemaker potentials of cultured ICCs from mouse small intestine. Future research should attempt to identify the components responsible for the effects of HHT and to determine their mechanisms of action *in vitro* and *in vivo*.

COMMENTS

Background

Interstitial cells of Cajal (ICCs) are the pacemaker cells that generate slow

waves in the gastrointestinal (GI) tract. Hwangryunhaedok-tang (HHT) (consisting of *Coptidis Rhizoma*, *Scutellariae Radix*, *Phellodendri Cortex* and *Gardeniae Fructus*) is a traditional herbal medicine used to treat GI disorders. However, despite the considerable use of HHT in traditional herbal medicine to treat GI dysfunction, little is known of the effects of a water extract of HHT (HHTe) on the pacemaker potentials of mouse small intestinal ICCs.

Research frontiers

The prokinetic effects of HHTe are mediated by the induction of pacemaker potentials in ICCs.

Innovations and breakthroughs

HHTe dose-dependently depolarizes ICC pacemaker potentials through the 5-HT₃ and 5-HT₄ receptors *via* external and internal Ca²⁺ regulation as well as *via* G protein-, PKC- and Rho kinase-independent pathways.

Applications

HHT may be a new target for pharmacological treatment of GI motility disorders.

Peer-review

The authors investigated the effects of HHT (a traditional herbal medicine) on the pacemaker potentials of mouse ICCs, the results of study suggest that HHT dose-dependently depolarizes ICC pacemaker potentials through 5-HT₃ and 5-HT₄ receptors *via* external and internal Ca²⁺ regulation and *via* G protein- and MLCK-dependent pathways. These data are somewhat interesting.

REFERENCES

- Zhang J, Wider B, Shang H, Li X, Ernst E. Quality of herbal medicines: challenges and solutions. *Complement Ther Med* 2012; **20**: 100-106 [PMID: 22305255 DOI: 10.1016/j.ctim.2011.09.004]
- Normile D. Asian medicine. The new face of traditional Chinese medicine. *Science* 2003; **299**: 188-190 [PMID: 12522228 DOI: 10.1126/science.299.5604.188]
- Xue T, Roy R. Studying traditional Chinese medicine. *Science* 2003; **300**: 740-741 [PMID: 12730581 DOI: 10.1126/science.300.5620.740]
- Jiang WY. Therapeutic wisdom in traditional Chinese medicine: a perspective from modern science. *Trends Pharmacol Sci* 2005; **26**: 558-563 [PMID: 16185775 DOI: 10.1016/j.tips.2005.09.006]
- Liu S, Yi LZ, Liang YZ. Traditional Chinese medicine and separation science. *J Sep Sci* 2008; **31**: 2113-2137 [PMID: 18615809 DOI: 10.1002/jssc.200800134]
- Lu T, Song J, Huang F, Deng Y, Xie L, Wang G, Liu X. Comparative pharmacokinetics of baicalin after oral administration of pure baicalin, *Radix scutellariae* extract and Huang-Lian-Jie-Du-Tang to rats. *J Ethnopharmacol* 2007; **110**: 412-418 [PMID: 17110066 DOI: 10.1016/j.jep.2006.09.036]
- Lu J, Wang JS, Kong LY. Anti-inflammatory effects of Huang-Lian-Jie-Du decoction, its two fractions and four typical compounds. *J Ethnopharmacol* 2011; **134**: 911-918 [PMID: 21296144 DOI: 10.1016/j.jep.2011.01.049]
- Yue R, Zhao L, Hu Y, Jiang P, Wang S, Xiang L, Liu W, Zhang W, Liu R. Rapid-resolution liquid chromatography TOF-MS for urine metabolomic analysis of collagen-induced arthritis in rats and its applications. *J Ethnopharmacol* 2013; **145**: 465-475 [PMID: 23183090 DOI: 10.1016/j.jep.2012.11.010]
- Seo CS, Kim OS, Kim JH, Shin HK. Simultaneous quantification and antiatherosclerosis effect of the traditional Korean medicine, Hwangryunhaedok-tang. *BMC Complement Altern Med* 2015; **15**: 108 [PMID: 25880755 DOI: 10.1186/s12906-015-0632-5]
- Yu YL, Lu SS, Yu S, Liu YC, Wang P, Xie L, Wang GJ, Liu XD. Huang-lian-jie-du-decoction modulates glucagon-like peptide-1 secretion in diabetic rats. *J Ethnopharmacol* 2009; **124**: 444-449 [PMID: 19497359 DOI: 10.1016/j.jep.2009.05.027]
- Ohta Y, Kongo-Nishimura M, Hayashi T, Kishikawa T. Effect of Oren-gedoku-to (Huanglian-Jie-Du-Tang) extract on disruption of hepatic antioxidant defense systems in rats treated with

- D-galactosamine. *J Ethnopharmacol* 2004; **94**: 323-329 [PMID: 15325738 DOI: 10.1016/j.jep.2004.06.004]
- 12 **Hsu YL**, Kuo PL, Tzeng TF, Sung SC, Yen MH, Lin LT, Lin CC. Huang-lian-jie-du-tang, a traditional Chinese medicine prescription, induces cell-cycle arrest and apoptosis in human liver cancer cells in vitro and in vivo. *J Gastroenterol Hepatol* 2008; **23**: e290-e299 [PMID: 18522681 DOI: 10.1111/j.1440-1746.2008.05390.x]
 - 13 **Hwang YS**, Shin CY, Huh Y, Ryu JH. Hwangryun-Hae-Dok-tang (Huanglian-Jie-Du-Tang) extract and its constituents reduce ischemia-reperfusion brain injury and neutrophil infiltration in rats. *Life Sci* 2002; **71**: 2105-2117 [PMID: 12204769]
 - 14 **Zhang Q**, Ye YL, Yan YX, Zhang WP, Chu LS, Wei EQ, Yu YP. [Protective effects of Huanglian-Jiedu-Tang on chronic brain injury after focal cerebral ischemia in mice]. *Zhejiang Da Xue Xue Bao Yi Xue Ban* 2009; **38**: 75-80 [PMID: 19253432]
 - 15 **Yamakawa J**, Ishigaki Y, Takano F, Takahashi T, Yoshida J, Moriya J, Takata T, Tatsuno T, Sasaki K, Ohta T, Takegami T, Yoshizaki F. The Kampo medicines Orengedokuto, Bofutsushosan and Boiogito have different activities to regulate gene expressions in differentiated rat white adipocytes: comprehensive analysis of genetic profiles. *Biol Pharm Bull* 2008; **31**: 2083-2089 [PMID: 18981578]
 - 16 **Ohta Y**, Kobayashi T, Nishida K, Sasaki E, Ishiguro I. Preventive effect of Oren-gedoku-to (Huanglian-Jie-Du-Tang) extract on the development of stress-induced acute gastric mucosal lesions in rats. *J Ethnopharmacol* 1999; **67**: 377-384 [PMID: 10617076]
 - 17 **Zhou H**, Mineshita S. The effect of Oren-gedoku-to on experimental colitis in rats. *J Pharm Pharmacol* 1999; **51**: 1065-1074 [PMID: 10528991]
 - 18 **Watanabe-Fukuda Y**, Yamamoto M, Miura N, Fukutake M, Ishige A, Yamaguchi R, Nagasaki M, Saito A, Imoto S, Miyano S, Takeda J, Watanabe K. Orengedokuto and berberine improve indomethacin-induced small intestinal injury via adenosine. *J Gastroenterol* 2009; **44**: 380-389 [PMID: 19319464 DOI: 10.1007/s00535-009-0005-2]
 - 19 **Huizinga JD**, Thuneberg L, Klüppel M, Malysz J, Mikkelsen HB, Bernstein A. W/kit gene required for interstitial cells of Cajal and for intestinal pacemaker activity. *Nature* 1995; **373**: 347-349 [PMID: 7530333 DOI: 10.1038/373347a0]
 - 20 **Kim BJ**, Lim HH, Yang DK, Jun JY, Chang IY, Park CS, So I, Stanfield PR, Kim KW. Melastatin-type transient receptor potential channel 7 is required for intestinal pacemaker activity. *Gastroenterology* 2005; **129**: 1504-1517 [PMID: 16285951 DOI: 10.1053/j.gastro.2005.08.016]
 - 21 **Kim HJ**, Wie J, So I, Jung MH, Ha KT, Kim BJ. Menthol Modulates Pacemaker Potentials through TRPA1 Channels in Cultured Interstitial Cells of Cajal from Murine Small Intestine. *Cell Physiol Biochem* 2016; **38**: 1869-1882 [PMID: 27160463 DOI: 10.1159/000445549]
 - 22 **He CL**, Burgart L, Wang L, Pemberton J, Young-Fadok T, Szurszewski J, Farrugia G. Decreased interstitial cell of cajal volume in patients with slow-transit constipation. *Gastroenterology* 2000; **118**: 14-21 [PMID: 10611149]
 - 23 **He CL**, Soffer EE, Ferris CD, Walsh RM, Szurszewski JH, Farrugia G. Loss of interstitial cells of cajal and inhibitory innervation in insulin-dependent diabetes. *Gastroenterology* 2001; **121**: 427-434 [PMID: 11487552]
 - 24 **Zárate N**, Mearin F, Wang XY, Hewlett B, Huizinga JD, Malagelada JR. Severe idiopathic gastroparesis due to neuronal and interstitial cells of Cajal degeneration: pathological findings and management. *Gut* 2003; **52**: 966-970 [PMID: 12801952]
 - 25 **Gim H**, Nam JH, Lee S, Shim JH, Kim HJ, Ha KT, Kim BJ. Quercetin Inhibits Pacemaker Potentials via Nitric Oxide/cGMP-Dependent Activation and TRPM7/ANO1 Channels in Cultured Interstitial Cells of Cajal from Mouse Small Intestine. *Cell Physiol Biochem* 2015; **35**: 2422-2436 [PMID: 25967872 DOI: 10.1159/000374043]
 - 26 **Kim H**, Kim I, Lee MC, Kim HJ, Lee GS, Kim H, Kim BJ. Effects of Hwangryunhaedok-tang on gastrointestinal motility function in mice. *World J Gastroenterol* 2017; **23**: 2705-2715 [PMID: 28487607 DOI: 10.3748/wjg.v23.i15.2705]
 - 27 **Gershon MD**, Tack J. The serotonin signaling system: from basic understanding to drug development for functional GI disorders. *Gastroenterology* 2007; **132**: 397-414 [PMID: 17241888 DOI: 10.1053/j.gastro.2006.11.002]
 - 28 **Liu HN**, Ohya S, Nishizawa Y, Sawamura K, Iino S, Syed MM, Goto K, Imaizumi Y, Nakayama S. Serotonin augments gut pacemaker activity via 5-HT3 receptors. *PLoS One* 2011; **6**: e24928 [PMID: 21949791 DOI: 10.1371/journal.pone.0024928]
 - 29 **Shahi PK**, Choi S, Zuo DC, Yeum CH, Yoon PJ, Lee J, Kim YD, Park CG, Kim MY, Shin HR, Oh HJ, Jun JY. 5-hydroxytryptamine generates tonic inward currents on pacemaker activity of interstitial cells of cajal from mouse small intestine. *Korean J Physiol Pharmacol* 2011; **15**: 129-135 [PMID: 21860590 DOI: 10.4196/kjpp.2011.15.3.129]
 - 30 **Komori S**, Kawai M, Takewaki T, Ohashi H. GTP-binding protein involvement in membrane currents evoked by carbachol and histamine in guinea-pig ileal muscle. *J Physiol* 1992; **450**: 105-126 [PMID: 1432705]
 - 31 **Ogata R**, Inoue Y, Nakano H, Ito Y, Kitamura K. Oestradiol-induced relaxation of rabbit basilar artery by inhibition of voltage-dependent Ca channels through GTP-binding protein. *Br J Pharmacol* 1996; **117**: 351-359 [PMID: 8789390]
 - 32 **Lee JH**, Kim SY, Kwon YK, Kim BJ, So I. Characteristics of the cholecystokinin-induced depolarization of pacemaker activity in cultured interstitial cells of Cajal from murine small intestine. *Cell Physiol Biochem* 2013; **31**: 542-554 [PMID: 23571358 DOI: 10.1159/000350075]
 - 33 **Torihashi S**, Horisawa M, Watanabe Y. c-Kit immunoreactive interstitial cells in the human gastrointestinal tract. *J Auton Nerv Syst* 1999; **75**: 38-50 [PMID: 9935268]
 - 34 **Rumessen JJ**, Vanderwinden JM. Interstitial cells in the musculature of the gastrointestinal tract: Cajal and beyond. *Int Rev Cytol* 2003; **229**: 115-208 [PMID: 14669956]
 - 35 **Sanders KM**, Koh SD, Ro S, Ward SM. Regulation of gastrointestinal motility--insights from smooth muscle biology. *Nat Rev Gastroenterol Hepatol* 2012; **9**: 633-645 [PMID: 22965426 DOI: 10.1038/nrgastro.2012.168]
 - 36 **Sanders KM**, Ward SM. Kit mutants and gastrointestinal physiology. *J Physiol* 2007; **578**: 33-42 [PMID: 17095561 DOI: 10.1113/jphysiol.2006.122473]
 - 37 **Vanderwinden JM**, Rumessen JJ. Interstitial cells of Cajal in human gut and gastrointestinal disease. *Microsc Res Tech* 1999; **47**: 344-360 [PMID: 10602294 DOI: 10.1002/(SICI)1097-0029(19991201)47:5<344::AID-JEMT6>3.0.CO;2-1]
 - 38 **Taniyama K**, Makimoto N, Furuichi A, Sakurai-Yamashita Y, Nagase Y, Kaibara M, Kanematsu T. Functions of peripheral 5-hydroxytryptamine receptors, especially 5-hydroxytryptamine4 receptor, in gastrointestinal motility. *J Gastroenterol* 2000; **35**: 575-582 [PMID: 10955595]
 - 39 **Wouters MM**, Farrugia G, Schemann M. 5-HT receptors on interstitial cells of Cajal, smooth muscle and enteric nerves. *Neurogastroenterol Motil* 2007; **19** Suppl 2: 5-12 [PMID: 17620082 DOI: 10.1111/j.1365-2982.2007.00963.x]
 - 40 **Murthy KS**, Makhlof GM. Opioid mu, delta, and kappa receptor-induced activation of phospholipase C-beta 3 and inhibition of adenylyl cyclase is mediated by Gi2 and G(o) in smooth muscle. *Mol Pharmacol* 1996; **50**: 870-877 [PMID: 8863832]
 - 41 **Murthy KS**, Zhou H, Grider JR, Brautigan DL, Eto M, Makhlof GM. Differential signalling by muscarinic receptors in smooth muscle: m2-mediated inactivation of myosin light chain kinase via Gi3, Cdc42/Rac1 and p21-activated kinase 1 pathway, and m3-mediated MLC20 (20 kDa regulatory light chain of myosin II) phosphorylation via Rho-associated kinase/myosin phosphatase targeting subunit 1 and protein kinase C/CPI-17 pathway. *Biochem J* 2003; **374**: 145-155 [PMID: 12733988 DOI: 10.1042/BJ20021274]
 - 42 **Kendig DM**, Grider JR. Serotonin and colonic motility. *Neurogastroenterol Motil* 2015; **27**: 899-905 [PMID: 26095115 DOI: 10.1111/nmo.12617]
 - 43 **Vanuytsel T**, Tack JF, Boeckstaens GE. Treatment of abdominal

- pain in irritable bowel syndrome. *J Gastroenterol* 2014; **49**: 1193-1205 [PMID: 24845149 DOI: 10.1007/s00535-014-0966-7]
- 44 **Spiller R.** Recent advances in understanding the role of serotonin in gastrointestinal motility in functional bowel disorders: alterations in 5-HT signalling and metabolism in human disease. *Neurogastroenterol Motil* 2007; **19** Suppl 2: 25-31 [PMID: 17620085 DOI: 10.1111/j.1365-2982.2007.00965.x]
- 45 **Hoffman JM,** Tyler K, MacEachern SJ, Balemba OB, Johnson AC, Brooks EM, Zhao H, Swain GM, Moses PL, Galligan JJ, Sharkey KA, Greenwood-Van Meerveld B, Mawe GM. Activation of colonic mucosal 5-HT(4) receptors accelerates propulsive motility and inhibits visceral hypersensitivity. *Gastroenterology* 2012; **142**: 844-854.e4 [PMID: 22226658 DOI: 10.1053/j.gastro.2011.12.041]
- 46 **Derkach V,** Surprenant A, North RA. 5-HT₃ receptors are membrane ion channels. *Nature* 1989; **339**: 706-709 [PMID: 2472553 DOI: 10.1038/339706a0]
- 47 **Hannon J,** Hoyer D. Molecular biology of 5-HT receptors. *Behav Brain Res* 2008; **195**: 198-213 [PMID: 18571247 DOI: 10.1016/j.bbr.2008.03.020]
- 48 **Herbrechter R,** Ziemba PM, Hoffmann KM, Hatt H, Werner M, Gisselmann G. Identification of Glycyrrhiza as the rikkunshito constituent with the highest antagonistic potential on heterologously expressed 5-HT_{3A} receptors due to the action of flavonoids. *Front Pharmacol* 2015; **6**: 130 [PMID: 26191003 DOI: 10.3389/fphar.2015.00130]
- 49 **Gershon MD.** Review article: serotonin receptors and transporters -- roles in normal and abnormal gastrointestinal motility. *Aliment Pharmacol Ther* 2004; **20** Suppl 7: 3-14 [PMID: 15521849 DOI: 10.1111/j.1365-2036.2004.02180.x]
- 50 **Costedio MM,** Hyman N, Mawe GM. Serotonin and its role in colonic function and in gastrointestinal disorders. *Dis Colon Rectum* 2007; **50**: 376-388 [PMID: 17195902 DOI: 10.1007/s10350-006-0763-3]
- 51 **Chang L.** Review article: epidemiology and quality of life in functional gastrointestinal disorders. *Aliment Pharmacol Ther* 2004; **20** Suppl 7: 31-39 [PMID: 15521853 DOI: 10.1111/j.1365-2036.2004.02183.x]
- 52 **Talley NJ.** Functional gastrointestinal disorders as a public health problem. *Neurogastroenterol Motil* 2008; **20** Suppl 1: 121-129 [PMID: 18402649 DOI: 10.1111/j.1365-2982.2008.01097.x]
- 53 **Glessner MR,** Heller DA. Changes in related drug class utilization after market withdrawal of cisapride. *Am J Manag Care* 2002; **8**: 243-250 [PMID: 11915974]
- 54 **Pasricha PJ.** Desperately seeking serotonin... A commentary on the withdrawal of tegaserod and the state of drug development for functional and motility disorders. *Gastroenterology* 2007; **132**: 2287-2290 [PMID: 17570201 DOI: 10.1053/j.gastro.2007.04.057]
- 55 **Rawsthorne P,** Shanahan F, Cronin NC, Anton PA, Löfberg R, Bohman L, Bernstein CN. An international survey of the use and attitudes regarding alternative medicine by patients with inflammatory bowel disease. *Am J Gastroenterol* 1999; **94**: 1298-1303 [PMID: 10235210 DOI: 10.1111/j.1572-0241.1999.01080.x]
- 56 **Seeff LB,** Lindsay KL, Bacon BR, Kresina TF, Hoofnagle JH. Complementary and alternative medicine in chronic liver disease. *Hepatology* 2001; **34**: 595-603 [PMID: 11526548 DOI: 10.1053/jhep.2001.27445]
- 57 **Park KJ,** Hennig GW, Lee HT, Spencer NJ, Ward SM, Smith TK, Sanders KM. Spatial and temporal mapping of pacemaker activity in interstitial cells of Cajal in mouse ileum in situ. *Am J Physiol Cell Physiol* 2006; **290**: C1411-C1427 [PMID: 16381798 DOI: 10.1152/ajpcell.00447.2005]
- 58 **Lee HT,** Hennig GW, Park KJ, Bayguinov PO, Ward SM, Sanders KM, Smith TK. Heterogeneities in ICC Ca²⁺ activity within canine large intestine. *Gastroenterology* 2009; **136**: 2226-2236 [PMID: 19268670 DOI: 10.1053/j.gastro.2009.02.060]
- 59 **Zhu MH,** Kim TW, Ro S, Yan W, Ward SM, Koh SD, Sanders KM. A Ca(2+)-activated Cl(-) conductance in interstitial cells of Cajal linked to slow wave currents and pacemaker activity. *J Physiol* 2009; **587**: 4905-4918 [PMID: 19703958 DOI: 10.1113/jphysiol.2009.176206]

P- Reviewer: Baker S, Liu S, Xu WX **S- Editor:** Gong ZM

L- Editor: A **E- Editor:** Li D





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>



ISSN 1007-9327

