

Name of Journal: World Journal of Orthopedics

Manuscript NO: 33884

Manuscript Type: Original Article

Retrospective Study

Total joint replacement in inhibitor-positive haemophilia: Long-term outcome analysis in fifteen patients

Danielson Heidi, Lassila Riitta, Ylinen Pekka, Yrjönen Timo

Abstract

Aim: To collect data from joint replacement in inhibitor patients, evaluate haemostatic and patient outcomes, and analyse the costs.

Methods: We report our 21-year, single-centre cumulative experience of 15 joint arthroplasties in six inhibitor patients.

Results: Two low responder inhibitor patients were in the early days treated with FVIII, whereas bypassing agents were used in the rest of the high responder patients. The primary haemostatic outcome was good in 8/15, fair in 4/15 and poor in 3/15 operations.

Match Overview

- | | | |
|---|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 1 | Crossref 23 words
"Abstracts", Haemophilia, 06/17/2010 | 1% |
| 2 | Crossref 15 words
Kokkola, Arto, Pentti Sipponen, Perttu Arkkila, Heidi Danielson, and Pauli Puolakkainen. "Does the eradication of Helicobacter ..." | <1% |
| 3 | Publications 13 words
"Reports from Maggiore Hospital Describe Recent Advances in Hematology Research (Optimising musculoskeletal, Obesity, Fit ..." | <1% |
| 4 | Crossref 10 words
"ABSTRACTS OF THE XXIV CONGRESS OF THE INTERNATIONAL SOCIETY ON THROMBOSIS AND HAEMOSTASIS", Jo ..." | <1% |

找到约 456,000 条结果 (用时 0.50 秒)

Long-term results of primary total knee replacement in patients with ...

<https://www.ncbi.nlm.nih.gov/pubmed/15634817> - 翻译此页

作者: M Silva - 2005 - 被引用次数: 118 - 相关文章

Long-term results of primary total knee replacement in patients with hemophilia. ... (1) Hemophilia Treatment Center, Orthopaedic Hospital, 2400 South Flower ... Prosthetic survival was calculated with use of Kaplan-Meier survivorship analysis. ... The prevalence of infection in HIV-positive and HIV-negative patients was 17% ...

Successful Management of Total Knee Replacement in a High ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: R Dolatkah - 2013 - 被引用次数: 1 - 相关文章

2013年1月5日 - The patient had a history of high levels of factor VIII inhibitor (> 5.0 Bethesda ... It can provide long-term cost savings as it reduces the bleeding frequency ... outcomes following total knee replacement in hemophilic patients have been ... Nevertheless, approximately 20%–25% of hemophilia A patients and ...

Hip and Knee Replacement in the HIV positive patient - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: ER Anand - 2012 - 被引用次数: 1 - 相关文章

Arthroplasty, the process of joint replacement, is used to relieve pain ... early and late sepsis, and aseptic loosening in cases of long-term survivors. ... Thomason et al published results of 23 total knee arthroplasties in 15 haemophilic patients, ... Mean post-lymphocyte analysis at 6 years: TLC - 1.98, CD4 - 350 mm³, CD8 ...

Declining trends in invasive orthopedic interventions for people with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4958548/> - 翻译此页

作者: P TOBASE - 2016 - 相关文章

Scholar

About 80 results (0.09 sec)

Articles

[HTML] Randomized, prospective clinical trial of recombinant factor VIIa for secondary prophylaxis in **hemophilia patients** with inhibitors

Case law BA Konkle, LS Ebbesen, E Erhardtson... - ... of Thrombosis and ..., 2007 - Wiley Online Library

My library

... period. Table 1. **Patient** characteristics. 90 µg kg⁻¹ recombinant factor VII (rFVIIa), 270 µg kg⁻¹ rFVIIa, **Total** ... months. ‡ The orthopedic **joint** score was based on pain and examinations of **joints** (ankle, **knee**, and elbow). Median ...

Cited by 256 Related articles All 5 versions Cite Save

Any time

Since 2017

Since 2016

Since 2013

Custom range...

Early **long-term** FEIBA prophylaxis in **haemophilia A patients** with inhibitor after failing immune tolerance induction: A prospective clinical case series

CE Ettingshausen, W Kreuz - **Haemophilia**, 2010 - Wiley Online Library

... In two, retrospective studies with a combined **total** of 10 **patients**, and the median incidence of **joint** bleeding ... Pre-existing **joint** damage was present in six of those **patients**. ... documented in the

third retrospective study of seven **patients** with a median of three target **joints** each, as ...

Cited by 41 Related articles All 5 versions Cite Save

Sort by relevance

Sort by date

 include patents include citations Create alert

Iliopsoas haemorrhage in **patients** with **haemophilia**: **results** from one centre

C Balkan, K Kavakli, D Karapinar - **Haemophilia**, 2005 - Wiley Online Library

... Mean **total** dose of FVIII concentrate was calculated as 915 U kg⁻¹ (range: 380–1400 ... of the hip

and the quadriceps muscle was impaired causing the restriction of **knee** extension ... In

conclusion,

pain around the hip **joint**, femoral neuropathy and hip flexion contracture in a **patient** ...

Cited by 66 Related articles All 4 versions Cite Save

Intracranial haemorrhage in the Italian population of **haemophilia patients** with and without inhibitors

E Zanon, A Iorio, A Rocino, A Artoni, R Santoro... - ..., 2012 - Wiley Online Library

... Next article in issue: Factor IX **replacement** to cover **total knee replacement surgery** in **haemophilia**

B: a single ... **Replacement** therapy in the 74 inhibitor negative **patients** was carried out for a whole ...

is summarized in Table 4. rFVIIa was used in 14 **patients** (mean **total** dose used ...

Cited by 27 Related articles All 5 versions Cite Save

Practice patterns in **haemophilia A** therapy—global progress towards optimal care



全部

图片

地图

购物

更多

设置

工具

不限语言 ▾

时间不限 ▾

精确匹配 ▾

清除

Variation in FVIII / FIX activity in haemophilia: classification and clinical ...

<https://dspace.library.uu.nl/bitstream/handle/1874/202592/uijl.pdf?...2> - 翻译此页

作者: IEM den Uijl - 2011 - 相关文章

13. Chapter 3: **Analysis** of low frequency bleeding data: the association of **joint** Little is known about **long-term outcome in moderate haemophilia**. Chapter 7 ...

[PDF] Final Program - International Society on Thrombosis and Haemostasis

www.isth.org/resource/resmgr/PastMeetingProgramBooks/2009_Final_Program.pdf?...

For the first time since the Washington Congress in 1999, the ISTH returns to the thrombosis and hemostasis prevalent in the Asian-Pacific area. The **joint** discuss the role of molecular genetic **analysis in making a diagnosis of (VWD)**. 75 **PATIENTS UNDERGOING TOTAL HIP AND TOTAL KNEE REPLACEMENT**.

TOC - JournalTOCs

www.journaltoCs.ac.uk/index.php?action=tocs&journalISSN=1351... ▾ 翻译此页

ResultsA total of 81 **patients with haemophilia** and 243 controls were included ... Functional **analysis** was performed in plasma fibrinogen to characterize fibrin polymerization **Outcome** was documented by assessment of the annual **joint** bleeding rate ... and should not be recommended as a **long term** option for treatment.

ABSTRACTS - Wiley Online Library

onlinelibrary.wiley.com/doi/10.1111/jth.13325/pdf

2016年5月25日 - ysis of von Willebrand factor (vWF) in **patients**, we used rats as a model to ... ratio); plasma vWF multimer **analysis**; and platelet counts and plasma fibrinogen of AnXa to reverse anticoagulation for each direct fXa inhibitor. **Total** Background: New approaches to **Haemophilia A replacement** therapy.