

# World Journal of *Gastroenterology*

*World J Gastroenterol* 2017 July 28; 23(28): 5041-5256



**EDITORIAL**

- 5041** Colorectal cancer in young adults: A difficult challenge

*Campos FG*

- 5045** Precision medicine: In need of guidance and surveillance

*Lin JZ, Long JY, Wang AQ, Zheng Y, Zhao HT*

**REVIEW**

- 5051** Barrett's oesophagus: Current controversies

*Amadi C, Gatenby P*

- 5068** Gastrointestinal neuroendocrine peptides/amines in inflammatory bowel disease

*El-Salhy M, Solomon T, Hausken T, Gilja OH, Hatlebakk JG*

- 5086** Colorectal cancer screening: An updated review of the available options

*Issa IA, Noureddine M*

- 5097** Therapeutic potential of flavonoids in inflammatory bowel disease: A comprehensive review

*Salaritabar A, Darvishi B, Hadjiakhoondi F, Manayi A, Sureda A, Nabavi SF, Fitzpatrick LR, Nabavi SM, Bishayee A*

**ORIGINAL ARTICLE****Basic Study**

- 5115** Myo-inositol reduces  $\beta$ -catenin activation in colitis

*Bradford EM, Thompson CA, Goretsky T, Yang GY, Rodriguez LM, Li L, Barrett TA*

- 5127** Involvement of CRF2 signaling in enterocyte differentiation

*Ducarouge B, Pelissier-Rota M, Powell R, Buisson A, Bonaz B, Jacquier-Sarlin M*

- 5146** Bone marrow-derived monocyte infusion improves hepatic fibrosis by decreasing osteopontin, TGF- $\beta$ 1, IL-13 and oxidative stress

*de Souza VCA, Pereira TA, Teixeira VW, Carvalho H, de Castro MCAB, D'assunção CG, de Barros AF, Carvalho CL, de Lorena VMB, Costa VMA, Teixeira AAC, Figueiredo RCBQ, de Oliveira SA*

- 5158** Single amino acid mutation of SR-BI decreases infectivity of hepatitis C virus derived from cell culture in a cell culture model

*Gao R, Gao W, Xu G, Xu J, Ren H*

- 5167 Fibroblast-derived CXCL12/SDF-1 $\alpha$  promotes CXCL6 secretion and co-operatively enhances metastatic potential through the PI3K/Akt/mTOR pathway in colon cancer

*Ma JC, Sun XW, Su H, Chen Q, Guo TK, Li Y, Chen XC, Guo J, Gong ZQ, Zhao XD, Qi JB*

#### Case Control Study

- 5179 Association between *CYP24A1* polymorphisms and the risk of colonic polyps and colon cancer in a Chinese population

*Chen XQ, Mao JY, Li WB, Li J, Yang H, Qian JM, Li JN*

#### Retrospective Cohort Study

- 5187 Clinical significance of glycemic parameters on venous thromboembolism risk prediction in gastrointestinal cancer

*Guadagni F, Riondino S, Formica V, Del Monte G, Morelli AM, Lucchetti J, Spila A, D'Alessandro R, Della-Morte D, Ferroni P, Roselli M*

- 5196 Sex-dependent difference in the effect of metformin on colorectal cancer-specific mortality of diabetic colorectal cancer patients

*Park JW, Lee JH, Park YH, Park SJ, Cheon JH, Kim WH, Kim TI*

#### Retrospective Study

- 5206 Sex-influenced association of non-alcoholic fatty liver disease with colorectal adenomatous and hyperplastic polyps

*Chen QF, Zhou XD, Sun YJ, Fang DH, Zhao Q, Huang JH, Jin Y, Wu JS*

#### Observational Study

- 5216 Development and validation of a simple and multifaceted instrument, GERD-TEST, for the clinical evaluation of gastroesophageal reflux and dyspeptic symptoms

*Nakada K, Matsuhashi N, Iwakiri K, Oshio A, Joh T, Higuchi K, Haruma K*

- 5229 Modified B-ultrasound method for measurement of antral section only to assess gastric function and guide enteral nutrition in critically ill patients

*Liu Y, Gao YK, Yao L, Li L*

#### Prospective Study

- 5237 Chronic liver failure-consortium acute-on-chronic liver failure and acute decompensation scores predict mortality in Brazilian cirrhotic patients

*Picon RV, Bertol FS, Tovo CV, de Mattos ÁZ*

#### CASE REPORT

- 5246 Refractory hepatic encephalopathy in a patient with hypothyroidism: Another element in ammonia metabolism

*Díaz-Fontenla F, Castillo-Pradillo M, Díaz-Gómez A, Ibáñez-Samaniego L, Gancedo P, Guzmán-de-Villoria JA, Fernández-García P, Bañares-Cañizares R, García-Martínez R*

- 5253** Endoscopic occlusion with silicone spigots for the closure of refractory esophago-bronchiole fistula after esophagectomy

*Uesato M, Kono T, Akutsu Y, Murakami K, Kagaya A, Muto Y, Nakano A, Aikawa M, Tamachi T, Amagai H, Arasawa T, Muto Y, Matsubara H*

## Contents

*World Journal of Gastroenterology*  
Volume 23 Number 28 July 28, 2017

### ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Edoardo G Giannini, FACG, MD, PhD, Assistant Professor, Department of Internal Medicine, University of Genoa, 16132 Genova, Italy

### AIMS AND SCOPE

*World Journal of Gastroenterology* (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7<sup>th</sup>, 14<sup>th</sup>, 21<sup>st</sup>, and 28<sup>th</sup> each month. The *WJG* Editorial Board consists of 1375 experts in gastroenterology and hepatology from 68 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

### INDEXING/ABSTRACTING

*World Journal of Gastroenterology* (*WJG*) is now indexed in Current Contents<sup>®</sup>/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch<sup>®</sup>), Journal Citation Reports<sup>®</sup>, Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2017 edition of Journal Citation Reports<sup>®</sup> cites the 2016 impact factor for *WJG* as 3.365 (5-year impact factor: 3.176), ranking *WJG* as 29<sup>th</sup> among 79 journals in gastroenterology and hepatology (quartile in category Q2).

### FLYLEAF

#### I-IX Editorial Board

### EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*  
Responsible Electronic Editor: *Fen-Fen Zhang*  
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Yuan Qi*  
Proofing Editorial Office Director: *Jin-Lei Wang*

NAME OF JOURNAL  
*World Journal of Gastroenterology*

ISSN  
ISSN 1007-9327 (print)  
ISSN 2219-2840 (online)

LAUNCH DATE  
October 1, 1995

FREQUENCY  
Weekly

EDITORS-IN-CHIEF  
**Damian Garcia-Olmo, MD, PhD, Doctor, Professor, Surgeon**, Department of Surgery, Universidad Autonoma de Madrid; Department of General Surgery, Fundacion Jimenez Diaz University Hospital, Madrid 28040, Spain

**Stephen C Strom, PhD, Professor**, Department of Laboratory Medicine, Division of Pathology, Karolinska Institutet, Stockholm 141-86, Sweden

**Andrzej S Tarnawski, MD, PhD, DSc (Med), Professor of Medicine, Chief Gastroenterology**, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach,

CA 90822, United States

EDITORIAL BOARD MEMBERS  
All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE  
Jin-Lei Wang, Director  
Yuan Qi, Vice Director  
Ze-Mao Gong, Vice Director  
*World Journal of Gastroenterology*  
Baishideng Publishing Group Inc  
7901 Stoneridge Drive, Suite 501,  
Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [editorialoffice@wjgnet.com](mailto:editorialoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>

PUBLISHER  
Baishideng Publishing Group Inc  
7901 Stoneridge Drive, Suite 501,  
Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>

<http://www.wjgnet.com>

PUBLICATION DATE  
July 28, 2017

COPYRIGHT  
© 2017 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT  
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS  
Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION  
<http://www.f6publishing.com>





## Colorectal cancer in young adults: A difficult challenge

Fábio Guilherme Campos

Fábio Guilherme Campos, Division of Colorectal Surgery, Department of Gastroenterology, Hospital das Clínicas, University of São Paulo, São Paulo, SP 01411000, Brazil

**Author contributions:** Campos FG contributed to the manuscript.

**Conflict-of-interest statement:** The authors have no conflicts of interest to report.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Invited manuscript

**Correspondence to:** Dr. Fábio Guilherme Campos, Professor, Division of Colorectal Surgery, Department of Gastroenterology, Hospital das Clínicas, University of São Paulo, Rua Padre João Manoel, 222, Cj 120, Cerqueira César, São Paulo, SP 01411000, Brazil. [fgmcampos@terra.com.br](mailto:fgmcampos@terra.com.br)  
Telephone: +55-11-30610108  
Fax: +55-11-30610108

Received: March 21, 2017

Peer-review started: March 22, 2017

First decision: April 20, 2017

Revised: June 7, 2017

Accepted: July 12, 2017

Article in press: July 12, 2017

Published online: July 28, 2017

### Abstract

Sporadic colorectal cancer (CRC) is traditionally diagnosed after the sixth decade of life, and current recommendations for surveillance include only patients older than 50 years of age. However, an increasing incidence of CRC in patients less than 40 years of age has been reported. This occurrence has been attributed

to different molecular features and low suspicion of CRC in young symptomatic individuals. When confronting young-onset CRC with older patients, issues such as biological aggressiveness, stage at diagnosis and clinical outcomes seem to differ in many aspects. In the future, the identification of the molecular profile underlying the early development of sporadic CRC will help to plan tailored screening recommendations and improve management. Besides that, differential diagnosis with CRC linked with hereditary syndromes is necessary to provide adequate patient treatment and family screening. Until we find the answers to some of these doubts, doctors should raise suspicion when evaluating a young adult and be aware of this risk and consequences of a late diagnosis.

**Key words:** Colorectal cancer; Young age; Hereditary; Prognosis

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** Colorectal cancer (CRC) is traditionally considered a disease affecting people with more than 50 years of age. However, numerous researches have detected a rising incidence of CRC in young people, mainly rectal cancer. This finding raises the need for increasing clinical suspicion when evaluating symptoms of a young patient. Furthermore, these groups of patients must be aware of this possibility.

Campos FG. Colorectal cancer in young adults: A difficult challenge. *World J Gastroenterol* 2017; 23(28): 5041-5044  
Available from: URL: <http://www.wjgnet.com/1007-9327/full/v23/i28/5041.htm> DOI: <http://dx.doi.org/10.3748/wjg.v23.i28.5041>

### INTRODUCTION

Colorectal cancer (CRC) is the most common gastrointestinal neoplasia. As it is widely considered a

disease that affects people after the 5<sup>th</sup> decade of life, screening is not indicated before 50 years of age, when the risk is lower<sup>[1]</sup>. Over the past years, the increasing use of screening colonoscopy in adults aged 50-75 years has declined CRC incidence by at least 4% per year due to resection of adenomas, also improving the detection of more early lesions<sup>[2]</sup>.

The finding of CRC in adolescents or young adults has always raised attention due to issues such as the emotional impact at diagnosis, the disease behavior and the possibility to be associated with genetic diseases<sup>[3]</sup>.

However, the definition of age to be considered young is controversial. In an analysis of 6425 patients from 55 publications, O'Connell *et al*<sup>[4]</sup> found that 37 manuscripts considered young those below 40 years of age, while 14 (25%) and 4 (7%) defined as young those below 30 and 35 years of age, respectively. In the literature, most publications refer the CRC incidence in patients with less than 40 years of age<sup>[4-6]</sup>.

The incidence of CRC diagnosed before 40 years of age varies from 0.8% to 15%<sup>[5]</sup>. Series published during the last decade reveal great variation due to biases associated with experiences from one institution or reference centers. Recently, a greater incidence of young age CRC has been documented<sup>[7]</sup>. Within this group, a proper investigation to discharge hereditary forms of CRC is fundamental.

So far, age is not universally accepted as an independent prognostic factor. Otherwise, CRC diagnosis in young people is always difficult, as both patients and physicians underestimate symptoms and postpone diagnosis and management<sup>[6]</sup>.

## EPIDEMIOLOGY

CRC lifetime risk is around 5%<sup>[8]</sup>. Since 1998, the CRC overall incidence has decreased in the United States, as a probable effect of CRC screening<sup>[9]</sup>. However, in a contrast to these overall trends, this pattern is different among young adults.

Studies from *Surveillance Epidemiology and End Result* have presented important findings regarding the epidemiologic trends of CRC in different age groups. Bailey *et al*<sup>[10]</sup> reported a growing incidence rate (10%) among patients from 20 to 50 years and a 20% reduction after 50 years. Using resource data from 1987 to 2006, Davis *et al*<sup>[11]</sup> found raising incidences within patients aged 20-49 years, especially among those between 40-44 years (from 10.7 in 1988 to 17.9 per 100 thousand in 2006). From 1973 to 2005, Meyer *et al*<sup>[12]</sup> identified 7661 CRC patients with less than 40 years in a retrospective study. Age variation throughout time revealed stable rates for colon and increased rates for rectal cancer. Consequently, the authors emphasize that symptomatic young adults should be endoscopically investigated to avoid a late diagnosis, as malignant tumors in young predominates in segments distal to the splenic flexure.

Also, data from the American Cancer Society also noticed an increase in the global incidence from 1992 to 2005 among adults between 20 and 49 years, demonstrating a 3.5% increase per years among men and 2.9% per year in women<sup>[13]</sup>. More recently, a report from the National Cancer Database identified a consistent incidence increase from 1998 to 2007 by analyzing a group of young-onset (64068, 11%) and later-onset (52480, 89%) CRC<sup>[14]</sup>.

## CLINICO-PATHOLOGICAL AND GENETIC ASPECTS

There exists lots of discussion regarding the disparities related to age at CRC diagnosis and tumor biology, recurrence rates, treatment and outcomes. Most publications emphasize that the incidence of young-onset CRC has increased mainly between 40-49 years, when they are more likely to be found in the distal colon and rectum and also advanced stages at presentation.

Although controversial, young patients have been considered to have a more aggressive biological behavior and worst prognosis<sup>[15-17]</sup>. Furthermore, It has been also described a greater prevalence of mucinous and less differentiated tumors<sup>[18]</sup> within this group, characteristics also associated with bad prognosis. For these reasons, the question of considering 40 years as the basis for colonoscopic surveillance has been a constant matter of debate, mainly for men<sup>[19]</sup>.

This behavior is generally attributed to the discovery of a more advanced disease, as stages III and IV predominate among the young<sup>[14,18,20]</sup>. Our group evaluated the question if a late diagnosis could result from less diagnostic efforts in an apparent health group, and we found that symptoms duration was equal (13.8 mo vs 14.5 mo,  $P = 0.5$ ) among the young and control group<sup>[5]</sup>. Others<sup>[21]</sup> have similarly emphasized that the greater proportion of patients with advanced-stage could not be simply explained by delay in diagnosis.

These results suggest that identification of high-risk young people for screening and recognition of alert symptoms is now a real medical problem. For this, complaints such as persistent rectal bleeding, abdominal pain or anemia should require endoscopic work-ups even in average-risk young people.

One of the main challenges in a young population is to distinguish sporadic from the hereditary forms of CRC. Overall, only 2%-5% of CRC are caused by highly penetrant genes<sup>[22]</sup>, and 15%-20% of CRC in young age population are hereditary<sup>[23,24]</sup>. An additional problem is that a familiar history of CRC in almost one fifth of hereditary syndromes patients is not recognized<sup>[14,25]</sup>. Besides this, when diagnosed with a CRC, a young patient should undergo genetic tests, even in the absence of clinical phenotype or normal MSI-IHC studies<sup>[26]</sup>.

So far, knowledge regarding the molecular features of sporadic young-onset CRC is limited. It is generally thought that they derive from a cumulative effect of multiple genetic variants displaying variable penetrance. Probably, the molecular profile in young is heterogeneous and different from late-onset CRC patients<sup>[27]</sup>.

## FINAL COMMENTS AND PERSPECTIVES

Important information may be extracted from the present data. Within a young group, CRC is usually diagnosed later and potentially associated with worst prognosis. Thus, a greater suspicion rate is necessary when evaluating young patients with common symptoms. Moreover, educational and preventive programs should provide adequate information regarding alert symptoms and risk populations.

The recognition that CRC incidence at an early age may be correlated with modern dietary factors and epidemic obesity requires improvement in medical awareness and population information about these risk factors<sup>[28]</sup>. Moreover, young people may develop sporadic cancers that are not usually detected through current screening programs, suggesting a possible drawback in the CRC screening recommendations adopted so far. This fact raised the idea to lower the screening age of those not associated with a high risk (although there is no consensus about it so far), even though the impact of such attitude on early detection, costs and survival has not been fully appreciated yet<sup>[29]</sup>. In addition, there are lots of barriers to overcome in terms of adherence to preventive colonoscopy at a young age.

In terms of colonoscopic surveillance for risk groups, both patients and physicians should be aware about the importance of establishing a family pedigree. This is based on previous knowledge and investigations demonstrating that the existence of a young family member or a first-degree relative with CRC is risk factors for advanced lesions (including cancer)<sup>[30]</sup>.

Eventually, young age may affect therapeutic decisions, as patients with sporadic cancers before 50 years have been considered for a subtotal colectomy, although this decision not always translates into a greater survival<sup>[31]</sup>. Otherwise, the recognition of a hereditary syndrome would certainly support the indication of a total colectomy and suggest familial surveillance, besides the absence of prospective randomized trials comparing extended and segmental resections<sup>[32]</sup>. This tendency correlates with the fact that the presence of mismatch repair deficiency raises the risk of subsequent metachronous neoplasia<sup>[33]</sup>.

Aside from those involved in Lynch Syndrome, many other genes (APC, KRAS, P53, BRAF) may influence tumor genesis and biology. In this context, mutations in FBXW7 and POLE genes were found to prevail in younger patients with CRC<sup>[34]</sup>. Thus, the most crucial challenge for the future is the understanding of how

genetic profile may affect CRC incidence and outcomes of treatment interventions in different age cohorts.

In the absence of genetic information, however, individual decisions should be taken on the basis of health status, family history, opportunity to undergo postoperative follow-up, quality of anal sphincters and patient consent information.

## REFERENCES

- 1 **Brenner H**, Altenhofen L, Hoffmeister M. Sex, age, and birth cohort effects in colorectal neoplasms: a cohort analysis. *Ann Intern Med* 2010; **152**: 697-703 [PMID: 20513827 DOI: 10.7326/0003-4819-152-11-201006010-00002]
- 2 **Siegel RL**, Miller KD, Jemal A. Cancer statistics, 2015. *CA Cancer J Clin* 2015; **65**: 5-29 [PMID: 25559415 DOI: 10.3322/caac.21254]
- 3 **Eliezer D**, Hadley DW, Koehly LM. Exploring psychological responses to genetic testing for Lynch Syndrome within the family context. *Psychooncology* 2014; **23**: 1292-1299 [DOI: 10.1002/pon.3551]
- 4 **O'Connell JB**, Maggard MA, Liu JH, Etzioni DA, Livingston EH, Ko CY. Do young colon cancer patients have worse outcomes? *World J Surg* 2004; **28**: 558-562 [PMID: 15366745 DOI: 10.1007/s00268-004-7306-7]
- 5 **Lupinacci RM**, Campos FG, Araújo SE, Imperiale AR, Seid VE, Habr-Gama A, Kiss DR, Gama-Rodrigues J. Análise comparativa das características clínicas, anátomo-patológicas e sobrevida entre pacientes com câncer colo-retal abaixo e acima de 40 anos de idade. *Rev. bras. Coloproct* 2003; **23**: 155-162
- 6 **Fazeli MS**, Adel MG, Lebaschi AH. Colorectal carcinoma: a retrospective, descriptive study of age, gender, subsite, stage, and differentiation in Iran from 1995 to 2001 as observed in Tehran University. *Dis Colon Rectum* 2007; **50**: 990-995 [PMID: 17525859 DOI: 10.1007/s10350-007-0248-z]
- 7 **Taggash D**, Rehil N, Sharma S, Flynn JC, Damadi A. Colorectal cancer: are the "young" being overlooked? *Am J Surg* 2013; **205**: 312-316; discussion 316 [PMID: 23414955 DOI: 10.1016/j.amjsurg.2012.10.016]
- 8 **Sag AA**, Selcukbiricik F, Mandel NM. Evidence -based medical oncology and interventional radiology paradigms for liver dominant colorectal cancer metastases. *World J Gastroenterol* 2016; **22**: 3127-3149 [PMID: 27003990 DOI: 10.3748/wjg.v22.i11.3127]
- 9 **Edwards BK**, Ward E, Kohler BA, Ehemann C, Zaubler AG, Anderson RN, Jemal A, Schymura MJ, Lansdorp-Vogelaar I, Seeff LC, van Ballegooijen M, Goede SL, Ries LA. Annual report to the nation on the status of cancer, 1975-2006, featuring colorectal cancer trends and impact of interventions (risk factors, screening, and treatment) to reduce future rates. *Cancer* 2010; **116**: 544-573 [PMID: 19998273 DOI: 10.1002/cncr.24760]
- 10 **Bailey CE**, Hu CY, You YN, Bednarski BK, Rodriguez-Bigas MA, Skibber JM, Cantor SB, Chang GJ. Increasing disparities in the age-related incidences of colon and rectal cancers in the United States, 1975-2010. *JAMA Surg* 2015; **150**: 17-22 [PMID: 25372703 DOI: 10.1001/jamasurg.2014.1756]
- 11 **Davis DM**, Marcet JE, Frattini JC, Prather AD, Mateka JJ, Nfonsam VN. Is it time to lower the recommended screening age for colorectal cancer? *J Am Coll Surg* 2011; **213**: 352-361 [PMID: 21737316 DOI: 10.1016/j.jamcollsurg.2011.04.033]
- 12 **Meyer JE**, Narang T, Schnoll-Sussman FH, Pochapin MB, Christos PJ, Sherr DL. Increasing incidence of rectal cancer in patients aged younger than 40 years: an analysis of the surveillance, epidemiology, and end results database. *Cancer* 2010; **116**: 4354-4359 [PMID: 20734460 DOI: 10.1002/cncr.25432]
- 13 **Siegel RL**, Jemal A, Ward EM. Increase in incidence of colorectal cancer among young men and women in the United States. *Cancer Epidemiol Biomarkers Prev* 2009; **18**: 1695-1698 [PMID:



- 19505901 DOI: 10.1158/1055-9965.EPI-09-0186]
- 14 **You YN**, Xing Y, Feig BW, Chang GJ, Cormier JN. Young-onset colorectal cancer: is it time to pay attention? *Arch Intern Med* 2012; **172**: 287-289 [PMID: 22157065 DOI: 10.1001/archinternmed.2011.602]
  - 15 **Fancher TT**, Palesty JA, Rashidi L, Dudrick SJ. Is gender related to the stage of colorectal cancer at initial presentation in young patients? *J Surg Res* 2011; **165**: 15-18 [PMID: 21074787 DOI: 10.1016/j.jss.2010.08.054]
  - 16 **Al-Barrak J**, Gill S. Presentation and outcomes of patients aged 30 years and younger with colorectal cancer: a 20-year retrospective review. *Med Oncol* 2011; **28**: 1058-1061 [PMID: 20680521 DOI: 10.1007/s12032-010-9639-4]
  - 17 **da Fonseca LM**, da Luz MM, Lacerda-Filho A, Cabral MM, da Silva RG. Colorectal carcinoma in different age groups: a histopathological analysis. *Int J Colorectal Dis* 2012; **27**: 249-255 [PMID: 21845417 DOI: 10.1007/s00384-011-1299-0]
  - 18 **Chou CL**, Chang SC, Lin TC, Chen WS, Jiang JK, Wang HS, Yang SH, Liang WY, Lin JK. Differences in clinicopathological characteristics of colorectal cancer between younger and elderly patients: an analysis of 322 patients from a single institution. *Am J Surg* 2011; **202**: 574-582 [PMID: 21872205 DOI: 10.1016/j.amjsurg.2010.10.014]
  - 19 **Kolligs FT**, Crispin A, Munte A, Wagner A, Mansmann U, Göke B. Risk of advanced colorectal neoplasia according to age and gender. *PLoS One* 2011; **6**: e20076 [PMID: 21629650 DOI: 10.1371/journal.pone.0020076]
  - 20 **Du CZ**, Zhang JS, Li M, Zhao J, Peng YF, Yao YF, Xue WC, Gu J. [Comparison of pathologic and clinical characteristics of young and old patients with advanced rectal cancer after neoadjuvant radiotherapy]. *Zhonghua Waike Zazhi* 2010; **48**: 1616-1620 [PMID: 21211254]
  - 21 **Chen FW**, Sundaram V, Chew TA, Ladabaum U. Advanced-Stage Colorectal Cancer in Persons Younger Than 50 Years Not Associated With Longer Duration of Symptoms or Time to Diagnosis. *Clin Gastroenterol Hepatol* 2017; **15**: 728-737.e3 [PMID: 27856366 DOI: 10.1016/j.cgh.2016.10.038]
  - 22 **Jasperson KW**, Tuohy TM, Neklason DW, Burt RW. Hereditary and familial colon cancer. *Gastroenterology* 2010; **138**: 2044-2058 [PMID: 20420945 DOI: 10.1053/j.gastro.2010.01.054]
  - 23 **Chang DT**, Pai RK, Rybicki LA, Dimaio MA, Limaye M, Jayachandran P, Koong AC, Kunz PA, Fisher GA, Ford JM, Welton M, Shelton A, Ma L, Arber DA, Pai RK. Clinicopathologic and molecular features of sporadic early-onset colorectal adenocarcinoma: an adenocarcinoma with frequent signet ring cell differentiation, rectal and sigmoid involvement, and adverse morphologic features. *Mod Pathol* 2012; **25**: 1128-1139 [PMID: 22481281 DOI: 10.1038/modpathol.2012.61]
  - 24 **Silla IO**, Rueda D, Rodríguez Y, García JL, de la Cruz Vigo F, Perea J. Early-onset colorectal cancer: a separate subset of colorectal cancer. *World J Gastroenterol* 2014; **20**: 17288-17296 [PMID: 25516639 DOI: 10.3748/wjg.v20.i46.17288]
  - 25 **Mork ME**, You YN, Ying J, Bannon SA, Lynch PM, Rodriguez-Bigas MA, Vilar E. High Prevalence of Hereditary Cancer Syndromes in Adolescents and Young Adults With Colorectal Cancer. *J Clin Oncol* 2015; **33**: 3544-3549 [PMID: 26195711 DOI: 10.1200/JCO.2015.61.4503]
  - 26 **Lynch PM**. How helpful is age at colorectal cancer onset in finding hereditary nonpolyposis colorectal cancer? *Clin Gastroenterol Hepatol* 2011; **9**: 458-460 [PMID: 21440092 DOI: 10.1016/j.cgh.2011.03.016]
  - 27 **Ballester V**, Rashtak S, Boardman L. Clinical and molecular features of young-onset colorectal cancer. *World J Gastroenterol* 2016; **22**: 1736-1744 [PMID: 26855533 DOI: 10.3748/wjg.v22.i5.1736]
  - 28 **Hubbard JM**, Grothey A. Adolescent and young adult colorectal cancer. *J Natl Compr Canc Netw* 2013; **11**: 1219-1225 [PMID: 24142823]
  - 29 **Deen KI**, Silva H, Deen R, Chandrasinghe PC. Colorectal cancer in the young, many questions, few answers. *World J Gastrointest Oncol* 2016; **8**: 481-488 [PMID: 27326317 DOI: 10.4251/wjgo.v8.i6.481]
  - 30 **Forsberg AM**, Hagel E, Jaramillo E, Rubio CA, Björck E, Lindblom A. Predicting Outcome in Colonoscopic High-risk Surveillance. *Anticancer Res* 2015; **35**: 4813-4819 [PMID: 26254372]
  - 31 **Klos CL**, Montenegro G, Jamal N, Wise PE, Fleshman JW, Safar B, Dharmarajan S. Segmental versus extended resection for sporadic colorectal cancer in young patients. *J Surg Oncol* 2014; **110**: 328-332 [PMID: 24888987 DOI: 10.1002/jso.23649]
  - 32 **Herzig DO**, Buie WD, Weiser MR, You YN, Rafferty JF, Feingold D, Steele SR. Clinical Practice Guidelines for the Surgical Treatment of Patients With Lynch Syndrome. *Dis Colon Rectum* 2017; **60**: 137-143 [PMID: 28059909 DOI: 10.1097/DCR.0000000000000785]
  - 33 **Aronson M**, Holter S, Semotiuk K, Winter L, Pollett A, Gallinger S, Cohen Z, Gryfe R. DNA Mismatch Repair Status Predicts Need for Future Colorectal Surgery for Metachronous Neoplasms in Young Individuals Undergoing Colorectal Cancer Resection. *Dis Colon Rectum* 2015; **58**: 645-652 [PMID: 26200678 DOI: 10.1097/DCR.0000000000000391]
  - 34 **Kothari N**, Teer JK, Abbott AM, Srikumar T, Zhang Y, Yoder SJ, Brohl AS6, Kim RD, Reed DR, Shibata D. Increased incidence of FBXW7 and POLE proofreading domain mutations in Young adult colorectal cancers. *Cancer* 2016; **122**: 2828-2835 [PMID: 27244218 DOI: 10.1002/cncr.30082]

**P-Reviewer:** Milone M, Lakatos PL, Sag AA **S-Editor:** Qi Y  
**L-Editor:** A **E-Editor:** Zhang FF





Published by **Baishideng Publishing Group Inc**  
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA  
Telephone: +1-925-223-8242  
Fax: +1-925-223-8243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>



ISSN 1007-9327

