

Colonoscopy procedural volume increases adenoma and polyp detection rates

全部 图片 视频 新闻 地图 图书

找到约 59,000 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[Leadership training to improve adenoma detection rate in screening ...](#)

gut.bmj.com/content/early/2015/02/09/gutjnl-2014-307503

10 Feb 2015 ... Colonoscopies performed June to December 2012 (early ... Suboptimal adenoma detection at colonoscopy is associated with increased risk of interval ... as involvement of a trainee or video recording of the colonoscopy, 8–11 or an ... Reliability of adenoma detection rate is based on procedural volume.

[\[PDF\] Leadership training to improve adenoma detection rate in ... - Gut](#)

gut.bmj.com/content/gutjnl/early/2015/02/.../gutjnl-2014-307503.full.pdf

10 Feb 2015 ... ABSTRACT. Objective Suboptimal adenoma detection rate (ADR) at colonoscopy is associated with increased risk of interval colorectal cancer.

[DETECTION RATES OF PREMALIGNANT POLYPS DURING ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4340745/>

10 Jan 2015 ... Standards for detection of adenomas during screening colonoscopy are ... Detection rates of both types of premalignant polyps improved over time This represents approximately 18% of colonoscopy volume in our Both overall ADR and sessile serrated adenoma detection rates increased over time for ...

[Association of trainee participation with adenoma and polyp ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5437386/>

16 May 2017 ... The ADR and PDR increased with increasing fellow training level (P for trend < 0.05). ... Keywords: Screening colonoscopy, Adenoma detection rate, Polyp ... quality of the procedure, but rather increases adenoma and polyp detection. ... The primary aim of our study was to investigate whether GI fellows at ...

[Screening and Surveillance of the Early Detection of Colorectal ...](#)

<https://gi.org/.../screening-and-surveillance-of-the-early-detection-of-colorectal-cancer-and-adenomatous-polyps/>

The most common and clinically important polyps are adenomatous polyps, which ... adults using different tests, with FSIG rates declining, colonoscopy rates increasing. When either the test, the testing procedure, or both have very low test In a recent retrospective

Name of Journal: World Journal of Gastroenterology

Manuscript NO: 34973

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Colonoscopy procedural volume increases adenoma and polyp detection rates in GI trainees

Emad Qayed, Ravi Vora, Sara Levy, Roberd M Bostick

Abstract

AIM

To investigate changes in polyp detection throughout fellowship training, and estimate colonoscopy volume required to achieve the ADRs and PDRs of attending gastroenterologists.

METHODS

We reviewed colonoscopies from July 1, 2009 to June 30, 2014. Fellows' procedural logs

Match Overview

1 Internet 51 words
crawled on 18-Jun-2017
www.wjgnet.com

1%

Colonoscopy procedural volume increases adenoma and polyp detection rates

全部 图片 视频 新闻 地图 图书

找到约 27,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Google 学术: Colonoscopy procedural volume increases adenoma and polyp detection rates in GI trainees](#)

[... and adenoma detection during screening colonoscopy](#) - Barclay - 被引用次数: 1079

[... on polyp yield: implications for optimal colonoscopy ...](#) - Simmons - 被引用次数: 241

[Adenoma detection rate and risk of colorectal cancer ...](#) - Corley - 被引用次数: 467

[Leadership training to improve adenoma detection rate in screening ...](#)

[gut.bmj.com/content/early/2015/02/09/gutjnl-2014-307503](#)

10 Feb 2015 ... Colonoscopies performed June to December 2012 (early ... Suboptimal adenoma detection at colonoscopy is associated with increased risk of interval ... as involvement of a trainee or video recording of the colonoscopy, 8–11 or an ... Reliability of adenoma detection rate is based on procedural volume.

[\[PDF\] Expert opinions and scientific evidence for colonoscopy key ... - Gut](#)

[gut.bmj.com/content/gutjnl/early/2016/10/.../gutjnl-2016-312043.full.pdf](#)

8 Oct 2016 ... procedural volumes increasing annually throughout the world. ... patients with lower GI symptoms and is an integral ... ation and adenoma detection rates (ADR) and these ... tandem (same day) colonoscopies. 61 Miss rates for all polyps ... with 140 for colorectal surgery and gastroenterology trainees in.

[Level of Fellowship Training Increases Adenoma Detection Rates ...](#)

[www.cghjournal.org/article/S1542-3565\(10\)00075-3/fulltext](#)

1 Feb 2010 ... Similarly, the polyp detection rate was higher (OR, 1.28; 95% CI, 1.08–1.52) among ... Involvement of fellows in screening colonoscopies increases the ADR, primarily ... of a gastroenterology (GI) fellow enhances polyp or adenoma detection. ... higher ADR when a fellow was involved with the procedure.

[\[PDF\] Level of Fellowship Training Increases Adenoma Detection Rates](#)

[www.cghjournal.org/article/S1542-3565\(10\)00075-3/pdf](#)

lary, the polyp detection rate was higher (OR, 1.28; 95% CI, 1.08–1.52) ... GI fellow participation during colonoscopy affects the ADR. This study ... was involved with the procedure. However ... or high volume (more than 300 screening colonoscopies per ... slower

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 26,700 条结果 (用时 0.37 秒)

Google 学术 : Colonoscopy procedural volume increases adenoma and polyp detection rates in GI trainees

... and adenoma detection during screening colonoscopy - Barclay - 被引用次数 : 1083

... on polyp yield: implications for optimal colonoscopy ... - Simmons - 被引用次数 : 242

Adenoma detection rate and risk of colorectal cancer ... - Corley - 被引用次数 : 470

Level of Fellowship Training Increases Adenoma Detection Rates

[www.cghjournal.org/article/S1542-3565\(10\)00075-3/pdf](http://www.cghjournal.org/article/S1542-3565(10)00075-3/pdf) - 翻译此页

作者 : SL Peters - 2010 - 被引用次数 : 48 - 相关文章

larly, the polyp detection rate was higher (OR, 1.28; 95% CI, 1.08–1.52) ... GI fellow participation during colonoscopy affects the ADR.8. This study ... was involved with the procedure. However ... or high volume (more than 300 screening colonoscopies per- slower in first-year trainees, compared with third-year fellows,.

Level of Fellowship Training Increases Adenoma Detection Rates ...

[www.cghjournal.org/article/S1542-3565\(10\)00075-3/fulltext](http://www.cghjournal.org/article/S1542-3565(10)00075-3/fulltext) - 翻译此页

作者 : SL Peters - 2010 - 被引用次数 : 48 - 相关文章

2010年2月1日 - Similarly, the polyp detection rate was higher (OR, 1.28; 95% CI, 1.08–1.52) among ... Involvement of fellows in screening colonoscopies increases the ADR, primarily ... of a gastroenterology (GI) fellow enhances polyp or adenoma detection. ... higher ADR when a fellow was involved with the procedure.

Leadership training to improve adenoma detection rate in screening ...

gut.bmj.com > Online First - 翻译此页

作者 : MF Kaminski - 2015 - 被引用次数 : 38 - 相关文章

2015年2月10日 - Colonoscopies performed June to December 2012 (early ... Suboptimal adenoma detection at colonoscopy is associated with increased risk of interval ... as involvement of a trainee or video recording of the colonoscopy,8–11 or an Reliability of adenoma detection rate is based on procedural volume.

What is the criterion for high-performing colonoscopists? Being