

# World Journal of *Gastroenterology*

*World J Gastroenterol* 2017 October 7; 23(37): 6747-6922



**EDITORIAL**

- 6747 Microbial dysbiosis in spouses of ulcerative colitis patients: Any clues to disease pathogenesis?  
*Sorrentino D*

**REVIEW**

- 6750 Road to stemness in hepatocellular carcinoma  
*Flores-Téllez TNJ, Villa-Treviño S, Piña-Vázquez C*
- 6777 Intrahepatic vascular changes in non-alcoholic fatty liver disease: Potential role of insulin-resistance and endothelial dysfunction  
*Pasarin M, Abalde JG, Liguori E, Kok B, La Mura V*
- 6788 Epidemiological and clinical perspectives on irritable bowel syndrome in India, Bangladesh and Malaysia: A review  
*Rahman MM, Mahadeva S, Ghoshal UC*

**ORIGINAL ARTICLE****Basic Study**

- 6802 Estrogen receptor expression in chronic hepatitis C and hepatocellular carcinoma pathogenesis  
*Iyer JK, Kalra M, Kaul A, Payton ME, Kaul R*
- 6817 Glycosylation-related gene expression in HT29-MTX-E12 cells upon infection by *Helicobacter pylori*  
*Cairns MT, Gupta A, Naughton JA, Kane M, Clyne M, Joshi L*
- 6833 STAT3 deficiency prevents hepatocarcinogenesis and promotes biliary proliferation in thioacetamide-induced liver injury  
*Abe M, Yoshida T, Akiba J, Ikezono Y, Wada F, Masuda A, Sakaue T, Tanaka T, Iwamoto H, Nakamura T, Sata M, Koga H, Yoshimura A, Torimura T*
- 6845 Performance verification and comparison of TianLong automatic hypersensitive hepatitis B virus DNA quantification system with Roche CAP/CTM system  
*Li M, Chen L, Liu LM, Li YL, Li BA, Li B, Mao YL, Xia LF, Wang T, Liu YN, Li Z, Guo TS*

**Case Control Study**

- 6854 Association of insertion-deletions polymorphisms with colorectal cancer risk and clinical features  
*Marques D, Ferreira-Costa LR, Ferreira-Costa LL, Correa RS, Borges AMP, Ito FR, Ramos CCO, Bortolin RH, Luchessi AD, Ribeiro-dos-Santos A, Santos S, Silbiger VN*

**Retrospective Cohort Study**

- 6868 Hospital readmissions in decompensated cirrhotics: Factors pointing toward a prevention strategy

*Seraj SM, Campbell EJ, Argyropoulos SK, Wegermann K, Chung RT, Richter JM*

- 6877 Measurement of biological age may help to assess the risk of colorectal adenoma in screening colonoscopy

*Kim SJ, Kim BJ, Kang H*

**Retrospective Study**

- 6884 Prognostic factors of response to endoscopic treatment in painful chronic pancreatitis

*Tantau A, Mandrutiu A, Leucuta DC, Ciobanu L, Tantau M*

- 6894 *In vivo* histological diagnosis for gastric cancer using endocytoscopy

*Tsurudome I, Miyahara R, Funasaka K, Furukawa K, Matsushita M, Yamamura T, Ishikawa T, Ohno E, Nakamura M, Kawashima H, Watanabe O, Nakaguro M, Satou A, Hirooka Y, Goto H*

**CASE REPORT**

- 6902 Achalasia after bariatric Roux-en-Y gastric bypass surgery reversal

*Abu Ghanimeh M, Qasrawi A, Abughanimeh O, Albadarin S, Clarkston W*

- 6907 Persistent severe hypomagnesemia caused by proton pump inhibitor resolved after laparoscopic fundoplication

*Semb S, Helgstrand F, Hjørne F, Bytzer P*

- 6911 Rupture of small cystic pancreatic neuroendocrine tumor with many microtumors

*Sagami R, Nishikiori H, Ikuyama S, Murakami K*

**LETTERS TO THE EDITOR**

- 6920 Resistance of *Helicobacter pylori* to furazolidone and levofloxacin: A viewpoint

*Zamani M, Rahbar A, Shokri-Shirvani J*

**ABOUT COVER**

Editorial board member of *World Journal of Gastroenterology*, Toru Mizuguchi, MD, PhD, Associate Professor, Surgeon, Department of Surgery, Surgical Oncology and Science, Sapporo Medical University Hospital, Sapporo 060-8543, Hokkaido, Japan

**AIMS AND SCOPE**

*World Journal of Gastroenterology* (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7<sup>th</sup>, 14<sup>th</sup>, 21<sup>st</sup>, and 28<sup>th</sup> each month. The *WJG* Editorial Board consists of 1375 experts in gastroenterology and hepatology from 68 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

**INDEXING/ABSTRACTING**

*World Journal of Gastroenterology* (*WJG*) is now indexed in Current Contents<sup>®</sup>/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch<sup>®</sup>), Journal Citation Reports<sup>®</sup>, Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2017 edition of Journal Citation Reports<sup>®</sup> cites the 2016 impact factor for *WJG* as 3.365 (5-year impact factor: 3.176), ranking *WJG* as 29<sup>th</sup> among 79 journals in gastroenterology and hepatology (quartile in category Q2).

**FLYLEAF**

**I-IX Editorial Board**

**EDITORS FOR THIS ISSUE**

**Responsible Assistant Editor:** Xiang Li  
**Responsible Electronic Editor:** Yan Huang  
**Proofing Editor-in-Chief:** Lian-Sheng Ma

**Responsible Science Editor:** Li-Juan Wei  
**Proofing Editorial Office Director:** Jin-Lei Wang

**NAME OF JOURNAL**  
*World Journal of Gastroenterology*

**ISSN**  
ISSN 1007-9327 (print)  
ISSN 2219-2840 (online)

**LAUNCH DATE**  
October 1, 1995

**FREQUENCY**  
Weekly

**EDITORS-IN-CHIEF**  
**Damian Garcia-Olmo, MD, PhD, Doctor, Professor, Surgeon**, Department of Surgery, Universidad Autonoma de Madrid; Department of General Surgery, Fundacion Jimenez Diaz University Hospital, Madrid 28040, Spain

**Stephen C Strom, PhD, Professor**, Department of Laboratory Medicine, Division of Pathology, Karolinska Institutet, Stockholm 141-86, Sweden

**Andrzej S Tarnawski, MD, PhD, DSc (Med), Professor of Medicine, Chief Gastroenterology**, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach,

CA 90822, United States

**EDITORIAL BOARD MEMBERS**  
All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

**EDITORIAL OFFICE**  
Jin-Lei Wang, Director  
Yuan Qi, Vice Director  
Ze-Mao Gong, Vice Director  
*World Journal of Gastroenterology*  
Baishideng Publishing Group Inc  
7901 Stoneridge Drive, Suite 501,  
Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [editorialoffice@wjgnet.com](mailto:editorialoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>

**PUBLISHER**  
Baishideng Publishing Group Inc  
7901 Stoneridge Drive, Suite 501,  
Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [bpoffice@wjgnet.com](mailto:bpoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>

<http://www.wjgnet.com>

**PUBLICATION DATE**  
October 7, 2017

**COPYRIGHT**  
© 2017 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

**SPECIAL STATEMENT**  
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

**INSTRUCTIONS TO AUTHORS**  
Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

**ONLINE SUBMISSION**  
<http://www.f6publishing.com>

## Retrospective Cohort Study

# Measurement of biological age may help to assess the risk of colorectal adenoma in screening colonoscopy

Sang-Jung Kim, Beom Jin Kim, Hyun Kang

Sang-Jung Kim, Department of Internal Medicine, Bundang Jesaeng Hospital, Sungnam-si, Gyeonggi-do 13590, South Korea

Beom Jin Kim, Department of Internal Medicine, Chung-Ang University College of Medicine, Seoul 06973, South Korea

Hyun Kang, Department of Anesthesiology and Pain Medicine, Chung-Ang University College of Medicine, Seoul 06973, South Korea

ORCID number: Sang-Jung Kim (0000-0002-2530-5906); Beom Jin Kim (0000-0002-0938-6697); Hyun Kang (0000-0003-2844-5880).

**Author contributions:** Kim BJ designed research; Kim BJ and Kim SJ performed research; Kang H analyzed data; Kim SJ wrote the paper.

**Institutional review board statement:** This study protocol was approved by the Ethics Committees at Bundang Jesaeng Hospital (approval No. IMG 15-04).

**Informed consent statement:** Informed consent was waived by the IRB because the study used anonymous clinical data.

**Conflict-of-interest statement:** The authors declare no conflict of interest in this study.

**Data sharing statement:** No additional data are available.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Unsolicited manuscript

**Correspondence to:** Beom Jin Kim, MD, PhD, Associate Professor, Department of Internal Medicine, Chung-Ang University College of Medicine, 102 Heukseok-ro, Dongjak-gu,

Seoul 06973, South Korea. [kimbj@cau.ac.kr](mailto:kimbj@cau.ac.kr)  
Telephone: +82-2-62992060  
Fax: +82-2-62992064

Received: June 11, 2017

Peer-review started: June 12, 2017

First decision: July 13, 2017

Revised: August 15, 2017

Accepted: September 6, 2017

Article in press: September 5, 2017

Published online: October 7, 2017

## Abstract

### AIM

To investigate the clinical utility of biological age (BA) measurement in screening colonoscopy for the detection of colorectal adenomas in the average-risk population.

### METHODS

A consecutive series of asymptomatic subjects aged  $\geq 30$  years who underwent colonoscopy in routine check-ups were enrolled. Colorectal adenoma was classified according to size, number, and location. BAs were calculated using the MEDIAGE™ Biological Age Measurement System.

### RESULTS

A total of 2696 subjects were investigated (1876 men and 820 women). The mean chronological age (CA) was 46.0 years and the mean BA was 44.7 years. Metabolic syndrome (MS) was diagnosed in 218 subjects (8.1%). The prevalence of overall colorectal adenoma was 23.1% (622/2,696). When the subjects were divided into four groups based on BA ( $\leq 39$  years; 40-49 years; 50-59 years;  $\geq 60$  years), the prevalence of colorectal adenoma was increased as BA increased ( $P < 0.001$ ). Colorectal adenoma located in the proximal colon was more prevalent in the BA-dominant group (BA-CA  $\geq 5$



years) than the CA-dominant group ( $CA-BA \geq 5$  years) ( $P = 0.034$ ). When the subjects were categorized into four groups according to MS and age gap between BA and CA, the incidence of colorectal adenoma increased with MS and BA-dominance ( $P < 0.05$ ).

## CONCLUSION

Measurement of BA may help to assess the risk of colorectal adenoma in screening colonoscopy.

**Key words:** Chronological age; Biological age; Colorectal adenoma; Age gap

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** The present study was conducted on the basis of the hypothesis that colorectal adenoma recognized as a premalignant lesion of colorectal cancer is associated with the biological aging process, suggesting that biological age (BA) represents a biological activity in human. Thus, we intended to assess the risk of colorectal adenomas in the average-risk population from the biological perspective. As a result, the present study demonstrated that the prevalence of colorectal adenoma correlated well with BA, and was more common in the distal colon in BA -dominance.

Kim SJ, Kim BJ, Kang H. Measurement of biological age may help to assess the risk of colorectal adenoma in screening colonoscopy. *World J Gastroenterol* 2017; 23(37): 6877-6883 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v23/i37/6877.htm> DOI: <http://dx.doi.org/10.3748/wjg.v23.i37.6877>

## INTRODUCTION

Colorectal cancer (CRC) is one of the most common malignancies and a leading cause of cancer mortality in the world<sup>[1,2]</sup>. Colonoscopy is effective for detecting adenomatous polyps, and colonoscopy with polypectomy has been shown to significantly reduce the expected incidence of CRC by 76%–90% in multiple cohort studies<sup>[3,4]</sup>.

The guidelines of the American Society of Gastrointestinal Endoscopy for colorectal cancer screening and surveillance recommend that screening colonoscopy should begin at the age of 50 for average-risk individuals<sup>[5,6]</sup>. The current cutoff age is based mainly on the fact that the incidence of CRC begins to rise during the 6<sup>th</sup> decade of life<sup>[7]</sup>. However, given the long lead time associated with adenoma-to-carcinoma progression, the increased number of CRCs diagnosed in this age group may reflect the end result of adenomas present in individuals under 50 years old<sup>[6]</sup>. In fact, nearly 1 in 15 of CRCs develops in adults under 50 years of age<sup>[8]</sup>. Therefore, reasonable guidelines for

performing screening colonoscopy for colorectal cancer should be proposed for the average-risk population.

Aging is a highly individual process, and has a high degree of inter-individual and between-individual difference<sup>[9]</sup>. Chronological age (CA) is the age as determined by the passage of time since birth. However, biological age (BA) is determined by physiology rather than chronology<sup>[10]</sup>. As CA does not accurately reflect the functional state of the body, it does not provide an accurate indication of the aging process<sup>[11,12]</sup>. CA and BA may differ significantly, since both genetics and the environment contribute to BA<sup>[10]</sup>. Individuals might be healthier, more vital, and biologically younger than might be expected from their CA, and vice versa<sup>[13]</sup>.

If subjects with an average risk of colon cancer undergo colonoscopy according to BA, colon cancers may be prevented by colonoscopy with polypectomy for colorectal adenomas and premalignant lesions at the optimal age of life. Thus, it is necessary to stratify the risks and identify high-risk groups in a specialized manner. In addition, there might be a rationale by which to recommend screening colonoscopy for early cancer detection and prevention besides CA determination.

To date, no worldwide studies have been conducted to estimate the clinical application of BA as a new indicator in CRC screening. Hence, we intended to assess the risk for colorectal adenomas in the average-risk population from the biological perspective. Therefore, the aim of the present study was to investigate the clinical utility of BA measurement in screening colonoscopy for the detection of colorectal adenomas in the average-risk population.

## MATERIALS AND METHODS

### Subjects

We conducted this study on a consecutive series of asymptomatic subjects  $\geq 30$  years of age who underwent colonoscopy during a routine check-up at the Health Promotion Center of Bundang Jesaeng Hospital in Gyeonggi-do, South Korea between August 2011 and August 2012. The exclusion criteria were a history of colonic disease, such as CRC, polyps, and inflammatory bowel disease, and a history of colorectal surgery or colonoscopy within the previous 10 years. Information regarding age, sex, medical history of diabetes mellitus and hypertension, and family history of colon cancer was collected from a standardized questionnaire. The study was approved by the Institutional Review Board of the Bundang Jesaeng Hospital in Korea (IMG 15-04).

### Measurement of BA

BAs were calculated as described previously using models for predicting BA (MEDIAGE™ Biological Age Measurement System, MEDIAGE Research Center, Gyeonggi-do, South Korea)<sup>[14,15]</sup>. This measurement

system has been used as an adjunctive measure for easily predicting individual differences in health and aging status with the use of clinical parameters that are commonly used at many health check-up centers in South Korea.

#### Test battery parameter

We collected clinical profiles such as physical, biochemical, and hormonal parameters for the calculation of BA. A profile of physical parameters included height, weight, waist circumference, lean body mass, body fat measurement, body mass index, blood pressure, forced expiratory volume 1, and forced vital capacity. Waist circumference was measured at the high point of the iliac crest during minimal respiration. Blood samples were drawn from the antecubital space in the morning. Biochemical parameters included complete blood count with differential count, erythrocyte sedimentation rate, fasting blood glucose, liver function test, lipid profile, blood urea nitrogen, creatinine, bone turnover markers such as osteocalcin and deoxypyridinoline, and tumor markers such as prostate specific antigen, and total antioxidant status were measured using Randox TAS assay kits (Randox, United States) based on the reaction between metmyoglobin and hydrogen peroxide. Lastly, the hormonal parameters included thyroid-stimulating hormone, temporary threshold shift, sex hormone-binding globulin, follicle-stimulating hormone, insulin-like growth factor-1, and dehydroepiandrosterone sulphate.

#### Metabolic syndrome

Based on the ATP III-WPRO (Regional Office for the Western Pacific Region of WHO) definition according to the WPRO criteria, metabolic syndrome (MS) was defined as the presence of three or more of the following<sup>[16]</sup>: (1) abdominal obesity, waist circumference > 90 cm in men and > 80 cm in women; (2) hypertriglyceridemia,  $\geq 150$  mg/dL; (3) low high-density lipoprotein cholesterol level, < 40 mg/dL in men and < 50 mg/dL in women; (4) high blood pressure,  $\geq 130$  mm Hg systolic or  $\geq 85$  mm Hg diastolic; and (5) high serum fasting glucose level,  $\geq 110$  mg/dL.

#### Colonoscopy

We performed colonoscopies that reached the cecum after bowel preparation with 4 L of Colyte powder (TAEJOON PHARM Co., Ltd., South Korea). The Colyte powder was composed of polyethylene glycol (3350.29 g), anhydrous sodium sulfate (2.85 g), sodium hydrogen carbonate (0.84 g), sodium chloride (0.73 g), and potassium chloride (0.37 g). We examined colonoscopic features, including the size, location, number, and histologic findings of polyps. Polyp size was assessed using open colonoscope biopsy forceps (MTW-Endoskopie Manufaktur, Wesel, Germany). Adenoma size was classified into < 10 and  $\geq 10$  mm (the largest size was used for multiple adenomas). The

location of colorectal adenomas was divided into three categories: the proximal colon, including the cecum, ascending colon, and transverse colon; the distal colon, including the splenic flexure, descending colon, sigmoid colon, and rectum; and both sides of the colon. The number of adenomas was classified as either single or multiple ( $\geq 2$ ). Histological findings were classified according to their premalignant potential: colorectal adenomas included tubular, villous, or serrated adenomas; and controls had normal colonoscopic findings and non-polypoid benign lesions such as non-specific colitis or histologically confirmed hyperplastic polyps. In addition, advanced adenomas were defined as  $\geq 1$  cm in estimated diameter, containing  $\geq 25\%$  villous features, and/or high-grade dysplasia.

#### Statistical analysis

For continuous variables, data distribution was first evaluated for normality using the Shapiro-Wilk test. As all the variables did not pass normality test, data were analyzed using Mann-Whitney's *U* tests and presented as medians ( $P_{25}$ – $P_{75}$ ). Descriptive variables were analyzed using  $\chi^2$  analyses or Fisher's exact tests, as appropriate, and  $P < 0.05$  were considered statistically significant. Statistical analyses were conducted using SPSS 21.0 (IBM Corp., Armonk, NY, United States).

## RESULTS

We identified a total of 2699 subjects in the study period. Among them, three with colorectal carcinoma were excluded. Thus, the study population consisted of 2696 subjects. The demographic and clinical characteristics of the enrolled subjects are summarized in Table 1.


The subjects comprised 1876 men (69.6%) and 820 women (30.4%). The median age of the subjects in CA was 46.0 years, whereas the median age of the subjects in BA was 44.7 years old.

MS was diagnosed in 218 subjects (8.1%). When the subjects were divided into four groups based on BA ( $\leq 39$  years; 40–49 years; 50–59 years;  $\geq 60$  years), the prevalence of MS was increased with increasing BA ( $P < 0.001$ ). Interestingly, the gradient of prevalence of MS in BA was higher than that of CA.

The prevalence of overall colorectal adenoma was 23.1% (622/2696). The colonoscopic findings of the subjects according to location, size, and number are summarized in Table 2.

When the subjects were divided into four groups based on BA or CA ( $\leq 39$  years; 40–49 years; 50–59 years;  $\geq 60$  years), the prevalence of colorectal adenoma was increased as BA and CA increased ( $P < 0.001$ ) (Figure 1).

Clinical and colonoscopic characteristics were evaluated in accordance with the age gap between CA and BA. The prevalence of colorectal adenoma in



**Figure 1** Prevalence of colorectal adenoma according to chronological age and biological age. When the subjects were divided into four groups (<39 yr; 40-49 yr; 50-59 yr; ≥ 60 yr), the prevalence of colorectal adenoma increased with increasing biological age ( $P < 0.001$ ).

**Table 2** Characteristics of colorectal adenoma in subjects  $n$  (%)

All subjects ( $n = 2696$ )	
Colorectal adenoma	
Yes	622 (23.1)
No	2074 (76.9)
Location	
Proximal	489 (18.1)
Distal	244 (9.1)
Both	53 (2.0)
Size (mm)	
< 10	714 (26.5)
≥ 10	73 (2.7)
Number	
1	484 (18.0)
≥ 2	138 (5.1)

dominant age groups was assessed. The subjects were divided into two groups based on an age gap between CA and BA of more than 5 years: BA-dominant group ( $BA-CA \geq 5$  years) and CA-dominant group ( $CA-BA \geq 5$  years). The incidences of MS as well as high blood pressure, fasting glucose level, and HbA1c in the BA-dominant group were significantly higher than those in the CA-dominant group (Table 3). As shown in Table 4, colorectal adenoma located in the distal colon was more prevalent in the BA-dominant group than the CA-dominant group ( $P = 0.034$ ).

The prevalence of colorectal adenoma according to MS and age gap was evaluated. We categorized subjects into four groups according to MS and age gap between BA and CA. As a result, the incidence of colorectal adenoma increased with MS and BA-dominance: The incidence of colorectal adenoma was 21.5%, 22.8%, 34.5%, and 39.2% for the first group ( $MS^-$  and  $BA < CA$ ,  $n = 368/1711$ ), the second group ( $MS^-$  and  $BA \geq CA$ ,  $n = 175/767$ ), third group ( $MS^+$  and  $BA \geq CA$ ,  $n = 31/79$ ), and the fourth group ( $MS^+$  and  $BA < CA$ ,  $n = 48/139$ ) ( $P < 0.05$ ) (Figure 2).

**Table 1** Baseline characteristics of the subjects

	All subjects ( $n = 2696$ )	
	Chronological age	Biological age
Age [median ( $P_{25}$ - $P_{75}$ ), yr]	46.0 (39.0-54.0)	44.7 (38.2-53.1)
Gender		
Male	1876	(69.6)
Female	820	(30.4)
High blood pressure		
Yes (%)	544	(20.2)
No (%)	2152	(79.8)
Total cholesterol (mg/dL)	193.00	(172.00-217.75)
LDL cholesterol (mg/dL)	118.00	(99.00-141.00)
HDL cholesterol (mg/dL)	53.00	(45.0-64.00)
Triglyceride (mg/dL)	98.00	(66.00-152.00)
Fasting glucose (mg/dL)	85.00	(79.00-92.00)
HbA1c	5.40	(5.20-5.70)
CEA	1.55	(1.00-2.38)
Metabolic syndrome		
Yes	218	(8.1)
No	2478	(91.9)

Data expressed as median ( $P_{25}$ - $P_{75}$ ), or absolute number (%). High blood pressure,  $\geq 130/85$  mmHg or documented use of antihypertensive therapy. LDL: Low-density lipoprotein; HDL: High-density lipoprotein; HbA1c: Hemoglobin A1c; CEA: Carcinoembryonic antigen.

## DISCUSSION

The present study was conducted on the basis of the hypothesis that colorectal adenoma recognized as a premalignant lesion of colorectal cancer is associated with the biological aging process, suggesting that BA represents a biological activity in human. Thus, we intended to assess the risk of colorectal adenomas in the average-risk population from the biological perspective. As a result, the present study demonstrated that the prevalence of colorectal adenoma correlated well with BA, and was more common in the distal colon in BA-dominance. Furthermore, the incidence of MS correlated more precisely with BA compared with CA.

Aging in humans refers to a multidimensional process of physical, psychological, and social changes that occur from birth<sup>[14]</sup>. Aging is usually assessed in terms of the CA, which is defined as the time elapsed since birth. However, CA fails as an accurate indicator of the aging process<sup>[17]</sup>. BA estimates the functional status of an individual with respect to their chronological peers on the basis of how well they function in comparison with others of the same CA<sup>[18]</sup>. Different individual rates of aging lead to differences between CA and BA, and individual values of BA can therefore vary widely at any given CA and are ultimately expected to correspond to the inter-individual variations in longevity and timing and/or magnitude of sequelae of the aging process<sup>[19,20]</sup>. It has been proposed that BA may serve as an indicator of an individual's general health status, remaining healthy life span, and active life expectancy<sup>[21]</sup>. Considering that BA corresponds well with the aging process, BA may be a good indicator for starting screening colonoscopy at the most biologically



**Table 3 Clinical characteristics between biological age-dominant and chronological age-dominant groups**

	BA-dominant group ( <i>n</i> = 28)	CA-dominant group ( <i>n</i> = 76)	<i>P</i> value
Age (yr, mean ± SD)	42.00 (35.25-49.75)	52.00 (43.25-59.00)	0.001
Gender			0.013
Male	25 (89.3)	49 (64.5)	
Female	3 (10.3)	27 (35.5)	
High blood pressure			< 0.001
Yes (%)	11 (39.3)	6 (7.9)	
No (%)	17 (60.7)	70 (92.1)	
Fasting glucose (mg/dL)	88.00 (82.00-97.50)	81.00 (75.25-87.00)	< 0.001
HbA1c	5.55 (5.30-5.80)	5.30 (5.20-5.50)	0.001
CEA	1.91 (0.97-3.20)	1.67 (1.03-2.57)	0.858
Metabolic syndrome			< 0.001
Yes	7 (25.0)	1 (1.3)	
No	21 (75.0)	75 (98.7)	

High blood pressure systolic blood pressure/diastolic blood pressure > 140/90 mmHg. Data expressed as median (*P*<sub>25</sub>-*P*<sub>75</sub>), or absolute number (%). BA: Biological age; CA: Chronological age; HbA1c: Hemoglobin A1c; CEA: Carcinoembryonic antigen.

**Table 4 Colorectal adenoma between biological age-dominant and chronological age-dominant groups *n* (%)**


	BA-dominant group ( <i>n</i> = 28)	CA-dominant group ( <i>n</i> = 76)	<i>P</i> value
Colorectal adenoma			0.386
Yes	3 (10.7)	15 (19.7)	
No	25 (89.3)	61 (80.3)	
Location			0.034
Proximal	0 (0.0)	18 (23.7)	
Distal	4 (14.3)	6 (7.9)	
Both	1 (3.6)	1 (1.3)	
Size (mm)			0.088
< 10	3 (10.7)	23 (30.3)	
≥ 10	2 (7.1)	2 (2.6)	
Number			0.433
1	2 (7.1)	13 (17.1)	
≥ 2	1 (3.6)	2 (2.6)	

BA: Biological age; CA: Chronological age.

optimal time in an individual's life. If so, CRC may eventually be preventable using colonoscopy with polypectomy for premalignant lesions.

Many previous studies have revealed that colorectal adenoma is associated with MS. The clinical characteristics of MS, such as obesity, dyslipidemia, and insulin resistance, have been linked to an increased risk of CRC in several studies<sup>[16,22]</sup>. In this study, the subjects with MS had more colorectal adenomas compared to those without MS (44.9% vs 27.8%, *P* < 0.001). In particular, the prevalence of colorectal adenoma was increased along with MS and BA-dominance (*P* < 0.001) (Figure 2).

In this study, the prevalence of MS increased proportionally with BA compared with CA. This suggests that MS is related to BA more than to CA because the metabolic change is associated with aging. Soprano *et al.*<sup>[23]</sup> reported that MS can be considered an age-related disease by identifying a correlation between MS and aging. In addition, they highlighted aging-related dysfunctions that contribute to the



**Figure 2 The prevalence of colorectal adenoma according to metabolic syndrome and age gap.** The incidence of colorectal adenoma increased for the first group (MS- and BA < CA), the second group (MS- and BA ≥ CA), the third group (MS+ and BA ≥ CA), and the fourth group (MS+ and BA < CA) (*P* < 0.05). MS: Metabolic syndrome; BA: Biological age; CA: Chronological age.

onset of several metabolic diseases<sup>[23]</sup>. Therefore, physiological disorders such as MS should be measured based on BA rather than CA.

In this study, a CA-dominant subject was defined as an individual younger than their CA, showing that they are healthier than expected. On the other hand, a BA-dominant subject was defined as an individual older than their CA, showing that they are less healthy than expected. The prevalence of MS was higher in the BA-dominant group than in the CA-dominant group in this study. Therefore, this result also suggests the clinical application of BA to various metabolic diseases related with age.

In this study, the prevalence of colorectal adenoma increased with BA. As shown in Figure 1, the prevalence of colorectal adenoma rose sharply along with BA distribution rather than CA distribution with increasing age. The prevalence of colorectal adenoma was higher in the BA-dominant group than in the CA-

dominant group, although the difference between the two groups was not statistically significant. Interestingly, colorectal adenomas in the distal colon were more common in the BA-dominant group compared with the CA-dominant group ( $P = 0.034$ ).

Based on these results, BA deserves attention as an alternative indicator for screening for various cancers including CRC. Interestingly, the prevalence of colorectal adenoma in subjects aged < 50 years was as high as 19.9% (329/2696) in this study. Hence, the commencement of screening colonoscopy should be determined based on BA and physiological criteria rather than indiscriminately at 50 years of CA. This strategy will help in identifying those at risks of CRC at an early stage (before the age of 50 years).

This study was limited because the study population, who voluntarily attended health screening, may have been individuals more concerned about their health than average and trying harder to maintain their health than members of the general population. As a result, the age gap between BA and CA was not as large as expected. Therefore, further studies targeting members of the general population are needed in the future.

The present study highlights the important implications for the future planning of CRC screening in that the higher prevalence of BA-dominance was observed in subjects with colorectal adenoma. It should be emphasized that long-term prevention of both colorectal adenoma along with the biological aging process is of clinical importance.

In conclusion, measurement of BA may help to assess the risk of colorectal adenoma in screening colonoscopy. In particular, careful colonoscopic examination in the distal colon should be performed in BA-dominant individuals.

## COMMENTS

### Background

Current guidelines for colorectal cancer screening are based on chronological age (CA) of 50 years. However, a significant number of colorectal cancers have been found even under 50 years of age. Therefore, screening colonoscopy at a more reasonable age may be necessary from the biological perspective.

### Research frontiers

The authors investigated the clinical utility of biological age (BA) measurement in screening colonoscopy for the detection of colorectal adenomas in the average-risk population.

### Innovations and breakthroughs

The present study demonstrated that the prevalence of colorectal adenoma correlated well with BA, and was more common in the distal colon in BA-dominance. Furthermore, the incidence of metabolic syndrome correlated more precisely with BA compared with CA.

### Applications

Measurement of BA may help to assess the risk of colorectal adenoma in screening colonoscopy. In particular, careful colonoscopic examination in the distal colon should be performed in BA-dominant individuals.

## Terminology

BA is determined by physiology rather than chronology. BA estimates the functional status of an individual with respect to their chronological peers on the basis of how well they function in comparison with others of the same CA.

## Peer-review

The author aimed to investigate the clinical utility of BA measurement in screening colonoscopy for the detection of colorectal adenomas in the average-risk population. The present paper is indeed interesting.

## REFERENCES

- 1 **Shonka NA**, Anderson JR, Panwalkar AW, Reed EC, Steen PD, Ganti AK. Effect of diabetes mellitus on the epidemiology and outcomes of colon cancer. *Med Oncol* 2006; **23**: 515-519 [PMID: 17303910 DOI: 10.1385/MO:23:4:515]
- 2 **Chang CK**, Ulrich CM. Hyperinsulinaemia and hyperglycaemia: possible risk factors of colorectal cancer among diabetic patients. *Diabetologia* 2003; **46**: 595-607 [PMID: 12764580 DOI: 10.1007/s00125-003-1109-5]
- 3 **Boroff ES**, Gurudu SR, Hentz JG, Leighton JA, Ramirez FC. Polyp and adenoma detection rates in the proximal and distal colon. *Am J Gastroenterol* 2013; **108**: 993-999 [PMID: 23567353 DOI: 10.1038/ajg.2013.68]
- 4 **Hirota WK**, Zuckerman MJ, Adler DG, Davila RE, Egan J, Leighton JA, Qureshi WA, Rajan E, Fanelli R, Wheeler-Harbaugh J, Baron TH, Faigel DO; Standards of Practice Committee, American Society for Gastrointestinal Endoscopy. ASGE guideline: the role of endoscopy in the surveillance of premalignant conditions of the upper GI tract. *Gastrointest Endosc* 2006; **63**: 570-580 [PMID: 16564854 DOI: 10.1016/j.gie.2006.02.004]
- 5 **Ma J**, Giovannucci E, Pollak M, Leavitt A, Tao Y, Gaziano JM, Stampfer MJ. A prospective study of plasma C-peptide and colorectal cancer risk in men. *J Natl Cancer Inst* 2004; **96**: 546-553 [PMID: 15069117]
- 6 **Blake GJ**, Pradhan AD, Manson JE, Williams GR, Buring J, Ridker PM, Glynn RJ. Hemoglobin A1c level and future cardiovascular events among women. *Arch Intern Med* 2004; **164**: 757-761 [PMID: 15078645 DOI: 10.1001/archinte.164.7.757]
- 7 **Kim BJ**, Kim YH, Sinn DH, Kang KJ, Kim JY, Chang DK, Son HJ, Rhee PL, Kim JJ, Rhee JC. Clinical usefulness of glycosylated hemoglobin as a predictor of adenomatous polyps in the colorectum of middle-aged males. *Cancer Causes Control* 2010; **21**: 939-944 [PMID: 20373014 DOI: 10.1007/s10552-010-9543-4]
- 8 **Imperiale TF**, Kahi CJ, Stuart JS, Qi R, Born LJ, Glowinski EA, Rex DK. Risk factors for advanced sporadic colorectal neoplasia in persons younger than age 50. *Cancer Detect Prev* 2008; **32**: 33-38 [PMID: 18400417 DOI: 10.1016/j.cdp.2008.01.003]
- 9 **Aihie Sayer A**, Osmond C, Briggs R, Cooper C. Do all systems age together? *Gerontology* 1999; **45**: 83-86 [PMID: 9933730]
- 10 **Wiweko B**, Prawesti DM, Hestiantoro A, Sumapraja K, Natadisastra M, Baziad A. Chronological age vs biological age: an age-related normogram for antral follicle count, FSH and anti-Mullerian hormone. *J Assist Reprod Genet* 2013; **30**: 1563-1567 [PMID: 23955628 DOI: 10.1007/s10815-013-0083-1]
- 11 **Zhang WG**, Bai XJ, Sun XF, Cai GY, Bai XY, Zhu SY, Zhang M, Chen XM. Construction of an integral formula of biological age for a healthy Chinese population using principle component analysis. *J Nutr Health Aging* 2014; **18**: 137-142 [PMID: 24522464 DOI: 10.1007/s12603-013-0345-8]
- 12 **Zhang WG**, Zhu SY, Bai XJ, Zhao DL, Jian SM, Li J, Li ZX, Fu B, Cai GY, Sun XF, Chen XM. Select aging biomarkers based on telomere length and chronological age to build a biological age equation. *Age (Dordr)* 2014; **36**: 9639 [PMID: 24659482 DOI: 10.1007/s11357-014-9639-y]
- 13 **Freude G**, Jakob O, Martus P, Rose U, Seibt R. Predictors of the discrepancy between calendar and biological age. *Occup Med (Lond)* 2010; **60**: 21-28 [PMID: 19666964 DOI: 10.1093/occmed/

- kqp113]
- 14 **Bae CY**, Kang YG, Piao MH, Cho B, Cho KH, Park YK, Yu BY, Lee SW, Kim MJ, Lee SH, Kim YJ, Kim DH, Kim JS, Oh JE. Models for estimating the biological age of five organs using clinical biomarkers that are commonly measured in clinical practice settings. *Maturitas* 2013; **75**: 253-260 [PMID: 23642770 DOI: 10.1016/j.maturitas.2013.04.008]
  - 15 **Bae CY**, Kang YG, Suh YS, Han JH, Kim SS, Shim KW. A model for estimating body shape biological age based on clinical parameters associated with body composition. *Clin Interv Aging* 2013; **8**: 11-18 [PMID: 23293513 DOI: 10.2147/CIA.S38220]
  - 16 **Kim JH**, Lim YJ, Kim YH, Sung IK, Shim SG, Oh SO, Park SS, Yang S, Son HJ, Rhee PL, Kim JJ, Rhee JC, Choi YH. Is metabolic syndrome a risk factor for colorectal adenoma? *Cancer Epidemiol Biomarkers Prev* 2007; **16**: 1543-1546 [PMID: 17684126 DOI: 10.1158/1055-9965.EPI-07-0199]
  - 17 **Finkel D**, Whitfield K, McGue M. Genetic and environmental influences on functional age: a twin study. *J Gerontol B Psychol Sci Soc Sci* 1995; **50**: P104-P113 [PMID: 7757832]
  - 18 **Borkan GA**, Norris AH. Assessment of biological age using a profile of physical parameters. *J Gerontol* 1980; **35**: 177-184 [PMID: 6967883]
  - 19 **Anstey KJ**, Lord SR, Smith GA. Measuring human functional age: a review of empirical findings. *Exp Aging Res* 1996; **22**: 245-266 [PMID: 8872080 DOI: 10.1080/03610739608254010]
  - 20 **Mitnitski AB**, Graham JE, Mogilner AJ, Rockwood K. Frailty, fitness and late-life mortality in relation to chronological and biological age. *BMC Geriatr* 2002; **2**: 1 [PMID: 11897015]
  - 21 **Borkan GA**, Norris AH. Biological age in adulthood: comparison of active and inactive U.S. males. *Hum Biol* 1980; **52**: 787-802 [PMID: 6970713]
  - 22 **Kim BC**, Shin A, Hong CW, Sohn DK, Han KS, Ryu KH, Park BJ, Nam JH, Park JW, Chang HJ, Choi HS, Kim J, Oh JH. Association of colorectal adenoma with components of metabolic syndrome. *Cancer Causes Control* 2012; **23**: 727-735 [PMID: 22450737 DOI: 10.1007/s10552-012-9942-9]
  - 23 **Soprano M**, Rusciano MR, Ciccarelli M, Maione AS, Formisano P, Illario M. Metabolic Syndrome and Aging: Calcium Signaling as Common Regulator. *Curr Diabetes Rev* 2015; **12**: 84-89 [PMID: 26201430]

**P- Reviewer:** Riss S, Shi HY, Lakatos PL **S- Editor:** Ma YJ

**L- Editor:** A **E- Editor:** Huang Y





Published by **Baishideng Publishing Group Inc**  
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA  
Telephone: +1-925-223-8242  
Fax: +1-925-223-8243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>



ISSN 1007-9327



9 771007 932045