

# World Journal of *Gastroenterology*

*World J Gastroenterol* 2017 October 7; 23(37): 6747-6922



**EDITORIAL**

- 6747 Microbial dysbiosis in spouses of ulcerative colitis patients: Any clues to disease pathogenesis?  
*Sorrentino D*

**REVIEW**

- 6750 Road to stemness in hepatocellular carcinoma  
*Flores-Téllez TNJ, Villa-Treviño S, Piña-Vázquez C*
- 6777 Intrahepatic vascular changes in non-alcoholic fatty liver disease: Potential role of insulin-resistance and endothelial dysfunction  
*Pasarin M, Abraldes JG, Liguori E, Kok B, La Mura V*
- 6788 Epidemiological and clinical perspectives on irritable bowel syndrome in India, Bangladesh and Malaysia: A review  
*Rahman MM, Mahadeva S, Ghoshal UC*

**ORIGINAL ARTICLE****Basic Study**

- 6802 Estrogen receptor expression in chronic hepatitis C and hepatocellular carcinoma pathogenesis  
*Iyer JK, Kalra M, Kaul A, Payton ME, Kaul R*
- 6817 Glycosylation-related gene expression in HT29-MTX-E12 cells upon infection by *Helicobacter pylori*  
*Cairns MT, Gupta A, Naughton JA, Kane M, Clyne M, Joshi L*
- 6833 STAT3 deficiency prevents hepatocarcinogenesis and promotes biliary proliferation in thioacetamide-induced liver injury  
*Abe M, Yoshida T, Akiba J, Ikezono Y, Wada F, Masuda A, Sakaue T, Tanaka T, Iwamoto H, Nakamura T, Sata M, Koga H, Yoshimura A, Torimura T*
- 6845 Performance verification and comparison of TianLong automatic hypersensitive hepatitis B virus DNA quantification system with Roche CAP/CTM system  
*Li M, Chen L, Liu LM, Li YL, Li BA, Li B, Mao YL, Xia LF, Wang T, Liu YN, Li Z, Guo TS*

**Case Control Study**

- 6854 Association of insertion-deletions polymorphisms with colorectal cancer risk and clinical features  
*Marques D, Ferreira-Costa LR, Ferreira-Costa LL, Correa RS, Borges AMP, Ito FR, Ramos CCO, Bortolin RH, Luchessi AD, Ribeiro-dos-Santos A, Santos S, Silbiger VN*

**Retrospective Cohort Study**

- 6868** Hospital readmissions in decompensated cirrhotics: Factors pointing toward a prevention strategy

*Seraj SM, Campbell EJ, Argyropoulos SK, Wegermann K, Chung RT, Richter JM*

- 6877** Measurement of biological age may help to assess the risk of colorectal adenoma in screening colonoscopy

*Kim SJ, Kim BJ, Kang H*

**Retrospective Study**

- 6884** Prognostic factors of response to endoscopic treatment in painful chronic pancreatitis

*Tantau A, Mandrutiu A, Leucuta DC, Ciobanu L, Tantau M*

- 6894** *In vivo* histological diagnosis for gastric cancer using endocytoscopy

*Tsurudome I, Miyahara R, Funasaka K, Furukawa K, Matsushita M, Yamamura T, Ishikawa T, Ohno E, Nakamura M, Kawashima H, Watanabe O, Nakaguro M, Satou A, Hirooka Y, Goto H*

**CASE REPORT**

- 6902** Achalasia after bariatric Roux-en-Y gastric bypass surgery reversal

*Abu Ghanimeh M, Qasrawi A, Abughanimeh O, Albadarin S, Clarkston W*

- 6907** Persistent severe hypomagnesemia caused by proton pump inhibitor resolved after laparoscopic fundoplication

*Semb S, Helgstrand F, Hjørne F, Bytzer P*

- 6911** Rupture of small cystic pancreatic neuroendocrine tumor with many microtumors

*Sagami R, Nishikiori H, Ikuyama S, Murakami K*

**LETTERS TO THE EDITOR**

- 6920** Resistance of *Helicobacter pylori* to furazolidone and levofloxacin: A viewpoint

*Zamani M, Rahbar A, Shokri-Shirvani J*

**ABOUT COVER**

Editorial board member of *World Journal of Gastroenterology*, Toru Mizuguchi, MD, PhD, Associate Professor, Surgeon, Department of Surgery, Surgical Oncology and Science, Sapporo Medical University Hospital, Sapporo 060-8543, Hokkaido, Japan

**AIMS AND SCOPE**

*World Journal of Gastroenterology* (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7<sup>th</sup>, 14<sup>th</sup>, 21<sup>st</sup>, and 28<sup>th</sup> each month. The *WJG* Editorial Board consists of 1375 experts in gastroenterology and hepatology from 68 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

**INDEXING/ABSTRACTING**

*World Journal of Gastroenterology* (*WJG*) is now indexed in Current Contents<sup>®</sup>/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch<sup>®</sup>), Journal Citation Reports<sup>®</sup>, Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2017 edition of Journal Citation Reports<sup>®</sup> cites the 2016 impact factor for *WJG* as 3.365 (5-year impact factor: 3.176), ranking *WJG* as 29<sup>th</sup> among 79 journals in gastroenterology and hepatology (quartile in category Q2).

**FLYLEAF**

**I-IX Editorial Board**

**EDITORS FOR THIS ISSUE**

**Responsible Assistant Editor:** Xiang Li  
**Responsible Electronic Editor:** Yan Huang  
**Proofing Editor-in-Chief:** Lian-Sheng Ma

**Responsible Science Editor:** Li-Juan Wei  
**Proofing Editorial Office Director:** Jin-Lei Wang

**NAME OF JOURNAL**  
*World Journal of Gastroenterology*

**ISSN**  
ISSN 1007-9327 (print)  
ISSN 2219-2840 (online)

**LAUNCH DATE**  
October 1, 1995

**FREQUENCY**  
Weekly

**EDITORS-IN-CHIEF**  
**Damian Garcia-Olmo, MD, PhD, Doctor, Professor, Surgeon**, Department of Surgery, Universidad Autonoma de Madrid; Department of General Surgery, Fundacion Jimenez Diaz University Hospital, Madrid 28040, Spain

**Stephen C Strom, PhD, Professor**, Department of Laboratory Medicine, Division of Pathology, Karolinska Institutet, Stockholm 141-86, Sweden

**Andrzej S Tarnawski, MD, PhD, DSc (Med), Professor of Medicine, Chief Gastroenterology**, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach,

CA 90822, United States

**EDITORIAL BOARD MEMBERS**  
All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

**EDITORIAL OFFICE**  
Jin-Lei Wang, Director  
Yuan Qi, Vice Director  
Ze-Mao Gong, Vice Director  
*World Journal of Gastroenterology*  
Baishideng Publishing Group Inc  
7901 Stoneridge Drive, Suite 501,  
Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [editorialoffice@wjgnet.com](mailto:editorialoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>

**PUBLISHER**  
Baishideng Publishing Group Inc  
7901 Stoneridge Drive, Suite 501,  
Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [bpoffice@wjgnet.com](mailto:bpoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>

<http://www.wjgnet.com>

**PUBLICATION DATE**  
October 7, 2017

**COPYRIGHT**  
© 2017 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

**SPECIAL STATEMENT**  
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

**INSTRUCTIONS TO AUTHORS**  
Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

**ONLINE SUBMISSION**  
<http://www.f6publishing.com>

## Microbial dysbiosis in spouses of ulcerative colitis patients: Any clues to disease pathogenesis?

Dario Sorrentino

Dario Sorrentino, IBD Center, Division of Gastroenterology, Virginia Tech Carilion School of Medicine, Roanoke, VA 24016, United States

Dario Sorrentino, Department of Clinical and Experimental Medical Sciences, University of Udine School of Medicine, 33100 Udine, Italy

ORCID number: Dario Sorrentino (0000-0002-2694-8823).

Author contributions: Sorrentino D solely wrote the manuscript

Conflict-of-interest statement: The author declares no conflict of interest related to this publication.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Dario Sorrentino, MD, FRACP, Professor of Medicine, IBD Center, Division of Gastroenterology, Virginia Tech Carilion School of Medicine, 3 Riverside Circle, Roanoke, VA 24016, United States. [drsorrentino@carilionclinic.org](mailto:drsorrentino@carilionclinic.org)  
Telephone: +1-540-5261057  
Fax: +1-540-9859418

Received: June 22, 2017

Peer-review started: June 24, 2017

First decision: July 13, 2017

Revised: July 30, 2017

Accepted: August 25, 2017

Article in press: August 25, 2017

Published online: October 7, 2017

### Abstract

A number of alterations have been found within the gut

microbial profile of patients with inflammatory bowel diseases when compared with the healthy population; however, it is unclear whether such dysbiosis is the cause or simply the consequence of the disease state. In ulcerative colitis, the environment seems to play a crucial role in disease etiology since monozygotic twins show a concordance rate of only 8%-10% - though it is unclear whether it does so by acting through the microbiome. In this study, the authors investigated the influence of cohabitation on the gut microbial community in healthy partners of ulcerative colitis patients - with the intent of clarifying some of these issues. As expected, ulcerative colitis patients had a significant dysbiosis and alterations in microbial metabolism. Interestingly, these abnormal fecal microbial communities were relatively similar amongst patients and their spouses. Thus, this study shows that the microbial profile might be partially transferred from ulcerative colitis patients to healthy individuals. Whether this finding impacts on disease development or has any implication for the role of the microbiome in inflammatory bowel disease etiology remains to be determined.

**Key words:** Ulcerative colitis; Cohabitation; Spouses; Gut microbiome

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** Dysbiosis in inflammatory bowel diseases is common. However, microbial dysbiosis could be a consequence, rather than cause of the disease state. In this study the authors detected dysbiosis and altered microbial metabolism not only in ulcerative colitis patients but also in their healthy cohabiting partners. Therefore, the microbiome might be partially transferred from ulcerative colitis patients to healthy individuals. Since spouses of ulcerative colitis patients do not have an increased risk of developing the disease this study suggests that dysbiosis might be an effect rather than the cause of the disease. Overall, the precise role of the microbiome in inflammatory bowel



disease etiology remains unknown.

Sorrentino D. Microbial dysbiosis in spouses of ulcerative colitis patients: Any clues to disease pathogenesis? *World J Gastroenterol* 2017; 23(37): 6747-6749 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v23/i37/6747.htm> DOI: <http://dx.doi.org/10.3748/wjg.v23.i37.6747>

Man has coexisted with microbes since the beginning of time. We bear trillions of microbes in our body. Indeed, this symbiotic microbial cell population (or microbiome) has been reported to outnumber that of the human host by 10:1 - though more recent estimates appear to validate such ratio only when considering human nucleate cells<sup>[1]</sup>. The largest portion of the microbiome is located in the gastrointestinal tract and functions as a digestive organ, a training tool for the immune system, and a guardian of harmful infections<sup>[2]</sup>. A number of factors can affect the microbial community in the gut--both in healthy and diseased states-- and include diet, age, medical therapy, smoking, and genetics<sup>[3]</sup>. Many human diseases have been linked to an altered gut microbiota. Among those, the most studied are the inflammatory bowel diseases (IBD) - both Ulcerative colitis (UC) and Crohn's disease (CD). Several alterations (dysbiosis) have been found in the gut microbial profile of IBD patients including a decrease in diversity, *Bacteroides*, *Firmicutes*, *Clostridia*, *Ruminococcaceae*, *Bifidobacterium*, *Lactobacillus*, and *F. prausnitzii*; an increase in *Gammaproteobacteria*; and the presence of adherent invasive *E. coli* and *Fusobacterium*<sup>[2]</sup>. Altered microbial function has also been reported in IBD patients-specifically, butyrate, butanoate and propanoate metabolism are decreased while the oxidative stress, the type II secretion system, secretion of toxins, the transport of both amino acids and sulfates are increased<sup>[2]</sup>.

Despite these impressive and reproducible alterations in the gut microbiome of IBD patients, it is unclear whether such dysbiosis is the cause or simply the consequence (*i.e.*, bacterial adaptation for survival) of the disease state. This is a major mechanistic issue with a potential crucial impact on development of new medications and on IBD management. Current evidence supports both views. For example, disease activity seems to affect the microbial profile in IBD while microbial alterations similar to those described in IBD have been described in patients with nonspecific intestinal injury<sup>[4]</sup>. These observations suggest that the microbial profile in IBD might be the result of local changes in the intestinal mucosa following the disease onset. On the other hand, dysbiosis also seems to be associated with genetic polymorphisms. In addition, immune mediated colitis can be elicited in genetically susceptible mice with single commensal bacterial species<sup>[4]</sup>. Thus, these findings highlight the potential role of the microbiome in IBD etiology.

As for the individual IBD's, studies in monozygotic twins show a concordance rate of 44%-55% in CD and only 8%-10% in UC<sup>[5]</sup>. Thus, the environment seems to play a crucial role in disease etiology - especially in UC - though it is unclear whether it does so by acting through the microbiome.

In the paper published in the 25<sup>th</sup> issue of the Journal, Chen *et al*<sup>[6]</sup> studied the influence of cohabitation on the gut microbial community in healthy partners of UC patients using 16S rRNA amplicon sequencing. Fecal samples were collected from 8 UC patients and their healthy partners at Lishui People's Hospital in China. Fecal microbial communities showed a higher similarity among UC patients than in their healthy partners (*i.e.*, the partners' group was more heterogeneous). As a whole, UC patients had a lower relative abundance of bacteria belonging to the *Firmicutes* - especially *Blautia*, *Clostridium*, *Coprococcus* and *Roseburia*. In addition, the microbiome of UC patients showed a greater lipid and nucleotide metabolism. Microbiota dysbiosis and altered microbial metabolism were detected in both UC patients and their healthy partners, with the most relevant genera in the latter group being *Akkermansia*, *Bacteroides*, *Escherichia*, *Lactobacillales*, *Klebsiella*, and *Parabacteroides*. Importantly, using 3 different types of analysis the authors showed that the entire microbiota of UC patients was not significantly divergent from those of their partners. Healthy partners of UC patients also showed an increased microbial membrane transport and metabolism of cofactors and vitamins.

The authors concluded that microbial composition and metabolism in healthy partners of UC patients may be impacted by cohabitation.

This is a small study without follow up which would have possibly clarified if the altered microbiome in spouses of UC patients may eventually impact on their health and bear any clinical meaning. No information is given regarding the number of samples per study subject and whether samples were collected the same day(s) by patients and their spouses. Furthermore, there were differences in age and sex distribution between the UC group and the spouses' group. This difference might bear implications for microbial profile heterogeneity<sup>[3]</sup>. Also, based on C-reactive protein levels, the disease activity was not uniformly distributed among UC patients - another potential source of heterogeneity in the microbial profile results. The methodology used by the authors in this study - 16S rRNA gene sequencing - is sound and more accurate than culture dependent methods used in many similar studies<sup>[7]</sup>. The authors' conclusions are reasonable and apparently support an important role of the environment in changing the gut microbiome - in this case, the UC microbiome modifying the bacterial profile of their healthy cohabiting partners. The study does not clarify whether the transfer of the microbiome might occur by direct contact or by other means.

To lend more weight to the authors' conclusion

in this and similar types of studies, IBD and any unrelated upper GI condition should be duly excluded in cohabiting spouses. While disease is unlikely in the absence of symptoms, the hypothetical concomitant presence of IBD in the spouses could clearly explain the similarities between their microbial profile and that of their patient partners. Furthermore, in the Chen *et al*<sup>[6]</sup> study only the stool bacteria were tested; however, the stool microbiome might be different from the intestinal tissue microbiome - which more closely represents the intestinal mucosa flora<sup>[2]</sup>. Finally, the microbial profile of individuals cohabiting with IBD patients should also be compared with that of healthy controls living in the same geographical area and socio-economic conditions to minimize the impact of unrelated environmental factors. The microbial profile of the Asian population might differ from that of the Western population<sup>[8]</sup> - an important factor to keep in mind in comparative studies.

So, does this study suggest that the microbiome dysbiosis in IBD might be a primary event rather than the result of the disease? The answer is no. This study shows that the environment might influence the microbial profile in healthy individuals; however, the impact of such change on disease development remains unclear. Likely, dysbiosis in and by itself is insufficient to cause disease. Indeed, there is no evidence that cohabitation with IBD patients might increase the risk of disease in the spouses, so the results of this study only indicate that the microbiome can be partially transmitted from individual to individual. The final microbial profile in each individual - whether healthy or not - might be the result of the very complex nature of the microbial development process<sup>[9]</sup>, which evolves over time and is affected by a number of factors<sup>[3]</sup>.

## CONCLUSION

The findings of this study support the well-known observation that cohabitation is indeed a way to share our microbial cell population with family members,

or even with our dogs<sup>[10]</sup>. Whether this impacts disease development or points to a specific role of the microbiome on IBD etiology remains to be determined.

## REFERENCES

- 1 **Sender R**, Fuchs S, Milo R. Are We Really Vastly Outnumbered? Revisiting the Ratio of Bacterial to Host Cells in Humans. *Cell* 2016; **164**: 337-340 [PMID: 26824647 DOI: 10.1016/j.cell.2016.01.013]
- 2 **Kostic AD**, Xavier RJ, Gevers D. The microbiome in inflammatory bowel disease: current status and the future ahead. *Gastroenterology* 2014; **146**: 1489-1499 [PMID: 24560869 DOI: 10.1053/j.gastro.2014.02.009]
- 3 **Blaser MJ**, Falkow S. What are the consequences of the disappearing human microbiota? *Nat Rev Microbiol* 2009; **7**: 887-894 [PMID: 19898491 DOI: 10.1038/nrmicro2245]
- 4 **Sartor RB**. Genetics and environmental interactions shape the intestinal microbiome to promote inflammatory bowel disease versus mucosal homeostasis. *Gastroenterology* 2010; **139**: 1816-1819 [PMID: 21029802 DOI: 10.1053/j.gastro.2010.10.036]
- 5 **Halfvarson J**, Bodin L, Tysk C, Lindberg E, Järnerot G. Inflammatory bowel disease in a Swedish twin cohort: a long-term follow-up of concordance and clinical characteristics. *Gastroenterology* 2003; **124**: 1767-1773 [PMID: 12806610]
- 6 **Chen GL**, Zhang Y, Wang WY, Ji XL, Meng F, Xu PS, Yang NM, Ye FQ, Bo XC. Partners of patients with ulcerative colitis exhibit a biologically relevant dysbiosis in fecal microbial metacommunities. *World J Gastroenterol* 2017; **23**: 4624-4631 [PMID: 28740351 DOI: 10.3748/wjg.v23.i25.4624]
- 7 **Janda JM**, Abbott SL. 16S rRNA gene sequencing for bacterial identification in the diagnostic laboratory: pluses, perils, and pitfalls. *J Clin Microbiol* 2007; **45**: 2761-2764 [PMID: 17626177]
- 8 **Prideaux L**, Kang S, Wagner J, Buckley M, Mahar JE, De Cruz P, Wen Z, Chen L, Xia B, van Langenberg DR, Lockett T, Ng SC, Sung JJ, Desmond P, McSweeney C, Morrison M, Kirkwood CD, Kamm MA. Impact of ethnicity, geography, and disease on the microbiota in health and inflammatory bowel disease. *Inflamm Bowel Dis* 2013; **19**: 2906-2918 [PMID: 24240708 DOI: 10.1097/01.MIB.0000435759.05577.12]
- 9 **Dalal SR**, Chang EB. The microbial basis of inflammatory bowel diseases. *J Clin Invest* 2014; **124**: 4190-4196 [PMID: 25083986 DOI: 10.1172/JCI72330]
- 10 **Song SJ**, Lauber C, Costello EK, Lozupone CA, Humphrey G, Berg-Lyons D, Caporaso JG, Knights D, Clemente JC, Nakielnny S, Gordon JI, Fierer N, Knight R. Cohabiting family members share microbiota with one another and with their dogs. *Elife* 2013; **2**: e00458 [PMID: 23599893 DOI: 10.7554/eLife.00458]

**P- Reviewer:** Gazouli M, Hamilton M, Lakatos PL, Wine E **S- Editor:** Gong ZM **L- Editor:** A **E- Editor:** Huang Y





Published by **Baishideng Publishing Group Inc**  
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA  
Telephone: +1-925-223-8242  
Fax: +1-925-223-8243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>



ISSN 1007-9327



9 771007 932045