



Factors associated with carcinoid syndrome in patients with gastrointestinal



学术搜索

找到约 21,200 条结果 (用时0.13秒)

我的著作引用情况

文章

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

我的图书馆

Carcinoid tumors an analysis of 2837 cases

JD Godwin - *Cancer*, 1975 - Wiley Online Library

... B Grossman, Appendiceal neuroendocrine neoplasms: diagnosis and management, *Endocrine-Related Cancer*, 2016, 23 ... Jung Son, Young-Ho Kim, Evaluation of the risk factors associated with rectal ... Al-Efraji, Mohammed A. Aljama, Hagen Fritz Kennecke, Association of dose ...

被引用次数: 1476 相关文章 所有 7 个版本 引用 保存

时间不限

2017以来

2016以来

2013以来

自定义范围...

Everolimus plus octreotide long-acting repeatable for the treatment of advanced neuroendocrine tumours associated with carcinoid syndrome (RADIANT-2): a ...

ME Pavel, JD Hainsworth, E Baudin, M Peeters... - *The Lancet*, 2011 - Elsevier

... 4, 5, 6 and 7 Autocrine activation of the mTOR signalling pathway, mediated through insulin-like growth factor I, has been associated with neuroendocrine tumour ... Current tumour-related symptoms †, 170 (79%), 172 (81%). Organ type involved ‡. Liver, 198 (92%), 196 (92%). ...

被引用次数: 635 相关文章 所有 26 个版本 引用 保存

[PDF] pagespro-orange.fr

按相关性排序

按日期排序

One hundred years after "carcinoid": epidemiology of and prognostic factors for neuroendocrine tumors in 35,825 cases in the United States

JC Yao, M Hassan, A Phan, C Dagohoy... - *Journal of clinical ...*, 2008 - ascopubs.org

... are also known as high-grade NETs, are aggressive and associated with poor ... Whether changes in dietary habits, environmental factors, and use of certain medications such ... receptor radiotherapy and systemic agents targeting vascular endothelial growth factor and mammalian ...

被引用次数: 2484 相关文章 所有 10 个版本 引用 保存

[HTML] ascopubs.org

不限语言

中文网页

简体中文网页

包括专利

包含引用

Prognostic factors and survival in endocrine tumor patients: comparison between gastrointestinal and pancreatic localization

F Panzuto, S Nasoni, M Falconi... - *...-Related Cancer*, 2005 - Soc Endocrinology

... To our knowledge, the role of sstr as a prognostic factor has not previously been ... at the time of diagnosis, irrespective of the presence of an associated syndrome, and of the ... and the presence of distant (particularly extra-hepatic) metastases; therefore, these factors should always ...

被引用次数: 343 相关文章 所有 12 个版本 引用 保存

[HTML] endocrinology-journals.org

创建快讯

Activity of sunitinib in patients with advanced neuroendocrine tumors

MH Kulke, HJ Lenz, NJ Meropol, J Posey... - *Journal of Clinical ...*, 2008 - ascopubs.org

... with advanced carcinoid tumors; moreover, their prolonged use is often associated with toxicity. ... β, stem-cell factor receptor, glial cell line-derived neurotrophic factor receptor (rearranged ... accident (0.9%), and hyponatremia (0.9%; Table 2). A single treatment-related death was ...

被引用次数: 558 相关文章 所有 8 个版本 引用 保存

[HTML] ascopubs.org

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 35217

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Factors associated with carcinoid syndrome in patients with gastrointestinal neuroendocrine tumors

Bei-Lei Cai, Michael S Broder, Eunice Chang, Ting-Jian Yan, David C Metz

Abstract

AIM

To discover unknown factors associated with carcinoid syndrome (CS) with the goal of earlier diagnosis of CS.

METHODS

In this retrospective case-control study using US administrative claims, patients (≥ 18 years) newly diagnosed with gastrointestinal neuroendocrine tumors (GI NETs) without CS (controls) were exactly matched to patients with CS (cases) based on NET

Match Overview

1

Crossref 27 words

Joshi, Kruti, Xiaoyun Pan, Rosa Wang, Erru Yang, and Carmela Benson. "Healthcare Resource Utilization of Second-

1%

SIMILAR



Factors associated with carcinoid syndrome in patients with gastrointestinal



学术搜索

找到约 21,200 条结果 (用时0.05秒)

我的著作引用情况

文章

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

我的图书馆

Carcinoid tumors an analysis of 2837 cases

JD Godwin - *Cancer*, 1975 - Wiley Online Library

... B Grossman, Appendiceal neuroendocrine neoplasms: diagnosis and management, *Endocrine-Related Cancer*, 2016, 23 ... Jung Son, Young-Ho Kim, Evaluation of the risk factors associated with rectal ... Al-Efraji, Mohammed A. Aljama, Hagen Fritz Kennecke, Association of dose ... 被引用次数: 1476 相关文章 所有 7 个版本 引用 保存

时间不限

2017以来

2016以来

2013以来

自定义范围...

Everolimus plus octreotide long-acting repeatable for the treatment of advanced neuroendocrine tumours associated with carcinoid syndrome (RADIANT-2): a ...

ME Pavel, JD Hainsworth, E Baudin, M Peeters... - *The Lancet*, 2011 - Elsevier

... 4, 5, 6 and 7 Autocrine activation of the mTOR signalling pathway, mediated through insulin-like growth factor I, has been associated with neuroendocrine tumour ... Current tumour-related symptoms †, 170 (79%), 172 (81%). Organ type involved ‡. Liver, 198 (92%), 196 (92%) ... 被引用次数: 635 相关文章 所有 26 个版本 引用 保存

[PDF] pagespro-orange.fr

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

One hundred years after "carcinoid": epidemiology of and prognostic factors for neuroendocrine tumors in 35,825 cases in the United States

JC Yao, M Hassan, A Phan, C Dagohoy... - *Journal of clinical ...*, 2008 - ascopubs.org

... are also known as high-grade NETs, are aggressive and associated with poor ... Whether changes in dietary habits, environmental factors, and use of certain medications such ... receptor radiotherapy and systemic agents targeting vascular endothelial growth factor and mammalian ... 被引用次数: 2484 相关文章 所有 10 个版本 引用 保存

[HTML] ascopubs.org

包括专利

包含引用

创建快讯

Prognostic factors and survival in endocrine tumor patients: comparison between gastrointestinal and pancreatic localization

F Panzuto, S Nasoni, M Falconi... - *...-Related Cancer*, 2005 - Soc Endocrinology

... To our knowledge, the role of sstr as a prognostic factor has not previously been ... at the time of diagnosis, irrespective of the presence of an associated syndrome, and of the ... and the presence of distant (particularly extra-hepatic) metastases; therefore, these factors should always ... 被引用次数: 343 相关文章 所有 12 个版本 引用 保存

[HTML] endocrinology-journals.org

Activity of sunitinib in patients with advanced neuroendocrine tumors

MH Kulke, HJ Lenz, NJ Meropol, J Posey... - *Journal of Clinical ...*, 2008 - ascopubs.org

... with advanced carcinoid tumors; moreover, their prolonged use is often associated with toxicity. ... β, stem-cell factor receptor, glial cell line-derived neurotrophic factor receptor (rearranged ... accident (0.9%), and hyponatremia (0.9%; Table 2). A single treatment-related death was ... 被引用次数: 558 相关文章 所有 8 个版本 引用 保存

[HTML] ascopubs.org

[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 108,000 条结果 (用时 0.59 秒)

Google 学术: Factors associated with carcinoid syndrome in patients with gastrointestinal neuroendocrine tumors

Carcinoid tumors an analysis of 2837 cases - Godwin - 被引用次数: 1478

... neuroendocrine tumours associated with carcinoid ... - Pavel - 被引用次数: 638

... of and prognostic factors for neuroendocrine tumors in ... - Yao - 被引用次数: 2484

Predictive factors of carcinoid syndrome among patients with ...

ascopubs.org/doi/abs/10.1200/JCO.2017.35.15_suppl.e15690 - 翻译此页

作者: B Cai - 2017

Predictive factors of carcinoid syndrome among patients with gastrointestinal neuroendocrine tumors (GI NETs) ... Background: Carcinoid syndrome (CS) is associated with a significant symptom burden and deleterious impacts on quality of life ...

Prognosis and Survival in Patients With Gastrointestinal Tract ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: KO Shebani - 1999 - 被引用次数: 299 - 相关文章

Survival estimates for prognostic factors were calculated using Kaplan-Meier product ... Surgically treated patients with carcinoid tumor have an overall favorable 83% ... Carcinoid tumors are neuroendocrine tumors and, as such, are part of the The frequency of the specific carcinoid syndrome-associated symptoms was ...

Gastrointestinal Carcinoid Tumors Treatment (PDQ®) - PDQ Cancer ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > Bookshelf - 翻译此页

作者: PDQATE Board - 2015 - 相关文章

2015年7月8日 - General Information About Gastrointestinal (GI) Carcinoid Tumors ... Neuroendocrine cells have uniform nuclei and abundant granular or faintly staining ... Carcinoid syndrome, which occurs in fewer than 20% of patients with carcinoid ... Factors associated with progression of carcinoid heart