

Value of histomorphometric tumour thickness and smoothelin for conventional m-classification in early oesophageal adenocarcinoma

B Katharina Endhardt^{1*}, Bruno Märkl^{1*}, Andreas Probst², Tina Schaller¹, Daniela Aust³

¹Institute of Pathology, Klinikum Augsburg, Augsburg, Germany

²Department of Gastroenterology, Klinikum Augsburg, Augsburg, Germany

³Institute of Pathology, Universitätsklinikum Dresden, Dresden, Germany

Author contributions: Endhardt K, Märkl B, Probst A, and Aust D. contributed to study conception and design; Schaller T, contributed to data acquisition, data analysis, and interpretation; Endhardt K, Märkl B. and Aust D, contributed to data acquisition, data analysis, and interpretation writing of article; all authors contributed to editing, reviewing and final approval of the article.

Institutional review board statement:

The study has been approved by the internal review board of the Klinikum Augsburg (Review-Number: BKF 2015-13). This study has been performed according to the national rules (Bayerisches Krankenhausgesetz BayKrG – Art 27 (4))

Informed consent is not necessary for this retrospective study.

Conflict-of-interest statement: All authors declare that no conflicting interests (including but not limited to commercial, personal, political, intellectual or religious interests) exist.

Biostatistics statement:

All statistics have been performed and double checked by an experienced researcher (B.M) that published over 50 scientific papers with complex statistical calculations.

Bruno Märkl

15/08/2017

