

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 35413

Manuscript Type: ORIGINAL ARTICLE

Prospective Study

Effect of local wound infiltration with ropivacaine on postoperative pain relief and stress response reduction after open hepatectomy

Sun J *et al.* Wound infiltration after open hepatectomy

Jing-Xian Sun, Ke-Yun Bai, Yan-Feng Liu, Gang Du, Zhi-Hao Fu, Hao Zhang, Jin-Huan Yang, Ben Wang, Xiu-Yu Wang, Bin Jin

Abstract

AIM

To prospectively evaluate the effect of local wound infiltration with ropivacaine on postoperative pain relief and stress response reduction after

Match Overview

1	Internet 79 words crawled on 06-Jul-2016 www.wjgnet.com	3%
2	Crossref 34 words Kawaguchi, Yoshikuni, Yuichiro Otsuka, Hironori Kaneko, Motoki Nagai, Yukihiro Nomura, Mariko Yamamoto, M	1%
3	Internet 29 words crawled on 12-Mar-2016 ar.iiarjournals.org	1%
4	Crossref 27 words Xin, Yu, LiZhe Yong, and Yu Hong. "Efficacy of Postoperative Continuous Wound Infiltration With Local Anesthetics ...	1%
5	Crossref 18 words N. B. Scott. "Wound infiltration for surgery", <i>Anaesthesia</i> , 04/2010	1%
6	Internet 16 words crawled on 19-Jul-2017 www.jssm.org	1%
7	Crossref 16 words Calveley, Victoria L., Mohammad A. Khan, Ivan W. T. Yeung, Jake Vandyk, and Richard P. Hill. "Partial volume rat l	1%
8	Internet 15 words crawled on 16-Mar-2014 bj.a.oxfordjournals.org	<1%
9	Internet 12 words crawled on 05-Nov-2015 www.hormones.gr	<1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 3,950 条结果 (用时 0.62 秒)

Efficacy of Postoperative Continuous Wound Infiltration With Local ...

<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)作者: Y Xin - 2014 - 被引用次数: 3 - [相关文章](#)

2014年6月18日 - Forty patients undergoing open hepatectomy were enrolled in this ... Surgical wound infusion with ropivacaine after hepatectomy can ... Control of postoperative pain is of interest to surgeons, anesthetists, hospital administrators, and patients. ... and shorter half-life to reduce the risk of plasma accumulation.

Postoperative pain control using continuous i.m. bupivacaine infusion ...

<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)作者: EM Wong-Lun-Hing - 2014 - 被引用次数: 8 - [相关文章](#)

2013年10月23日 - Effective postoperative pain control will reduce the incidence of numerous ... However, the use of epidural analgesia after hepatectomy is still subject to debate. ... pain is the continuous infiltration of local anaesthetic using wound possibility that the local analgesic effect of the bupivacaine infusion may ...

[PDF] Local Anaesthetic Wound Infusion Catheters - Discipline of ...

anaesthetics.ukzn.ac.za/.../FMM_No6_6-3-15_Local_Anaesthetics_Wound_... - [翻译此页](#)

2015年3月6日 - Pain is relieved by Continuous Local Anaesthetic wound infusion into a ... nerve fibres and a decrease of the local inflammatory response to injury¹ related side effects, a reduction in PONV in 4 of 6 RCTs, although not and continuous wound infiltration for postoperative pain management after open.

[PDF] A comparison of pre- and post-surgical administration of ... - CiteSeerX

citeseerx.ist.psu.edu/viewdoc/download;jsessionid=...?doi=10.1.1... - [翻译此页](#)作者: JH Hayes - 1999 - 被引用次数: 26 - [相关文章](#)

analgesic treatment was a significant factor in the reduction in body weight ... administration of analgesics had no influence on the effect of bupivacaine ... operative pain and on the surgical stress response. Blocking of this afferent input by the use of local anaesthesia prior to surgery ... infiltration of a laparotomy wound with.

[PDF] A wound infiltration as a method of postoperative analgesia

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 3,950 条结果 (用时 0.48 秒)

Postoperative pain control using continuous i.m. bupivacaine infusion ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: EM Wong-Lun-Hing - 2014 - 被引用次数: 8 - 相关文章

2013年10月23日 - Effective postoperative pain control will reduce the incidence of numerous ...

However, the use of epidural analgesia after hepatectomy is still subject to debate. ... pain is the continuous infiltration of local anaesthetic using wound possibility that the local analgesic effect of the bupivacaine infusion may ...

Efficacy of Postoperative Continuous Wound Infiltration With Local ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: Y Xin - 2014 - 被引用次数: 3 - 相关文章

2014年6月18日 - Forty patients undergoing open hepatectomy were enrolled in this ... Surgical wound infusion with ropivacaine after hepatectomy can ... Control of postoperative pain is of interest to surgeons, anesthetists, hospital administrators, and patients. ... and shorter half-life to reduce the risk of plasma accumulation.

[PDF] Local Anaesthetic Wound Infusion Catheters - Discipline of ...

anaesthetics.ukzn.ac.za/.../FMM_No6_6-3-15_Local_Anaesthetics_Wound_... - 翻译此页

2015年3月6日 - Pain is relieved by Continuous Local Anaesthetic wound infusion into a ... nerve fibres and a decrease of the local inflammatory response to injury¹ related side effects, a reduction in PONV in 4 of 6 RCTs, although not and continuous wound infiltration for postoperative pain management after open.

[PDF] A wound infiltration as a method of postoperative analgesia

https://bib.irb.hr/datoteka/430938.kvolik_PB_241_2461.pdf - 翻译此页

作者: S Kvolik - 2009 - 被引用次数: 6 - 相关文章

New local anaesthetics with long duration of action and low ... the postoperative pain treatment is characterized by the use of different analge- ... oped to improve postoperative analgesia, reduce opioid consumption ... hip and knee replacement, after shoulder surgery and af- ... CFNB: 0.2% of ropivacaine at 7 mL/hr.

缺少字词: stress response hepatectomy

全部

图片

新闻

视频

购物

更多

设置

工具

找到约 3,960 条结果 (用时 1.45 秒)

Efficacy of Postoperative Continuous Wound Infiltration With Local ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: Y Xin - 2014 - 被引用次数: 5 - 相关文章

2014年6月18日 - Forty patients undergoing **open hepatectomy** were enrolled in this ... Surgical **wound** infusion with **ropivacaine** after hepatectomy can ... **Control** of **postoperative pain** is of interest to surgeons, anesthetists, hospital administrators, and patients. ... and shorter half-life to **reduce** the risk of plasma accumulation.

Postoperative pain control using continuous i.m. bupivacaine infusion ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: EM Wong-Lun-Hing - 2014 - 被引用次数: 8 - 相关文章

2013年10月23日 - Effective **postoperative pain control** will **reduce** the incidence of numerous ... However, the use of epidural analgesia **after hepatectomy** is still subject to debate. ... pain is the continuous **infiltration** of **local** anaesthetic using **wound** possibility that the **local** analgesic **effect** of the bupivacaine infusion may ...

[PDF] Local Anaesthetic Wound Infusion Catheters - Discipline of ...

anaesthetics.ukzn.ac.za/.../FMM_No6_6-3-15_Local_Anaesthetics_Wound_... ▼ 翻译此页

2015年3月6日 - Pain is relieved by Continuous **Local** Anaesthetic wound infusion into a ... nerve fibres and a decrease of the **local** inflammatory **response** to injury¹ related side **effects**, a **reduction** in PONV in 4 of 6 RCTs, although not and continuous **wound infiltration** for **postoperative pain management** after open.

[PDF] A comparison of pre- and post-surgical administration of ... - CiteSeerX

<citeseerx.ist.psu.edu/viewdoc/download;jsessionid...?doi=10.1.1...> ▼ 翻译此页

作者: JH Hayes - 1999 - 被引用次数: 26 - 相关文章

analgesic **treatment** was a significant factor in the **reduction** in body weight ... administration of