



Name of Journal: *World Journal of Gastrointestinal Oncology*

Manuscript NO: 35768

Manuscript Type: Original Article

Clinical Practice Study

Neoadjuvant hyperfractionated accelerated radiotherapy plus concomitant 5-FU infusion in locally advanced rectal cancer: A phase II study

Zeynep Gural, Sezer Saglam, Serap Yucel, Esra Kaytan-saglam, Oktar Asoglu, Cetin Ordu, Hediye Acun, Rasul Sharifov, Semen Onder, Ahmet Kizir, Ethem N Oral

Abstract

AIM

In this phase II study, we evaluated the efficacy and tolerability of neoadjuvant hyperfractionated accelerated radiotherapy (HART) and concurrent chemotherapy in

Match Overview

1	Internet 31 words crawled on 03-Jan-2010 www.ncbi.nlm.nih.gov	1%
2	Internet 19 words crawled on 13-Jan-2017 docplayer.net	1%
3	Publications 17 words Patel, Anush Puthillath, Ajithkumar Yang. "Neoadjuvant c... emoradiation for rectal cancer: is more better?(Clinical repo	1%
4	Internet 13 words crawled on 13-Mar-2017 repub.eur.nl	<1%
5	Crossref 11 words Lotz, J.P.. "Treatment of unresectable hepatocellular carcin oma with a combination of human recombinant @a-2b int...	<1%
6	Crossref 10 words Diaz-Gonzalez, J.A. "Preoperative chemoradiation with ora	<1%

学术搜索

文章

我的图书馆

时间不限

2017以来

2016以来

2013以来

自定义范围...

[HTML] Neoadjuvant hyperfractionated-accelerated radiotherapy with concomitant chemotherapy in esophageal cancer: phase II study

[S Saglam](#), [A Arifoglu](#), [EK Saglam](#), [F Tunca](#)... - Journal of ..., 2013 - ncbi.nlm.nih.gov

Purpose Concomitant use of chemotherapy and a radiation dose schedule that is more efficient compared to conventional radiotherapy may provide better outcomes in patients with esophageal cancer. This study aimed to assess the efficacy and tolerability of neoadjuvant cisplatin-based chemotherapy and hyperfractionated accelerated radiotherapy regimen in this group of patients.

被引用次数: 4 相关文章 所有 10 个版本 引用 保存

以上显示的是最相近的搜索结果。 [查看全部搜索结果](#)

按相关性排序

按日期排序

[关于 Google 学术搜索](#)[隐私权](#)[条款](#)[提供反馈](#)

不限语言

中文网页

简体中文网页

☒ 包括专利☐ 包含引用



Neoadjuvant hyperfractionated accelerated radiotherapy plus concomitant



全部

新闻

图片

购物

地图

更多

设置

工具

找到约 29,200 条结果 (用时 0.61 秒)

CPT-11 and concomitant hyperfractionated accelerated radiotherapy ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2360515/> - 翻译此页

作者 : V Voelter - 2006 - 被引用次数 : 22 - 相关文章

2006年9月12日 - In a prior phase I trial, we established the **neoadjuvant** regimen of 3 weekly ... Patients with **locally advanced rectal cancer** stage cT3/4 and/or cN+ without to **infusion** of 5FU alone in patients with stage II and III colon cancer (Andre **Phase II study** of **radiotherapy plus** concurrent irinotecan (CPT-11) and ...

Neoadjuvant capecitabine, bevacizumab and radiotherapy for locally ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4229928/> - 翻译此页

作者 : Y Miki - 2014 - 被引用次数 : 3 - 相关文章

2014年8月16日 - Patients with cT3/T4 **rectal cancer** were eligible. ... as preoperative CRT for Japanese patients with **locally advanced rectal cancer**. ... is non-inferior to 5-FU as a component of **neoadjuvant** RT for **rectal cancer** [12, 14, 15]. A **phase II study** of **neoadjuvant** bevacizumab **plus** capecitabine and **concomitant** ...

Rectal Cancer: Adjuvant Therapy for Rectal Cancer - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2789506/> - 翻译此页

作者 : SS Krishnamurthi - 2007 - 被引用次数 : 11 - 相关文章

Keywords: **Rectal cancer**, radiation, chemotherapy, adjuvant, review ... This **study** used a radiation dose of 45 Gy **plus** a 5.4 Gy boost. ... arms consisted of bolus **infusions** of 5-FU during the first 3 and last 3 days of **radiation therapy**. One small **phase II** trial tested a **hyperfractionated** boost following 45 Gy of conventional ...

Phase I Trial of Capecitabine and Weekly Irinotecan in Combination ...

ascopubs.org/doi/full/10.1200/jco.2005.04.171 - 翻译此页

作者 : RD Hofheinz - 2005 - 被引用次数 : 107 - 相关文章

[全部](#)[新闻](#)[图片](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 34,000 条结果 (用时 0.69 秒)

Neoadjuvant accelerated concomitant boost radiotherapy and ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/21555930> ▼ 翻译此页

作者: L Caravatta - 2012 - 被引用次数: 18 - 相关文章

Neoadjuvant accelerated concomitant boost radiotherapy and multidrug chemotherapy in locally advanced rectal cancer: a dose-escalation study. ... (55 Gy/5 wk) plus raltitrexed (cohort B), standard radiotherapy plus raltitrexed and ... On the basis of these results, this intensified regimen could be tested in a phase II study.

CPT-11 and concomitant hyperfractionated accelerated radiotherapy ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2360515/> - 翻译此页

作者: V Voelter - 2006 - 被引用次数: 22 - 相关文章

2006年9月12日 - 5-Fluorouracil (5FU) has been the only approved chemotherapy agent for ... neoadjuvant regimen of 3 weekly doses of CPT-11 with concomitant HART ... Patients with locally advanced rectal cancer stage cT3/4 and/or cN+ without Phase II study of radiotherapy plus concurrent irinotecan (CPT-11) and ...

CAPIRI-IMRT: a phase II study of concurrent capecitabine and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4353448/> ▼ 翻译此页

作者: G Cai - 2015 - 被引用次数: 5 - 相关文章

2015年2月28日 - Seventy-one patients diagnosed with recurrent rectal cancer who did not (range, 2–12) of 5-FU plus oxaliplatin or 6 cycles (range, 6–8) capecitabine plus oxaliplatin. to neoadjuvant chemoradiation for the treatment of locally advanced A Phase II study of preoperative radiotherapy and concomitant ...

Abeloff's Clinical Oncology E-Book - 第 1359 页 - Google 图书结果

<https://books.google.ae/books?isbn=1455728810> - 翻译此页

John E. Niederhuber, James O. Armitage, James H Doroshow - 2013 - Medical

Preoperative hyperfractionated accelerated radiotherapy (HART) in locally advanced rectal ... of patients with