

Risk  
factors  
for  
**surgical  
site  
infection  
after  
elective  
resection  
of  
the  
colon  
and  
rectum:**  
a  
single-  
center  
prospective  
study  
of  
2,809  
consecutive  
patients

R Tang,  
HH  
Chen,  
YL  
Wang,  
CR  
Changchien... -  
... of  
**surgery,**  
2001  
-  
[ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/11711111/)  
...

22  
Allogeneic  
leukocytes  
have  
a  
critical  
**role**  
in  
the  
induction  
of  
transfusion-  
induced  
...  
space  
SSI  
with  
or  
without

leakage  
is  
of  
clinical  
and  
pathogenetic  
**importance**  
because  
the

...  
Postoperative  
**infection**  
and  
natural  
killer  
cell  
**function**  
following  
blood  
transfusion  
in  
patients  
undergoing



Cited by 420  
Related articles  
All 9 versions

... of  
**oral**  
non-  
absorbable  
and  
intravenous  
**antibiotics**  
versus  
intravenous  
**antibiotics**  
alone  
in  
the  
**prevention**  
of  
**surgical**  
**site**  
**infections**  
**after**  
**colorectal**  
**surgery:**  
a  
meta- ...

CF  
Bellows,  
KT Mills,  
TN

Kelly,  
G  
Gagliardi -  
Techniques  
in  
coloproctology,  
2011  
-  
Springer

...  
of  
the  
studies  
that  
utilized  
only  
**oral**  
non-  
absorbable  
**antibiotics**  
is  
necessary  
in  
order  
to  
clarify  
their  
**role**  
in  
reducing

...  
In  
the  
RCT  
by  
Lewis  
and  
colleagues,  
the  
most  
**important**  
risk  
factor  
for  
wound  
**infection**  
was  
the  
presence  
of  
a

...  
**Importance**  
of  
the  
duration  
of  
the  
**surgical**  
procedure.

...



Cited by 64  
Related articles

[All 9 versions](#)

Antimicrobial  
prophylaxis  
in  
colorectal  
surgery:  
a  
systematic  
review  
of  
randomized  
controlled  
trials

F Song,  
AM  
Glenny -  
British  
Journal  
of  
**Surgery**,  
1998

-  
Wiley  
Online  
Library

...  
**antibiotic**  
regimes  
in  
the  
**treatment**  
of  
patients  
undergoing  
**elective**  
**colorectal**  
**surgery**  
in  
a  
district  
general  
hospital.

...  
Do  
preoperative  
**oral**  
**antibiotics**  
influence  
sepsis  
rates  
following  
**elective**  
**colorectal**  
**surgery**  
in  
patients  
receiving  
perioperative  
intravenous  
**prophylaxis**?

...



Cited by 435

[Related articles](#)[All 12 versions](#)

**Surgical  
site  
infections  
after  
colorectal  
surgery:  
do  
risk  
factors  
vary  
depending  
on  
the  
type  
of  
infection  
considered?**

J

Blumetti,

M

Luu,

G

Sarosi,

K

Hartless,

J

McFarlin... -

**Surgery,**

2007

-

Elsevier

...

4,

5,

6,

7,

8

and

11

Most

of

this

variation

is

probably

a

**function**

of

multiple

...

Although

there

is  
a  
generalized  
awareness  
and  
acceptance  
of  
the  
the **importance**  
of  
the  
choice  
of  
of **prophylactic**

...  
A  
second  
**important**  
issue  
is  
identification  
of  
an  
acceptable  
means  
of  
risk-  
adjusting  
SSI  
rates.

...



Cited by 133  
[Related articles](#)  
[All 9 versions](#)

**Preventing  
surgical  
site  
infections:  
a  
surgeon's  
perspective.**

RL  
Nichols -  
Emerging  
**infectious**  
diseases,  
2001  
-  
[ncbi.nlm.nih.gov](https://ncbi.nlm.nih.gov)

...  
This  
report  
stresses  
the  
**importance**  
of  
conducting  
a  
formal

epidemiologic  
investigation  
when  
a

...

**Oral**  
neomycin  
and  
erythromycin  
or  
metronidazole  
were  
combined  
with  
a  
perioperative  
parenteral  
**antibiotic**

...

It  
is  
**important**  
to  
use  
**antibiotic**  
agents  
with  
both  
facultative  
and  
anaerobic

...



Cited by 268  
[Related articles](#)  
[All 17 versions](#)

[Quality  
standard  
for  
antimicrobial  
prophylaxis  
in  
surgical  
procedures](#)

EP  
Dellinger,  
PA  
Gross,  
TL  
Barrett... -  
**Infection**  
Control  
& ...,  
1994  
-  
[cambridge.org](#)

...  
Recent  
reports  
have

confirmed  
the  
**importance**  
of  
this  
timing.20~21~28  
Indeed,  
pharmacokinetic  
data  
indicate  
the  
desirability

...  
no  
parenteral  
antimicrobial  
agent  
was  
administered,  
the  
form  
would  
be  
checked  
to  
indicate  
either  
that  
**oral**  
antimicrobial  
drugs

...  
**Prophylactic  
antibiotics.**

...



Cited by 605  
[Related articles](#)  
[All 24 versions](#)



Preoperative  
oral  
antibiotics  
reduce  
surgical  
site  
infection  
following  
elective  
colorectal  
resections

JA  
Cannon,  
LK  
Altom,  
RJ  
Deierhoi... -  
Diseases  
of  
the  
Colon ...,  
2012  
-  
journals.lww.com

...  
Skip  
Navigation  
Links  
Home  
>  
November  
2012  
-  
Volume  
55  
-  
Issue  
11  
>  
Preoperative  
**Oral**  
**Antibiotics**  
Reduce  
**Surgical**  
**Site**  
**Infection...**

...  
  
  
Cited by 85  
Related articles  
All 5 versions

Prophylactic  
oral  
antimicrobial  
agents  
in  
elective

colonic  
surgery:  
a  
controlled  
trial

J  
Goldring,  
W  
McNaught,  
A  
Scott,  
G  
Gillespie -  
The  
Lancet,  
1975  
-  
Elsevier

...  
Furthermore  
recent  
work7-  
10  
has  
highlighted  
the  
clinical  
**importance**  
of  
gram-  
negative  
anaerobic  
Bac

...  
**Oral**  
**Antibiotics**  
to  
Prevent  
**Surgical**  
**Site**  
**Infections**  
Following  
Colon  
**Surgery.**

...  
Quality  
assurance  
and  
the  
**role**  
of  
**infection**  
control:  
a  
retrospective  
study  
of  
hospital-  
acquired

...  
  
  
Cited by 210

[Related articles](#)[All 5 versions](#)

# Antimicrobial prophylaxis for surgical wounds: guidelines for clinical care

CP

Page,

JMA

Bohnen,

JR

Fletcher... -

... of

**Surgery,**

1993

-

[archsurg.jamanetwork.com](http://archsurg.jamanetwork.com)

...

of

NRC

classification

and

the

SENIC

risk

categories

seems

appropriate

as

a

guide

for

administering

prophy-

lactic

**antibiotics,**

for

directing

...

This

combination

is

a

particularly

**important**

consider-

ation

because

of

recent

enthusiasm

for

extending

prophy-

lactic

**antibiotic**

use  
to  
a  
...



Cited by 660  
Related articles  
All 8 versions

... of  
**antibiotic**  
**prophylaxis**  
in  
**elective**  
**colorectal**  
**surgery:**  
single  
dose  
vs  
3  
doses  
of  
a  
second-  
generation  
cephalosporin  
without  
metronidazole  
and  
**oral**  
**antibiotics**

S Fujita,  
N  
Saito,  
T  
Yamada,  
Y  
Takii,  
K  
Kondo... -

... of  
**surgery,**  
2007  
-  
jamanetwork.com

...  
Clarke  
JSCondon  
REBartlett  
JGGorbach  
SLNichols  
RLOchi  
S  
Preoperative  
**oral**  
**antibiotics**

reduce  
septic  
...  
bowel  
preparation  
for  
**elective  
colorectal  
surgery:**  
comparison  
among  
**oral,**  
systemic,  
and

...  
Coppa  
GFEng  
K  
Factors  
involved  
in  
**antibiotic**  
selection  
in  
**elective  
colon  
and  
rectal**

...



[Cited by 91](#)

[Related articles](#)

[All 17 versions](#)