

6
 Name of journal: *World Journal of Gastroenterology*

Manuscript NO: 35987

Manuscript Type: ORIGINAL ARTICLE

Basic Study

**miR-192-5p regulates lipid synthesis in non-alcoholic fatty liver disease
 through SCD-1 miR-192-5p regulates lipid synthesis through SCD-1**

Xiao-Lin Liu, Hai-Xia Cao, Bao-Can Wang, Feng-Zhi Xin, Rui-Nan Zhang, Da
 Zhou, Rui-Xu Yang, Ze-Hua Zhao, Qin Pan, Jian-Gao Fan

Abstract

AIM

To evaluate the levels of miR-192-5p in non-alcoholic fatty liver disease

Match Overview

1	Internet 21 words crawled on 15-Jun-2016 hrcak.srce.hr	<1%
2	Crossref 20 words Ding, Jiexia, Meng Li, Xingyong Wan, Xi Jin, Shaohua Chen, Chaohui Yu, and Youming Li. "Effect of miR-34a in l..."	<1%
3	Internet 16 words crawled on 14-Jul-2016 www.science.gov	<1%
4	Internet 16 words crawled on 12-Jun-2017 www.mdpi.com	<1%
5	Internet 15 words crawled on 17-Jul-2017 ajpgi.physiology.org	<1%
6	Internet 15 words crawled on 05-Oct-2017 real.mtak.hu	<1%
7	Internet 14 words crawled on 30-Aug-2017 jeb.biologists.org	<1%
8	Internet 14 words crawled on 26-Jun-2016 www.deepdyve.com	<1%

全部

图片

视频

新闻

更多

设置

工具

找到约 3,350 条结果 (用时 0.81 秒)

Downregulation of miR-192 causes hepatic steatosis and lipid ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/28483554> - 翻译此页

作者: Y Lin - 2017 - 被引用次数: 2 - 相关文章

1. Biochim Biophys Acta. 2017 May 5;1862(9):869-882. doi: 10.1016/j.bbailip.2017.05.001. [Epub ahead of print]. Downregulation of **miR-192** causes hepatic steatosis and **lipid** accumulation ... for bisphenol A-triggered **non-alcoholic fatty liver disease**. ... Using HepG2 cells, we further confirmed that **miR-192** directly acted on ...

Micro RNAs in the development of non-alcoholic fatty liver disease

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: GS Gerhard - 2015 - 被引用次数: 10 - 相关文章

2015年2月27日 - **Nonalcoholic fatty liver disease** (NAFLD) represents a spectrum of ... may **regulate** the development of hepatic steatosis **through** mechanisms involving the PPARA pathway. ... miRNAs including stearoyl-coenzyme A desaturase 1 (**Scd1**). ... Serum levels of **miR-34a**, **miR-122**, **miR-181a**, **miR-192**, **miR-200b**, ...

MicroRNAs as Biomarkers for Liver Disease and Hepatocellular ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: CN Hayes - 2016 - 被引用次数: 47 - 相关文章

2016年2月24日 - **miR-125a-5p** and **miR-1231** have been shown to directly target hepatitis B ... Figure 1. MicroRNA **synthesis** and function. MicroRNA genes are ... MicroRNAs from the **liver** may enter the serum passively **through** ... to be **up-regulated** in **liver** tissue, and **miR-122**, **miR-192**, **miR-155**, ... **Alcoholic Liver Disease**.

Molecular Pathogenesis of NASH - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: A Caligiuri - 2016 - 被引用次数: 17 - 相关文章

全部

图片

视频

新闻

更多

设置

工具

找到约 3,350 条结果 (用时 0.81 秒)

Downregulation of miR-192 causes hepatic steatosis and lipid ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/28483554> - 翻译此页

作者: Y Lin - 2017 - 被引用次数: 2 - 相关文章

1. Biochim Biophys Acta. 2017 May 5;1862(9):869-882. doi: 10.1016/j.bbali.2017.05.001. [Epub ahead of print]. Downregulation of **miR-192** causes hepatic steatosis and **lipid** accumulation ... for bisphenol A-triggered **non-alcoholic fatty liver disease**. ... Using HepG2 cells, we further confirmed that **miR-192** directly acted on ...

Micro RNAs in the development of non-alcoholic fatty liver disease

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: GS Gerhard - 2015 - 被引用次数: 10 - 相关文章

2015年2月27日 - **Nonalcoholic fatty liver disease** (NAFLD) represents a spectrum of ... may **regulate** the development of hepatic steatosis **through** mechanisms involving the PPARA pathway. ... miRNAs including stearoyl-coenzyme A desaturase 1 (**Scd1**). ... Serum levels of **miR-34a**, **miR-122**, **miR-181a**, **miR-192**, **miR-200b**, ...

MicroRNAs as Biomarkers for Liver Disease and Hepatocellular ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: CN Hayes - 2016 - 被引用次数: 47 - 相关文章

2016年2月24日 - **miR-125a-5p** and **miR-1231** have been shown to directly target hepatitis B ... Figure 1. MicroRNA **synthesis** and function. MicroRNA genes are ... MicroRNAs from the **liver** may enter the serum passively **through** ... to be **up-regulated** in **liver** tissue, and **miR-122**, **miR-192**, **miR-155**, ... **Alcoholic Liver Disease**.

Molecular Pathogenesis of NASH - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: A Caligiuri - 2016 - 被引用次数: 17 - 相关文章



miR-192-5p regulates lipid synthesis in non-alcoholic fatty liver disease throug



文章

时间不限

2017以来

2016以来

2013以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☐ 包括专利

☐ 包含引用

☒ 创建快讯

您的搜索 - **miR-192-5p regulates lipid synthesis in non-alcoholic fatty liver disease through SCD-1** miR-192-5p regulates lipid synthesis through SCD-1 - 与所有文章均不相符。

建议:

请检查输入字词有无错误。

请尝试其他的查询词

请改用较常见的字词。

请减少查询字词的数量。

[请向所有网络查询](#)