


学术搜索

获得 6 条结果 (用时0.03秒)

 我的著作引用情况

文章

我的图书馆

时间不限

2017以来

2016以来

2013以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☒ 包括专利☒ 包含引用☒ 创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Nuclear factor-κB—a pivotal transcription factor in chronic inflammatory diseases

PJ Barnes, M Karin - New England journal of medicine, 1997 - Mass Medical Soc

... alpha is not phosphorylated by Raf-1 or protein kinase C isozymes, but is a casein kinase II substrate. ... Cyclooxygenase-1 and -2 expression in rheumatoid synovial tissues: effects of interleukin-1β, phorbol ester, and corticosteroids. ... Adhesion molecules and inflammatory injury. ...

被引用次数: 5048 相关文章 所有 13 个版本 引用 保存

[PDF] academia.edu

The pathophysiology and treatment of sepsis

RS Hotchkiss, IE Karl - New England Journal of Medicine, 2003 - Mass Medical Soc

... Liver dysfunction rarely progresses to hepatic encephalopathy. ... bleeding episode, or a requirement for 3 or more units of blood), as compared with 2 percent of ... the criteria for sepsis specified by the Acute Physiology and Chronic Health Evaluation (APACHE II) scoring system. ...

被引用次数: 4153 相关文章 所有 24 个版本 引用 保存

[PDF] free.fr

Neuroprotective effects of geniposide on Alzheimer's disease pathology

WZ Liu, G Li, C Hölscher, L Li - Reviews in the Neurosciences, 2015 - degruyter.com

... After the deglycosylation of geniposide in the liver or intestine, genipin would undergo redox or phase II metabolism immediately ... and the AUC 0→4 h values in tissues were in the order of kidney>spleen>liver>heart>lung ... (2) Geniposide also suppresses the induction of apoptosis ...

被引用次数: 18 相关文章 所有 6 个版本 引用 保存

[PDF] semanticscholar.org

[HTML] A novel small molecule TLR4 antagonist (IAXO-102) negatively regulates non-hematopoietic toll like receptor 4 signalling and inhibits aortic aneurysms ...

C Huggins, S Pearce, F Peri, F Neumann, G Cockerill... - Atherosclerosis, 2015 - Elsevier

... 102-FITC was present in the bladder, small and large intestine, to some extent, in kidney or liver (data are ... pretreatment inhibited, to varying extent, the expression of 14/22 Angiotensin II driven proinflammatory proteins (Table 2 Supplement). ... Ang II/IAXO-102, 10, 0, 30, 1.05 ± 0.18 ...

被引用次数: 8 相关文章 所有 8 个版本 引用 保存

[HTML] sciencedirect.com

Potential Signaling Pathways Involved in the Clinical Application of Oxymatrine

ML Lu, XH Xiang, SH Xia - Phytotherapy Research, 2016 - Wiley Online Library

16

Name of journal: *World Journal of Gastroenterology*

Manuscript No.: 36223

Manuscript Type: Original Article

Basic Study

β -arrestin 2 attenuates lipopolysaccharide-induced liver injury via inhibition of TLR4/NF- κ B signaling pathway-mediated inflammation in mice

Jiang MP *et al.* β -arrestin 2 attenuates LPS-induced liver injury via inhibition of inflammation

Match Overview

1	Internet 119 words crawled on 22-Jul-2017 www.wjgnet.com	3%
2	Internet 48 words crawled on 07-Sep-2017 www.jbc.org	1%
3	Internet 38 words crawled on 27-Oct-2017 www.spandidos-publications.com	1%
4	Crossref 36 words N Suleymanova, C Crudden, T Shibano, C Worrall, I Oprea, A Tica, G A Calin, A Girmata, L Girmata. "Functional anta	1%
5	Crossref 29 words Wen, Shun-hang, Hu-jun Wu, Li Lin, Lei Chong, Li-li Zhu, Wei-xi Zhang, Hai-lin Zhang, and Chang-chong Li. "Adju	1%
6	Internet 28 words crawled on 18-Oct-2017 www.mdpi.com	1%
7	Internet 24 words crawled on 08-Dec-2009 www.freepatentsonline.com	1%

时间不限

2017 以来

2016 以来

2013 以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

☐ 创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Nuclear factor- κ B—a pivotal transcription factor in chronic inflammatory diseases

[PDF] academia.edu

PJ Barnes, M Karin - New England journal of medicine, 1997 - Mass Medical Soc

... alpha is not phosphorylated by Raf-1 or protein kinase C isozymes, but is a casein kinase II substrate ... Cyclooxygenase-1 and -2 expression in rheumatoid synovial tissues: effects of interleukin-1 β , phorbol ester, and corticosteroids ... Adhesion molecules and inflammatory injury ...

☆ 99 被引用次数: 5098 相关文章 所有 13 个版本

The pathophysiology and treatment of sepsis

[PDF] free.fr

RS Hotchkiss, IE Karl - New England Journal of Medicine, 2003 - Mass Medical Soc

... Liver dysfunction rarely progresses to hepatic encephalopathy ... bleeding episode, or a requirement for 3 or more units of blood), as compared with 2 percent of ... the criteria for sepsis specified by the Acute Physiology and Chronic Health Evaluation (APACHE II) scoring system ...

☆ 99 被引用次数: 4220 相关文章 所有 24 个版本

Neuroprotective effects of geniposide on Alzheimer's disease pathology

[PDF] semanticscholar.org

WZ Liu, G Li, C Hölscher, L Li - Reviews in the Neurosciences, 2015 - degruyter.com

... After the deglycosylation of geniposide in the liver or intestine, genipin would undergo redox or phase II metabolism immediately ... and the AUC 0→4 h values in tissues were in the order of kidney>spleen>liver>heart>lung ... (2) Geniposide also suppresses the induction of apoptosis ...

☆ 99 被引用次数: 20 相关文章 所有 6 个版本

[HTML] A novel small molecule TLR4 antagonist (IAXO-102) negatively regulates non-hematopoietic toll like receptor 4 signalling and inhibits aortic aneurysms ...

[HTML] sciencedirect.com

C Huggins, S Pearce, F Peri, F Neumann, G Cockerill... - Atherosclerosis, 2015 - Elsevier

... 102-FITC was present in the bladder, small and large intestine, to some extent, in kidney or liver (data are ... pretreatment inhibited, to varying extent, the expression of 14/22 Angiotensin II driven proinflammatory proteins (Table 2 Supplement) ... Ang II/IAXO-102, 10, 0, 30, 1.05 \pm 0.18 ...

找到约 2,670 条结果 (用时 0.47 秒)

β-Arrestin 2 Negatively Regulates Toll-like Receptor 4 (TLR4 ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4132806/> - 翻译此页

作者: H Li - 2014 - 被引用次数: 17 - 相关文章

2014年7月10日 - Deficiency of β-arrestin 2 and inhibition of p38 MAPK activity both ... amount of IL-10, and are more susceptible to LPS-induced septic shock which is ... which β-arrestin 2 negatively regulates TLR4-mediated inflammatory reactions. ... and attenuation of G protein-coupled receptor (GPCR) signaling through ...

Chikusetsusaponin V attenuates lipopolysaccharide-induced liver ...

www.tandfonline.com/doi/abs/10.3109/08923973.2016.1153109 ▼ 翻译此页

作者: YW Dai - 2016 - 被引用次数: 4 - 相关文章

Chikusetsusaponin V attenuates lipopolysaccharide-induced liver injury in mice ... to inhibit inflammatory responses in lipopolysaccharide (LPS)-induced macrophage cells ... the anti-inflammatory effects of CsV on LPS-induced acute liver injury in mice ... responses, partly via inhibiting NF-κB and MAPK signaling pathways.

S-Adenosylmethionine attenuates lipopolysaccharide-induced liver ...

europepmc.org/abstract/med/26202509 ▼ 翻译此页

作者: P Li - 2014 - 被引用次数: 9 - 相关文章

S-Adenosylmethionine attenuates lipopolysaccharide-induced liver injury by downregulating the Toll-like receptor 4 signal in Kupffer cells. Finally, SAM did not affect NF-κB-p65 translocation into the nucleus ($p > 0.05$), but significantly inhibited ... mediated by suppressing activation of the TLR4/MAPK signaling pathway.

The negative regulation of Toll-like receptor and associated pathways ...

www.nature.com > immunology & cell biology > review ▼

作者: T Lang - 2007 - 被引用次数: 117 - 相关文章

2007年7月10日 - Mal also associates with TRAF-6 to mediate transactivation of NF-κB. ... attenuate TLR signalling and therefore prevent acute inflammatory Their studies found by using β-arrestin-1 and β-arrestin-2 siRNA in HEK-TLR4 stable cells in the liver against LPS-induced liver failure, as mice injected with an ...

Beta-arrestin 2 negatively regulates sepsis-induced inflammation - Fan

onlinelibrary.wiley.com/doi/10.1111/j.1365-2567.2009.03185.x/full ▼ 翻译此页

作者: H Fan - 2010 - 被引用次数: 55 - 相关文章

2010年5月4日 - Binding to TLR4 leads to activation of a series of signalling proteins leading to ... 1 and