

找到约 2,590 条结果 (用时 0.65 秒)

Google 学术: Early Gastric Cancer Frequently Has High Expression of KK-LC-1, a Cancer-Testis Antigen

Cancer/testis (CT) antigens: potential targets for ... - Caballero - 被引用次数: 448

... of cancer/testis antigens expression in patients with ... - Shigematsu - 被引用次数: 61

... significance of expression of cancer/testis antigen and ... - Hanagiri - 被引用次数: 26

Frequent High Expression of Kita-Kyushu Lung Cancer Antigen-1 (KK ...

ar.iiarjournals.org/content/35/6/3575.full - 翻译此页

作者: A SHIDA - 2015 - 被引用次数: 3 - 相关文章

The present study demonstrated that KK-LC-1 is expressed in gastric cancer. Materials and Methods: We analyzed the expression of KK-LC-1 and cancer/testis antigens ... therapeutic approaches for treating patients with advanced gastric cancer. Hence, KK-LC-1 may have other epitope peptides that are recognized by ...

Correlation Between Expression of the Cancer/Testis Antigen KK-LC-1 ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5461451/> - 翻译此页

作者: T FUKUYAMA - 2017 - 相关文章

2017年5月3日 - Advanced - Journal list - Help ... KK-LC-1 was frequently expressed in gastric cancer caused by H. pylori infection. ... We previously identified Kita-kyushu lung cancer antigen-1 (KK-LC-1) as a new CTA that has an In only one case was KK-LC-1 not expressed despite a high IgG titer against H. pylori.

Clinical significance of cancer/testis antigens expression in patients ...

[www.lungcancerjournal.info/article/S0169-5002\(09\)00307-9/fulltext](http://www.lungcancerjournal.info/article/S0169-5002(09)00307-9/fulltext) - 翻译此页

作者: Y Shigematsu - 2010 - 被引用次数: 60 - 相关文章

2009年6月22日 - Advanced Search RNA isolation and RT-PCR analysis of cancer/testis antigens; 2.4. ... were expressed more frequently in squamous cell carcinoma than in ... KK-LC-1 was expressed in 32.1% of patients with adenocarcinoma and in ... such cancers from lung, breast, esophagus, stomach and colon.

Clinical significance of cancer/testis antigens expression in patients ...

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 36404

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Early gastric cancer frequently has high expression of KK-LC-1, a cancer-testis antigen

Futawatari N *et al.* CTA's clinical significance in gastric cancer

Nobue Futawatari, Takashi Fukuyama, Rui Yamamura, Akiko Shida, Yoshihito Takahashi, Yatsushi Nishi, Yoshinobu Ichiki, Noritada Kobayashi, Hitoshi Yamazaki, Masahiko Watanabe

Abstract

AIM

To assess cancer-testis antigens (CTAs) expression in gastric cancer patients and

Match Overview

1	Internet 26 words crawled on 31-Aug-2017 preview-jecr.biomedcentral.com	1%
2	Internet 18 words crawled on 01-Mar-2017 repub.eur.nl	1%
3	Publications 15 words "EGF RECEPTOR FOUND TO REGULATE MACROPHAGE... INFLAMMATION IN GUT.", States News Service, Oct 13 201	<1%
4	Crossref 14 words Matthias Wölfl. "Expression of MHC class I, MHC class II, and d cancer germline antigens in neuroblastoma", Cancer Imm	<1%
5	Internet 14 words crawled on 05-Oct-2017 real.mtak.hu	<1%
6	Internet 12 words crawled on 07-May-2016 f1000.com	<1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 2,510 条结果 (用时 0.74 秒)

Correlation Between Expression of the Cancer/Testis Antigen KK-LC-1 ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5461451/> - 翻译此页

作者: T FUKUYAMA - 2017 - 相关文章

2017年5月3日 - **Advanced** - Journal list - Help ... **KK-LC-1** was frequently expressed in gastric cancer caused by *H. pylori* infection. ... We previously identified Kita-kyushu lung cancer antigen-1 (**KK-LC-1**) as a new CTA that has an In only one case was **KK-LC-1** not expressed despite a high IgG titer against *H. pylori*.

Frequent High Expression of Kita-Kyushu Lung Cancer Antigen-1 - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/26026129> - 翻译此页

作者: A Shida - 2015 - 被引用次数: 3 - 相关文章

Frequent High Expression of Kita-Kyushu Lung Cancer Antigen-1 (**KK-LC-1**) in ... lung cancer antigen-1 (**KK-LC-1**) has been reported as not being expressed in ... cancer/testis antigens (CTAs) in surgical specimens of 49 gastric carcinomas.

缺少字词: early

Frequent High Expression of Kita-Kyushu Lung Cancer Antigen-1 (KK ...

<ar.iiarjournals.org/content/35/6/3575.full> - 翻译此页

作者: A SHIDA - 2015 - 被引用次数: 3 - 相关文章

The present study demonstrated that **KK-LC-1** is expressed in gastric cancer. ... Methods: We analyzed the expression of **KK-LC-1** and cancer/testis antigens (CTAs) in ... therapeutic approaches for treating patients with advanced gastric cancer. ... TAA designated Kita-Kyushu lung cancer antigen-1 (**KK-LC-1**), which has an ...

Identification of a New Cancer/Germline Gene, KK-LC-1, Encoding an ...

<cancerres.aacrjournals.org/content/66/9/4922> - 翻译此页

作者: T Fukuyama - 2006 - 被引用次数: 38 - 相关文章

2006年5月1日 - **Advanced Search** ... A 9-mer peptide (**KK-LC-176-84**; RQKRILVNL) was identified as an epitope peptide. ... of this gene was only expressed in the testis among normal tissues. ... The new antigen, designated as Kita-kyushu lung cancer antigen 1 ... The method used to establish the lung cancer cell line has ...

找到约 2,580 条结果 (用时 0.52 秒)

Correlation Between Expression of the Cancer/Testis Antigen KK-LC-1 ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5461451/> - 翻译此页

作者: T FUKUYAMA - 2017 - 相关文章

2017年5月3日 - **Advanced** - Journal list - Help ... KK-LC-1 was frequently expressed in gastric cancer caused by *H. pylori* infection. ... We previously identified Kita-kyushu lung cancer antigen-1 (KK-LC-1) as a new CTA that has an In only one case was KK-LC-1 not expressed despite a high IgG titer against *H. pylori*.

Frequent High Expression of Kita-Kyushu Lung Cancer Antigen-1 (KK ...

<ar.iiarjournals.org/content/35/6/3575.full> - 翻译此页

作者: A SHIDA - 2015 - 被引用次数: 3 - 相关文章

The present study demonstrated that KK-LC-1 is expressed in gastric cancer. ... Methods: We analyzed the expression of KK-LC-1 and cancer/testis antigens (CTAs) in ... therapeutic approaches for treating patients with advanced gastric cancer. ... TAA designated Kita-Kyushu lung cancer antigen-1 (KK-LC-1), which has an ...

Identification of a New Cancer/Germline Gene, KK-LC-1, Encoding an ...

<cancerres.aacrjournals.org/content/66/9/4922> ▼ 翻译此页

作者: T Fukuyama - 2006 - 被引用次数: 38 - 相关文章

2006年5月1日 - **Advanced** Search ... A 9-mer peptide (KK-LC-176-84; RQKRILVNL) was identified as an epitope peptide. ... of this gene was only expressed in the testis among normal tissues. ... The new antigen, designated as Kita-kyushu lung cancer antigen 1 ... The method used to establish the lung cancer cell line has ...

Cancer/testis antigen expression and autologous humoral immunity to ...

<cancerimmunolres.aacrjournals.org/content/canimmarch/4/1/11.figures-only> - 翻译此页

NY-ESO-1 expression detected by RT-PCR in gastric cancer specimens. ... SSX-4, and CT10 tended to be expressed more frequently in advanced stages.

Lung cancer-associated tumor antigens and the ... - Springer Link

<https://link.springer.com/content/pdf/10.1007/s11748-008-0433-6.pdf> ▼ 翻译此页

作者: K Yasumoto - 2009 - 被引用次数: 37 - 相关文章

Especially, patients with unresectable lung cancer have a high mor- ... apy in both early and advanced stage cancer patients.7 KK-LC-1 is also expressed in breast, esophageal, gastric, ... Table 3 Expression of cancer-testis antigens in non-small-cell lung cancer ... that tumor cells often cause a loss or down-regulation of.