



时间不限

2017以来

2016以来

2013以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

包括专利

包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

... performance and description of morphological features of focal nodular hyperplasia in Gd-EOB-DTPA-enhanced liver magnetic resonance imaging: results of a ...

CJ Zech, L Grazioli, J Breuer, MF Reiser... - Investigative ..., 2008 - journals.lww.com

Objectives: The aim of this prospective study was to evaluate the diagnostic performance of magnetic.

☆ 99 被引用次数: 227 相关文章 所有 7 个版本

[HTML] ... hepatobiliary phase MR imaging in the diagnosis of early-stage hepatocellular carcinoma: Comparison with dynamic triple-phase multidetector CT imaging

[HTML] wiley.com

H Haradome, L Grazioli, R Tinti... - ... Resonance Imaging, 2011 - Wiley Online Library

... detection technique (CARE Bolus; Siemens) to adjust optimal scan delay timing for ... Therefore, added interpretations of hepatobiliary phase images could significantly improve the diagnosis of ... lesions showing isointensity during portal venous or equilibrium phases, which could ...

☆ 99 被引用次数: 92 相关文章 所有 9 个版本

Qualitative and quantitative evaluation of hepatocellular carcinoma and cirrhotic liver enhancement using Gd-EOB-DTPA

BB Frericks, C Loddenkemper... - American Journal of ..., 2009 - Am Roentgen Ray Soc

... have reported a relevant number of isointense, and hence hardly visible, HCCs during equilibrium or delayed phases after gadopentetate ... to a slight overdosing in all but one patient (24-36 μmol/kg of body weight), and arterial phase imaging started at a fixed delay of 15 ...

☆ 99 被引用次数: 151 相关文章 所有 4 个版本

Hepatocellular carcinoma in a North American population: Does hepatobiliary MR imaging with Gd-EOB-DTPA improve sensitivity and confidence for diagnosis?

Name of journal: *World Journal of Gastroenterology*

Manuscript NO: 36562

Manuscript Type: Original Article

Retrospective study

Optimization of hepatobiliary phase delay time in ¹Gd-EOB-DTPA-enhanced magnetic resonance imaging for identification of hepatocellular carcinoma in patients with different severity of cirrhosis

Wu JW *et al.* Optimize HBP-DT of Gd-EOB-DTPA-enhanced MRI for HCC occurred in different cirrhosis

Jian-Wei Wu, Yue-Cheng Yu, Xian-Li Qu, Yan Zhang, Hong Gao

Abstract

AIM

To optimize the hepatobiliary phase delay time (HBP-DT) of ¹Gd-EOB-DTPA-enhanced

Match Overview

- | | | |
|---|---|-----|
| 1 | Crossref 209 words
"Abstracts of the 26th Annual Conference of APASL, February 15-19, 2017, Shanghai, China", <i>Hepatology International</i> | 4% |
| 2 | Internet 22 words
crawled on 24-Apr-2016
f1000.com | <1% |
| 3 | Internet 16 words
crawled on 07-Jan-2017
eprints.lib.okayama-u.ac.jp | <1% |
| 4 | Internet 16 words
crawled on 11-Nov-2017
pdfs.semanticscholar.org | <1% |

- 时间不限
- 2017以来
- 2016以来
- 2013以来
- 自定义范围...

- 按相关性排序
- 按日期排序

- 不限语言
- 中文网页
- 简体中文网页

- 包括专利
- 包含引用

- 创建快讯

小提示：只搜索中文(简体)结果，可在 学术搜索设置 指定搜索语言

... performance and description of morphological features of focal nodular hyperplasia in Gd-EOB-DTPA-enhanced liver magnetic resonance imaging: results of a ...

CJ Zech, L Grazioli, J Breuer, MF Reiser... - Investigative ..., 2008 - journals.lww.com
Objectives: The aim of this prospective study was to evaluate the diagnostic performance of magnetic.
☆ 99 被引用次数: 231 相关文章 所有 7 个版本

[HTML] ... hepatobiliary phase MR imaging in the diagnosis of early-stage hepatocellular carcinoma: Comparison with dynamic triple-phase multidetector CT imaging

[HTML] wiley.com

H Haradome, L Grazioli, R Tinti... - ... Resonance Imaging, 2011 - Wiley Online Library
... detection technique (CARE Bolus; Siemens) to adjust optimal scan delay timing for ... Therefore, added interpretations of hepatobiliary phase images could significantly improve the diagnosis of ... lesions showing isointensity during portal venous or equilibrium phases, which could ...
☆ 99 被引用次数: 94 相关文章 所有 9 个版本

Qualitative and quantitative evaluation of hepatocellular carcinoma and cirrhotic liver enhancement using Gd-EOB-DTPA

BB Frericks, C Loddenkemper... - American Journal of ..., 2009 - Am Roentgen Ray Soc
... have reported a relevant number of isointense, and hence hardly visible, HCCs during equilibrium or delayed phases after gadopentetate ... to a slight overdosing in all but one patient (24-36 μmol/kg of body weight), and arterial phase imaging started at a fixed delay of 15 ...
☆ 99 被引用次数: 153 相关文章 所有 4 个版本

Hepatocellular carcinoma in a North American population: Does hepatobiliary MR imaging with Gd-EOB-DTPA improve sensitivity and confidence for diagnosis?



找到约 13,900 条结果 (用时 0.67 秒)

Feasibility of 10-Minute Delayed Hepatocyte Phase Imaging ... - PLOS

journals.plos.org/plosone/article?id=10.1371/journal.pone.0167701 ▾ [翻译此页](#)

作者: I Jeon - 2016 - 被引用次数: 2 - [相关文章](#)

2016年12月9日 - Objectives To compare 10-minute (min) delayed hepatocyte **phase** imaging (HPI) using a 30° flip angle (FA) (10m-FA30) and 20-min delayed HPI using a 10° FA (20m-FA10) or 30° FA (20m-FA30) in **Gd-EOB-DTPA-enhanced MRI** in **patients** with chronic hepatitis or **cirrhosis**, in terms of lesion-to-liver ...

Diagnostic Accuracy of Gd-EOB-DTPA for Detection Hepatocellular ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5301959/> ▾ [翻译此页](#)

作者: M Imbriaco - 2017 - 被引用次数: 2 - [相关文章](#)

2017年2月1日 - The aim of the present study was to evaluate the diagnostic accuracy of hepato-**biliary** (HB) **phase MRI** with **Gd-EOB-DTPA**, compared with that of dynamic contrast-**enhanced magnetic resonance imaging** (DCEMRI) and dynamic contrast-**enhanced CT** (DCECT), for detection of HCC in **patients** with liver ...

Is 3-Tesla Gd-EOB-DTPA-Enhanced MRI with Diffusion-Weighted ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4223069/> - [翻译此页](#)

作者: B Maiwald - 2014 - 被引用次数: 15 - [相关文章](#)

2014年11月6日 - To compare 64-slice contrast-**enhanced** computed tomography (CT) with 3-Tesla **magnetic resonance imaging (MRI)** using **Gd-EOB-DTPA** for the diagnosis ... Based on the Child-Pugh-Classification, the **severity** of liver **cirrhosis** was classified as class A in 27 **patients**, class B in 16 **patients** and class C in 7 ...

Gd-EOB-DTPA enhanced MRI for hepatocellular carcinoma - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/16127248> ▾ [翻译此页](#)

作者: K Saito - 2005 - 被引用次数: 163 - [相关文章](#)

Magn Reson Med Sci. 2005;4(1):1-9. **Gd-EOB-DTPA enhanced MRI for hepatocellular carcinoma**: quantitative evaluation of tumor enhancement in **hepatobiliary phase**. Saito K(1), Kotake F, Ito N, Ozuki T, Mikami R, Abe K, Shimazaki Y. Author information: (1)Department of Radiology, Tokyo Medical University Kasumigaura ...

Focal hepatic lesions in Gd-EOB-DTPA enhanced MRI: the atlas

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3443279/> ▾ [翻译此页](#)

作者: JT Campos - 2012 - 被引用次数: 47 - [相关文章](#)

2012年6月15日 - Background. **Gd-EOB-DTPA** is a contrast agent with combined properties of a