

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 37870

Manuscript Type: ORIGINAL ARTICLE

Case Control Study

Serum interleukin-34 level can be an indicator of liver fibrosis in patients with chronic hepatitis B virus infection

Wang YQ *et al.* IL-34 and liver fibrosis in CHB patients

Yin-Qiu Wang, Wen-Jun Cao, Yu-Feng Gao, Jun Ye, Gui-Zhou Zou

Abstract

AIM

To investigate whether serum interleukin (IL)-34 levels are correlated with hepatic inflammation and fibrosis in patients with chronic hepatitis B virus

Match Overview

1	Crossref 178 words Dultz, G., L. Gerber, H. Farnik, A. Berger, J. Vermehren, T. Pileri, S. Zeuzem, A. Piiper, B. Kronenberger, and O. Waidman	5%
2	Crossref 97 words Huang, Haijun, Tiangang Wu, Jian Mao, Yongxing Fang, Jiajie Zhang, Lihua Wu, Shu Zheng, Biao Yang Lin, and Hong	3%
3	Internet 30 words crawled on 24-Nov-2017 www.wjgnet.com	1%
4	Crossref 19 words Maieron, A., P. Salzl, M. Peck-Radosavljevic, M. Trauner, S. Hametner, R. Schöfl, P. Ferenci, and M. Ferlitsch. "Von Wille	1%
5	Crossref 16 words Preisser, Laurence, Charline Miot, Hélène Le Guillou-Guillemette, Elodie Beaumont, Etienne D Foucher, Erwan Garo,	<1%

找到约 152,000 条结果 (用时 0.72 秒)

Interleukin-34 as a fibroblast-derived marker of liver fibrosis in patients ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4929441/> ▼ [翻译此页](#)

作者: H Shoji - 2016 - 被引用次数: 6 - [相关文章](#)

2016年7月1日 - Because we hypothesized that macrophage-related factors could serve as biomarkers of fibrosis, we examined serum levels of IL-34, M-CSF, and ... show that serum IL-34 can serve as a feasible diagnostic marker of fibrosis not only in patients with NAFLD but also in those with chronic HCV infection.

Association of Serum Autotaxin Levels with Liver Fibrosis in Patients ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5397977/> - [翻译此页](#)

作者: T Yamazaki - 2017 - 被引用次数: 3 - [相关文章](#)

2017年4月20日 - Association of Serum Autotaxin Levels with Liver Fibrosis in Patients with Chronic Hepatitis C. Tomoo Yamazaki,^{1,*} Satoru Joshita,^{1,*} ... We sought to clarify the diagnostic ability of ATX for liver fibrosis in 593 biopsy-confirmed hepatitis C virus (HCV)-infected patients. The diagnostic accuracy of ATX was ...

Serum Cytokeratin 18 M30 Levels in Chronic Hepatitis B Reflect Both ...

<https://www.hindawi.com/journals/mi/2017/3480234/> ▼ [翻译此页](#)

作者: M Świdarska - 2017

2017年7月9日 - We aimed to investigate an association between serum M30 CK-18 and histological activity and phase of HBV infection. 150 Caucasian ... It is important to note that frequent exacerbations of chronic hepatitis B (CHB) results in the progression of liver fibrosis which eventually leads to liver cirrhosis [2].

Rather than Rs1800796 Polymorphism, Expression of Interleukin-6 Is ...

<https://www.hindawi.com/journals/dm/2013/508023/> ▼ [翻译此页](#)

作者: S Tang - 2013 - 被引用次数: 8 - [相关文章](#)

2013年11月4日 - Rs1800796 polymorphism was genotyped in 641 Chinese Han patients with chronic HBV infection, including 23 IT, 25 IC, 292 CHB, 153 LC, and 148 ... the progression of chronic HBV



Serum interleukin-34 level can be an indicator of liver fibrosis in patients with ct



登录

全部

图片

新闻

视频

购物

更多

设置

工具

找到约 111,000 条结果 (用时 0.53 秒)

Interleukin-34 as a fibroblast-derived marker of liver fibrosis in patients ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4929441/> - 翻译此页

作者 : H Shoji - 2016 - 被引用次数 : 6 - 相关文章

2016年7月1日 - Preisser et al. recently reported that IL-34 and M-CSF increased in the sera of hepatitis C virus (HCV) infected patients with advanced liver fibrosis ... levels increased in association with the fibrosis stage in patients with chronic hepatitis C, suggesting that sCD163 can be useful as a marker of liver fibrosis.

Clinical predictors of liver fibrosis in patients with chronic hepatitis B ...

<https://www.ncbi.nlm.nih.gov/pubmed/29390144> - 翻译此页

作者 : JF Wu - 2018

2018年1月30日 - Clinical predictors of liver fibrosis in patients with chronic hepatitis B virus infection from children to adults. Wu JF(1), Song ... The AA genotype at rs16944 and the CC genotype at rs1143627 of the interleukin-1 β gene were associated with higher serum interleukin-1 β levels and liver fibrosis. Conclusions: ... 缺少字词 : 34 indicator

Predictors for advanced liver fibrosis in chronic hepatitis B virus ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5740558/> - 翻译此页

作者 : D Wang - 2017 - 相关文章

2017年9月29日 - The aim of the present study was to evaluate the predictors for advanced liver fibrosis in patients with chronic hepatitis B virus (HBV) infection with persistently normal alanine aminotransferase ... This division is based on HBV DNA viral load, HBeAg and serum levels of alanine aminotransferase (ALT) (4).

The Effect of Hepatitis B Virus Infection on Hepcidin Expression in ...

www.annclinlabsci.org/content/43/2/126.full - 翻译此页

作者 : X Wang - 2013 - 被引用次数 : 10 - 相关文章

Viral infections that disrupt liver function can be accompanied by changes in iron homeostasis, and iron

全部

图片

新闻

视频

购物

更多

设置

工具

找到约 453,000 条结果 (用时 0.48 秒)

The effect of hepatitis B virus infection on hepcidin expression in ...

<https://www.ncbi.nlm.nih.gov/pubmed/23694786> - 翻译此页

作者: XH Wang - 2013 - 被引用次数: 10 - 相关文章

Abnormal hepcidin expression has been reported following liver injury, including **liver cirrhosis**, alcoholic liver disease, and **chronic hepatitis B** and hepatitis C. However ... METHODS: We investigated **serum hepcidin levels** in 71 **patients** with different **HBV** DNA loads, 10 **patients** with hepatocellular carcinoma (HCC), and 13 ...

Interleukin-34 as a fibroblast-derived marker of liver fibrosis in patients ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4929441/> - 翻译此页

作者: H Shoji - 2016 - 被引用次数: 6 - 相关文章

2016年7月1日 - Preisser et al. recently reported that IL-34 and M-CSF increased in the sera of **hepatitis C virus** (HCV) **infected** patients with advanced **liver fibrosis** ... **levels** increased in association with the fibrosis stage in **patients with chronic hepatitis C**, suggesting that sCD163 **can** be useful as a marker of **liver fibrosis**.

Predictors for advanced liver fibrosis in chronic hepatitis B virus ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5740558/> - 翻译此页

作者: D Wang - 2017 - 相关文章

2017年9月29日 - The aim of the present study was to evaluate the predictors for advanced **liver fibrosis** in **patients with chronic hepatitis B virus** (HBV) **infection** with persistently normal alanine aminotransferase ... This division is based on **HBV** DNA **viral** load, HBeAg and **serum levels** of alanine aminotransferase (ALT) (4).

Zakim and Boyer's Hepatology: A Textbook of Liver Disease E-Book

<https://books.google.com/books?isbn=0323446574> - 翻译此页

Thomas D. Boyer, Arun J. Sanyal, Norah A Terrault - 2016 - Medical

Bedossa P: Intraobserver and interobserver variations in liver biopsy interpretation in **patients with chronic hepatitis C**. Hepatology 20:15-20, 1994. 7. Ratzliff V, et al: ... Hepatol 22:696-699, 1995.

Fontana RJ, et al: Relationship of **serum** fibrosis markers with **liver fibrosis** stage and collagen content in patients with advanced ...