

## Match Overview

- | Match | Source  | Words | Similarity |
|-------|---|-------|------------|
| 1     | Crossref 24 words<br>Dae Won Ma, Young Hoon Youn, Jae Jun Park, Jie-Hyun Kim, Hyojin Park. "Sa1092 Risk Factors of Electrocoagulation Syndrome after Endoscopic Submucosal Dissection of the Esophagus" | 24    | 1%         |
| 2     | Crossref 15 words<br>Manabu Takeuchi. "Strategy for endoscopic submucosal dissection of carcinoma of the abdominal esophagus", Esophagus  | 15    | 1%         |

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 37955

Manuscript Type: ORIGINAL ARTICLE

### Retrospective Study

**Risk factors of electrocoagulation syndrome after esophageal endoscopic submucosal dissection**

Dae Won Ma, Young Hoon Youn, Da Hyun Jung, Jae Jun Park, Jie-Hyun Kim, Hyojin Park

### Abstract

#### AIM

To investigate post endoscopic submucosal dissection electrocoagulation syndrome (PEECS) of the esophagus.

#### METHODS

We analyzed 55 consecutive cases with esophageal endoscopic submucosal

找到约 3,000 条结果 (用时 0.28 秒)

### Sa1092 Risk Factors of Electrocoagulation Syndrome After ...

[www.giejournal.org/article/S0016-5107\(17\)30600-4/abstract](http://www.giejournal.org/article/S0016-5107(17)30600-4/abstract) - 翻译此页

作者 : DW Ma - 2017

Some patients experience fever, chest pain and systemic inflammatory response after esophageal endoscopic submucosal dissection (ESD), despite of no obvious perforation. However, the risk factors of such post ESD electrocoagulation syndrome (PEECS) are not well known. This study aimed to evaluate the risk factors ...

### Su1612 Risk Factors of Electrocoagulation Syndrome After ...

[www.giejournal.org/article/S0016-5107\(16\)01189-5/fulltext](http://www.giejournal.org/article/S0016-5107(16)01189-5/fulltext) - 翻译此页

作者 : S Ito - 2016

Colorectal endoscopic submucosal dissection (ESD) is applied for the treatment of large colorectal superficial neoplasms; however, it is technically difficult, with perforation and delayed bleeding being well-known complications. However, there have been no large studies on electrocoagulation syndrome developing after ...

### Risk of electrocoagulation syndrome after endoscopic submucosal ...

<https://www.ncbi.nlm.nih.gov/pubmed/23990482> - 翻译此页

作者 : D Jung - 2013 - 被引用次数 : 21 - 相关文章

2013年8月29日 - BACKGROUND AND STUDY AIMS: The risk of post endoscopic submucosal dissection electrocoagulation syndrome (PEECS) is unknown. We aimed to investigate the incidence and clinicopathologic risk factors associated with PEECS after colorectal endoscopic submucosal dissection (ESD). PATIENTS ...

### Features of electrocoagulation syndrome after endoscopic ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/26202127> - 翻译此页

作者 : T Yamashina - 2016 - 被引用次数 : 14 - 相关文章

Features of electrocoagulation syndrome after endoscopic submucosal dissection for colorectal neoplasm. Yamashina T(1)(2), Takeuchi Y(1), Uedo N(1), ... The aim of this study was to evaluate the



[全部](#)[图片](#)[视频](#)[新闻](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 2,870 条结果 (用时 0.43 秒)

### Sa1092 Risk Factors of Electrocoagulation Syndrome After ...

[www.giejournal.org/article/S0016-5107\(17\)30600-4/abstract](http://www.giejournal.org/article/S0016-5107(17)30600-4/abstract) - 翻译此页

作者 : DW Ma - 2017

Some patients experience fever, chest pain and systemic inflammatory response after esophageal endoscopic submucosal dissection (ESD), despite of no obvious perforation. However, the risk factors of such post ESD electrocoagulation syndrome (PEECS) are not well known. This study aimed to evaluate the risk factors ...

### Risk Factors of Electrocoagulation Syndrome After Esophageal ...

[www.giejournal.org/article/S0016-5107\(17\)30600-4/pdf](http://www.giejournal.org/article/S0016-5107(17)30600-4/pdf) - 翻译此页

作者 : DW Ma - 2017

Sa1092. Risk Factors of Electrocoagulation Syndrome After. Esophageal Endoscopic Submucosal Dissection. Dae Won Ma\*, Young Hoon Youn, Jae Jun Park, Jie-Hyun Kim, Hyojin Park. Department of Internal Medicine, Gangnam Severance Hospital, Yonsei. University College of Medicine, Seoul, Korea (the Republic of).

### Risk of electrocoagulation syndrome after endoscopic submucosal ...

<https://www.ncbi.nlm.nih.gov/pubmed/23990482> - 翻译此页

作者 : D Jung - 2013 - 被引用次数 : 21 - 相关文章

2013年8月29日 - BACKGROUND AND STUDY AIMS: The risk of post endoscopic submucosal dissection electrocoagulation syndrome (PEECS) is unknown. We aimed to investigate the incidence and clinicopathologic risk factors associated with PEECS after colorectal endoscopic submucosal dissection (ESD). PATIENTS ...

### Features of electrocoagulation syndrome after endoscopic ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/26202127> ▼ 翻译此页

作者 : T Yamashina - 2016 - 被引用次数 : 14 - 相关文章

Features of electrocoagulation syndrome after endoscopic submucosal dissection for colorectal neoplasm. Yamashina T(1)(2), Takeuchi Y(1), Uedo N(1), ... The aim of this study was to evaluate the

[全部](#)[图片](#)[视频](#)[新闻](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 5,750 条结果 (用时 0.45 秒)

## Google 学术: Risk factors of electrocoagulation syndrome after esophageal endoscopic submucosal dissection

... for complications of endoscopic submucosal dissection ... - Mannen - 被引用次数: 149

Complications of colonoscopy - Fisher - 被引用次数: 179

... endoscopic submucosal dissection: is it suitable in ... - Uraoka - 被引用次数: 61

## Sa1092 Risk Factors of Electrocoagulation Syndrome After ...

[www.giejournal.org/article/S0016-5107\(17\)30600-4/abstract](http://www.giejournal.org/article/S0016-5107(17)30600-4/abstract) - 翻译此页

作者: DW Ma - 2017

Some patients experience fever, chest pain and systemic inflammatory response after esophageal endoscopic submucosal dissection (ESD), despite of no obvious perforation. However, the risk factors of such post ESD electrocoagulation syndrome (PEECS) are not well known. This study aimed to evaluate the risk factors ...

## Su1612 Risk Factors of Electrocoagulation Syndrome After ...

[www.giejournal.org/article/S0016-5107\(16\)01189-5/fulltext](http://www.giejournal.org/article/S0016-5107(16)01189-5/fulltext) - 翻译此页

作者: S Ito - 2016

Colorectal endoscopic submucosal dissection (ESD) is applied for the treatment of large colorectal superficial neoplasms; however, it is technically difficult, with perforation and delayed bleeding being well-known complications. However, there have been no large studies on electrocoagulation syndrome developing after ...

## Risk of electrocoagulation syndrome after endoscopic submucosal ...

<https://www.ncbi.nlm.nih.gov/pubmed/23990482> - 翻译此页

作者: D Jung - 2013 - 被引用次数: 21 - 相关文章

2013年8月29日 - BACKGROUND AND STUDY AIMS: The risk of post endoscopic submucosal dissection electrocoagulation syndrome (PEECS) is unknown. We aimed to investigate the incidence and clinicopathologic risk factors associated with PEECS after colorectal endoscopic submucosal dissection (ESD). PATIENTS ...

Features of electrocoagulation syndrome after endoscopic ... NCBI