

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 38458

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Post-polypectomy bleeding and thromboembolism risks associated with warfarin *vs* direct oral anticoagulants

Naohiro Yanagisawa, Naoyoshi Nagata, Kazuhiro Watanabe, Tatsuhiko Iida, Mariko Hamada, Sakurako Kobayashi, Takuro Shimbo, Junichi Akiyama, Naomi Uemura

Abstract

AIM

To verify the validity of the endoscopy guidelines for patients taking warfarin

Match Overview

- | Match Number | Source | Words | Similarity |
|--------------|--|----------|------------|
| 1 | Crossref | 31 words | 1% |
| | Toshiyuki Kubo, Kentaro Yamashita, Kei Onodera, Tomoya Iida, Yoshiaki Arimura, Masanori Nojima, Hiroshi Nakase ... | | |
| 2 | Internet | 19 words | <1% |
| | crawled on 20-Sep-2017
journals.plos.org | | |



全部

新闻

图片

视频

购物

更多

设置

工具

找到约 17,200 条结果 (用时 0.48 秒)

Therapeutic endoscopy-related GI bleeding and thromboembolic ... - Gut

gut.bmj.com > Online First - 翻译此页

作者 : N Nagata - 2017 - 相关文章

Oral anticoagulant users have a higher **risk** of GI **bleeding after** therapeutic endoscopy, whereas temporal discontinuation of oral anticoagulants may increase their **risk** of **thromboembolism**. The **risk** of endoscopy-related GI **bleeding or thromboembolism** may differ between **direct oral anticoagulants** (DOACs) and **warfarin**.

Direct oral anticoagulants and digestive bleeding: therapeutic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5424873/> - 翻译此页

作者 : D Deutsch - 2017 - 被引用次数 : 3 - 相关文章

2017年4月17日 - Resuming **or** not resuming **anticoagulation** with a DOAC **after bleeding or** a risky procedure depends on the thrombotic and **bleeding risk** as well as the procedure **involved**. This discussion should always involve the cardiologist and decisions should be taken by a pluridisciplinary team. Keywords: apixaban ...

Endoscopy in patients on antiplatelet or anticoagulant therapy ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4789831/> - 翻译此页

作者 : AM Veitch - 2016 - 被引用次数 : 107 - 相关文章

Endoscopy in patients on antiplatelet **or** anticoagulant therapy, including **direct oral anticoagulants**: British Society of Gastroenterology (BSG) and European Society of Even when temporarily discontinued, **warfarin** therapy is **associated** with an increased **risk** of **post-polypectomy bleeding** (PPB) and patients should be ...

Gastrointestinal endoscopy in patients on anticoagulant therapy and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5198250/> - 翻译此页

作者 : A Zullo - 2017 - 被引用次数 : 2 - 相关文章

2016年10月7日 - **Oral anticoagulants** need to be temporarily withheld, and consideration must be given to whether a periendoscopic "bridge" therapy, typically a low-molecular-weight heparin, should be used



Post-polypectomy bleeding and thromboembolism risks associated with w



全部

图片

新闻

视频

购物

更多

设置

工具

找到约 77,800 条结果 (用时 0.45 秒)

Therapeutic endoscopy-related GI bleeding and thromboembolic ... - Gut

gut.bmj.com > Online First - 翻译此页

作者: N Nagata - 2017 - 相关文章

Oral anticoagulant users have a higher **risk of GI bleeding after** therapeutic endoscopy, whereas temporal discontinuation **of** oral anticoagulants may increase their **risk of thromboembolism**. The **risk of endoscopy-related GI bleeding or thromboembolism** may differ between **direct oral anticoagulants** (DOACs) and **warfarin**.

Risk of Endoscopy-Related GI Bleeding and Thromboembolism ...

www.practiceupdate.com/.../risk-of...related...bleeding-and-thromboembolis... - 翻译此页

2017年10月4日 - **Risk of Endoscopy-Related GI Bleeding and Thromboembolism Associated With Warfarin or Direct Oral Anticoagulants**. ... retrospective cohort study compared **post-procedure bleeding** outcomes **in** patients on **warfarin versus** direct oral anti-coagulants (DOACs) undergoing high-risk endoscopic procedures.

Direct oral anticoagulants and digestive bleeding: therapeutic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5424873/> - 翻译此页

作者: D Deutsch - 2017 - 被引用次数: 3 - 相关文章

2017年4月17日 - The use **of direct oral anticoagulants** (DOACs) was an important step forward **in** the management **of** atrial fibrillation and venous **thromboembolism** (VTE). ... Resuming **or** not resuming anticoagulation with a DOAC **after bleeding or** a risky procedure depends on the thrombotic and



Post-polypectomy bleeding and thromboembolism risks associated with



全部

图片

新闻

视频

购物

更多

设置

工具

找到约 9,550 条结果 (用时 0.45 秒)

Therapeutic endoscopy-related GI bleeding and thromboembolic ... - Gut

gut.bmj.com › Online First - 翻译此页

作者: N Nagata - 2017 - 相关文章

Oral anticoagulant users have a higher **risk of GI bleeding after** therapeutic endoscopy, whereas temporal discontinuation **of** oral anticoagulants may increase their **risk of thromboembolism**. The **risk of endoscopy-related GI bleeding or thromboembolism** may differ between **direct oral anticoagulants** (DOACs) and **warfarin**.

Management of Anticoagulants and Antiplatelet Agents During ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/28344132> - 翻译此页

作者: LA Feagins - 2017 - 被引用次数: 2 - 相关文章

2017年3月24日 - The **risks of** precipitating **thromboembolic** complications if anticoagulants are stopped must be weighed against the **risk of postpolypectomy bleeding** if these ... For the thienopyridines, **warfarin**, and the **direct oral anticoagulants**, the decision to interrupt **or** continue these agents for endoscopy will involve ...

Direct oral anticoagulants and digestive bleeding: therapeutic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5424873/> - 翻译此页

作者: D Deutsch - 2017 - 被引用次数: 3 - 相关文章

2017年4月17日 - The use **of direct oral anticoagulants** (DOACs) was an important step forward **in** the management **of** atrial fibrillation and venous **thromboembolism** (VTE). ... Resuming **or** not resuming anticoagulation with a DOAC **after bleeding or** a risky procedure depends on the thrombotic and **bleeding risk** as well as ...

Endoscopy in patients on antiplatelet or anticoagulant therapy ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4789831/> - 翻译此页

作者: AM Veitch - 2016 - 被引用次数: 118 - 相关文章

Guidelines for the management **of** patients on **warfarin or direct oral anticoagulants** (DOAC) undergoing endoscopic procedures. Even when temporarily discontinued, **warfarin** therapy is **associated** with an increased **risk of post-polypectomy bleeding** (PPB) and patients should be advised **of** this. For patients with ...