



时间不限

2018以来

2017以来

2014以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☒ 包括专利☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

[HTML] Preoperative opioid misuse is associated with increased morbidity and mortality after elective orthopaedic surgery

ME Menendez, D Ring, BT Bateman - Clinical Orthopaedics and Related ..., 2015 - Springer

... The effect size for many of the associations that we reported are large (> 2 ... multidisciplinary approaches to effectively reduce opioid prescribing and associated adverse consequences in the ... Discharge Survey and Nationwide Inpatient Sample: the databases used affect results in ...

☆ 99 被引用次数: 51 相关文章 所有 10 个版本

[HTML] A flood of opioids, a rising tide of deaths

S Okie - New England Journal of Medicine, 2010 - Mass Medical Soc

... substances registered with the Drug Enforcement Administration (DEA) and about 4 million US patients receiving long-acting or extended-release opioids each year, the FDA's opioid REMS will affect far more ... 2017. Opiate Action on Sleep and Breathing ... (2016) Effect of opioid ...

☆ 99 被引用次数: 636 相关文章 所有 12 个版本

[HTML] Opioid abuse in chronic pain—misconceptions and mitigation strategies

ND Volkow, AT McLellan - New England Journal of Medicine, 2016 - Mass Medical Soc

... When combined, these pharmacologic properties affect the rapidity of onset, potency, and duration of both the ... associated with the opioid itself (type, dose, potency, and duration of action) and those ... or concurrent prescription of any medication that has a depressive effect on the ...

☆ 99 被引用次数: 196 相关文章 所有 7 个版本

Neuropsychological consequences of opiate use

SA Gruber, MM Silveri, DA Yurgelun-Todd - Neuropsychology review, 2007 - Springer

... buprenorphine and the buprenorphine/naloxone (B/N) combination produce long durations of action, which has ... Treatment with B/N did not affect overall performance on any of the cognitive



## Match Overview



There are no matching sources for this report.

**Name of Journal:** *World Journal of Orthopedics*

**Manuscript NO:** 38639

**Manuscript Type:** Original Article

*Retrospective Cohort Study*

**Effect of opioid dependence or abuse on opioid utilization after shoulder arthroplasty**

**Derek D Berglund, Samuel Rosas, Jennifer Kurowicki, Dragomir Mijic, Jonathan C Levy**

**Abstract**

*AIM*

To examine whether opioid dependence or abuse has an effect on opioid utilization after anatomic or reverse total shoulder arthroplasty.

找到约 211,000 条结果 (用时 0.30 秒)

## Google 学术: Effect of opioid dependence or abuse on opioid utilization after shoulder arthroplasty

Preoperative opioid misuse is associated with ... - Menendez - 被引用次数: 60

A flood of opioids, a rising tide of deaths - Okie - 被引用次数: 668

Opioid abuse in chronic pain—misconceptions and ... - Volkow - 被引用次数: 239

### [PDF] Preoperative opioid use associated with worse outcomes after ...

<https://sa1s3.patientpop.com/assets/docs/26094.pdf> ▼ 翻译此页

作者: BJ Morris - 2016 - 被引用次数: 15 - 相关文章

Keywords: Anatomic total shoulder arthroplasty; opioid use; glenohumeral joint osteoarthritis. Primary ... to potential abuse and misuse. ... the impact of preoperative opioid use on outcomes and patient satisfac- tion after TSA. We hypothesized that patients reporting preoperative opioid use would demonstrate lower preoper-

### Preoperative Opioid Misuse is Associated With Increased Morbidity ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4457771/> ▼ 翻译此页

作者: ME Menendez - 2015 - 被引用次数: 57 - 相关文章

2015年2月19日 - To characterize the association of opioid abuse and dependence with in-hospital mortality, morbidity, and resource use after orthopaedic surgery, we identified patients with a diagnosis of opioid-type dependence (ICD-9-CM codes 304.0x, 304.7x) or nondependent opioid abuse (ICD-9-CM code 305.5x).

### Preoperative Opioid Misuse is Associated With ... - Springer Link

<https://link.springer.com/content/pdf/10.1007/s11999-015-4173-5.pdf> - 翻译此页

作者: ME Menendez - 2015 - 被引用次数: 60 - 相关文章



找到约 282,000 条结果 (用时 0.44 秒)

## Google 学术 : Effect of opioid dependence or abuse on opioid utilization after shoulder arthroplasty

Preoperative opioid misuse is associated with ... - Menendez - 被引用次数 : 61

A flood of opioids, a rising tide of deaths - Okie - 被引用次数 : 672

Opioid abuse in chronic pain—misconceptions and ... - Volkow - 被引用次数 : 249

### Preoperative Opioid Misuse is Associated With Increased Morbidity ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4457771/> ▼ 翻译此页

作者 : ME Menendez - 2015 - 被引用次数 : 61 - 相关文章

2015年2月19日 - The prevalence of opioid abuse and dependence increased from 0.095% in ... surgery: primary THA (81.51), TKA (81.54), total shoulder arthroplasty (81.80, ... morbidity, and resource use after orthopaedic surgery, we identified patients ..... Yet little is known regarding the effect of this trend on perioperative ...

### [PDF] Preoperative opioid use associated with worse outcomes after ...

<https://sa1s3.patientpop.com/assets/docs/26094.pdf> ▼ 翻译此页

作者 : BJ Morris - 2016 - 被引用次数 : 15 - 相关文章

Keywords: Anatomic total shoulder arthroplasty; opioid use; glenohumeral joint ... associated pain.2,20  
The impact of opioid use is not limited to potential abuse and misuse. .... potentially to establish a dose-dependent impact of preop-

### Preoperative Opioid Misuse is Associated With ... - Springer Link

<https://link.springer.com/content/pdf/10.1007/s11999-015-4173-5.pdf> - 翻译此页

作者 : ME Menendez - 2015 - 被引用次数 : 61 - 相关文章

2015年2月19日 - Morbidity and Mortality After Elective Orthopaedic Surgery. Mariano E. Menendez ...  
lence and effect of high-risk opioid use (eg, abuse, dependence) in the ... prevalence of opioid abuse and dependence in patients undergoing major .... THA (81.51), TKA (81.54), total shoulder arthroplasty. (81.80, 81.88) ...

### Trends and risk factors for prolonged opioid use after ...

<https://online.boneandjoint.org.uk/.../0301-620X.100B1.BJJ-2017-0547.R1> - 翻译此页

作者 : NA Bedard - 2018 - 相关文章

opioid use after UKA, total knee arthroplasty (TKA) and total hip arthroplasty (THA). ... as anxiety/depression, smoking, back pain and substance abuse were found to significantly ... dependence, resulted in significantly prolonged opioid use after ... Patient risk factors were analysed for