

Name of journal: *World Journal of Gastroenterology*

Manuscript No: 38690

Manuscript Type: ORIGINAL ARTICLE

Retrospective study

Multikinase inhibitor-associated hand-foot skin reaction ³ as a predictor of outcomes in patients with hepatocellular carcinoma treated with sorafenib

Masanori Ochi, Toshiro Kamoshida, Atsushi Ohkawara, Haruka Ohkawara,
Nobushige Kakinoki, Shinji Hirai, Akinori Yanaka

Abstract

AIM

To investigate the relationship between the onsets of multikinase inhibitor (MKI)-associated hand-foot skin reaction (HFSR) and prognosis under

Match Overview

1	Internet 92 words crawled on 02-Mar-2017 theoncologist.alphamedpress.org	3%
2	Crossref 38 words "Abstracts", <i>Hepatology</i> , 10/2011	1%
3	Crossref 34 words Seok Hyun Son, Eun Young Park, Hee Hyun Park, Chul Seung Kay, Hong Seok Jang. "Pre-radiotherapy neutrophil-to-ly	1%
4	Internet 20 words crawled on 21-Sep-2017 www.wjgnet.com	1%
5	Internet 18 words crawled on 25-Dec-2016 office.wjgnet.com	1%
6	Crossref 11 words "The 6th Asia-Pacific Primary Liver Cancer Expert Meeting (APPLE 2015), Evidence and Consensus on HCC Manag...	<1%
7	Crossref 11 words Wörns, Marcus Alexander, and Peter Robert Galle. "Sorafenib for the treatment of hepatocellular carcinoma", <i>Hepatic O</i>	<1%



文章

获得 10 条结果 (用时0.09秒)

时间不限

2018以来

2017以来

2014以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

 包括专利 包含引用 创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Hand-foot skin reaction in patients treated with sorafenib: a clinicopathological study of cutaneous manifestations due to multitargeted kinase inhibitor therapy

CH Yang, WC Lin, CK Chuang... - British Journal of ..., 2008 - Wiley Online Library

... are warranted to elucidate the mechanisms of this novel multitargeted kinase inhibitor-associated skin reaction ... of the cutaneous side effects of sorafenib, a novel multi-kinase inhibitor ... Effect of Urea-Based Cream on Sorafenib-Associated Hand-Foot Skin Reactions in Patients ...

☆ 99 被引用次数: 152 相关文章 所有 7 个版本

[HTML] Sorafenib in advanced clear-cell renal-cell carcinoma

B Escudier, T Eisen, WM Stadler... - ... England Journal of ..., 2007 - Mass Medical Soc

... were predominantly of grade 1 or 2. The most common events were diarrhea, rash, fatigue, hand-foot skin reactions, alopecia, and ... AG-013736.21 Reported toxic effects of these agents include diarrhea, hypertension, fatigue, hand-foot skin reaction, and rash, although the ...

☆ 99 被引用次数: 4776 相关文章 所有 17 个版本

[HTML] Sorafenib-associated facial acneiform eruption

PR Cohen - Dermatology and therapy, 2015 - Springer

... Sorafenib is an oral multikinase inhibitor associated with the potential development of several ... Leukocytoclastic vasculitis masquerading as hand-foot syndrome in a patient treated with sorafenib ... K. Acne inversa-like lesions associated with the multi-kinase inhibitor sorafenib ...

☆ 99 被引用次数: 9 相关文章 所有 14 个版本

Understanding toxicities of targeted agents: implications for anti-tumor activity and management

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 16,000 条结果 (用时 0.51 秒)

Hand-foot skin reaction in patients treated with sorafenib: a ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/18070211> ▾ [翻译此页](#)

作者: CH Yang - 2008 - 被引用次数: 156 - [相关文章](#)

2007年12月6日 - Hand-foot skin reaction in patients treated with sorafenib: a ... Histopathological studies of skin toxicity associated with kinase inhibitors are currently unavailable. ... hand-foot skin reaction produced by the multikinase inhibitor sorafenib. ... and one with hepatocellular carcinoma (HCC)-received continuous, ...

Regorafenib-associated hand-foot skin reaction: practical advice on ...

<https://academic.oup.com/annonc/article/26/10/2017/143835> ▾ [翻译此页](#)

作者: B McLellan - 2015 - 被引用次数: 27 - [相关文章](#)

Regorafenib is an orally available, small-molecule multikinase inhibitor with international ... Management approaches for hand-foot skin reaction (HFSR) associated with licensed kinase inhibitors. Reference, Kinase inhibitor, Intervention, Outcome in advanced hepatocellular carcinoma patients treated with sorafenib.

Hand foot skin reaction in cancer patients treated with the multikinase ...

<https://academic.oup.com/annonc/article/19/11/1955/199487> ▾ [翻译此页](#)

作者: ME Lacouture - 2008 - 被引用次数: 199 - [相关文章](#)

2008年6月10日 - Hand foot skin reaction in cancer patients treated with the multikinase inhibitors sorafenib and sunitinib Sorafenib and sunitinib are two novel, small molecule multikinase inhibitors ... The associated cancer was renal cell carcinoma in eight cases (66%) and hepatocellular carcinoma for two cases (17%).

Hand-foot skin reaction (HFSR) and overall survival (OS) in the phase ...

ascopubs.org/doi/abs/10.1200/JCO.2018.36.4_suppl.412 - [翻译此页](#)

作者: J Bruix - 2018 - 被引用次数: 2

... for treatment of hepatocellular carcinoma (HCC) progressing on sorafenib. Background: Skin toxicity is a known adverse effect of multikinase inhibitors, and ... to be a predictor of OS in patients (pts) with HCC treated with sorafenib (Reig M, ... analysis explored whether HFSR with regorafenib was associated with OS in ...

Hand-foot skin reaction is a beneficial indicator of sorafenib therapy for ...

<https://www.tandfonline.com/doi/full/10.1080/17474124.2017.1373018> - [翻译此页](#)

作者: P Wang - 2018 - 被引用次数: 2 - [相关文章](#)

2017年9月3日 - KEYWORDS: Sorafenib, hepatocellular carcinoma, hand-foot skin reaction, whereas

找到约 14,700 条结果 (用时 0.49 秒)

Hand-foot skin reaction in patients treated with sorafenib: a ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/18070211> ▾ 翻译此页

作者: CH Yang - 2008 - 被引用次数: 156 - 相关文章

2007年12月6日 - **Hand-foot skin reaction** in patients treated with sorafenib: a ... Histopathological studies of skin toxicity associated with kinase inhibitors are currently unavailable. ... **hand-foot skin reaction** produced by the **multikinase inhibitor sorafenib**. ... and one with **hepatocellular carcinoma (HCC)**-received continuous, ...

Regorafenib-associated hand-foot skin reaction: practical advice on ...

<https://academic.oup.com/annonc/article/26/10/2017/143835> ▾ 翻译此页

作者: B McLellan - 2015 - 被引用次数: 27 - 相关文章

Regorafenib is an orally available, small-molecule **multikinase inhibitor** with ... a consistent and predictable adverse-event profile, with **hand-foot skin reaction (HFSR)** al. have developed and validated a **prediction** index for **sorafenib-related HFSR**, in advanced **hepatocellular carcinoma patients treated with sorafenib**.

Hand-Foot Skin Reaction is Associated with the Clinical Outcome in ...

<https://academic.oup.com/jjco/article/43/10/1023/851399> - 翻译此页

作者: K Nakano - 2013 - 被引用次数: 31 - 相关文章

2013年8月15日 - A **sorafenib-induced Hand-Foot skin reaction** in metastatic renal cell carcinoma ... to be associated with the clinical **outcome** in patients with mRCC treated with this drug (2). This probably occurred because the **inhibition** of the vascular ... **hepatocellular carcinoma (HCC)** (9) treated with **sorafenib**, the time to ...

Hand-foot skin reaction (HFSR) and overall survival (OS) in the phase ...

ascopubs.org/doi/abs/10.1200/JCO.2018.36.4_suppl.412 - 翻译此页

作者: J Bruix - 2018 - 被引用次数: 2

Hand-foot skin reaction (HFSR) and overall survival (OS) in the phase 3 ... for treatment of **hepatocellular carcinoma (HCC)** progressing on **sorafenib**. of **multikinase inhibitors**, and was shown to be a **predictor** of OS in patients (pts) ... analysis explored whether HFSR with regorafenib was associated with OS in RESORCE.