

ANSWERING REVIEWERS



Dear Editor,

Please find enclosed the edited manuscript in Word format (file name: 2429-review.doc).

Title: Gastric endoscopic submucosal dissection: From animal model to patient

Author: Nicolás González, Adolfo Parra-Blanco, Miguel Villa-Gómez, Alejandra Gamba, Andrés Taullard, Anaulina Silveira, Alberto Sanguinetti, Carolina Olano, Henry Cohen

Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 3877

The manuscript has been improved according to the suggestions of reviewers:

1 Format has been updated

2 Revision has been made according to the suggestions of the reviewer

(1) 1) As mentioned in the present study, a sequential ESD training program would be important as the education for endoscopists. However, for ESD treatment of gastric cancer, it should be more important to give them the training of endoscopic diagnosis, especially to diagnose demarcation line between cancer tissue and non-cancerous area. Please discuss about this matter. 2) Usually in Japan, the pattern of early gastric cancer is popular as depressive type, so-called IIC. However, in the present study, the authors performed ESD for patients with Ila type early gastric cancers.

(2) Had describe conventional EGD, the experience of the colonoscope, the experience of the gastric cancer diagnosis in detail. The information about the standard deviation would be more informative. The simple t-test is not appropriate for this study. It is not clear how patients were selected for ESD treatment.

3 References and typesetting were corrected

Thank you again for publishing our manuscript in the *World Journal of Gastroenterology*.

Sincerely yours,