

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 38868

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

The Impact of ³the Number of Examined Lymph Nodes on Outcomes in
Patients with Lymph Node-Negative Gallbladder Carcinoma

Dong-Xu Fan, Rong-Wei Xu, Yong-Cheng Li, Bing-Qing Zhao, Ming-Yu
Sun

Abstract

AIM

Match Overview

- | | | |
|---|--|----|
| 1 | Crossref 26 words
Minia Hellan. "The Impact of Lymph Node Number on Survival in Patients With Lymph Node-Negative Pancreatic Cancer" | 1% |
| 2 | Internet 12 words
crawled on 02-Feb-2017
www.health.gov.au | 1% |
| 3 | Crossref 12 words
Chuangang Tang, Pei Wang, Xiaoxin Li, Bingqing Zhao, Haichang Yang, Haifeng Yu, Changwen Li. "Lymph node status and survival in patients with gallbladder cancer: a meta-analysis" | 1% |

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 82,200 条结果 (用时 0.52 秒)

Impact of lymph node ratio as a valuable prognostic factor in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3594644/> - 翻译此页

作者: BG Choi - 2013 - 被引用次数: 11 - 相关文章

跳到RESULTS - Sixty six patients (67%) were node negative patients and positive LNs were found in 33 patients (33%) among the 99 patients who underwent LN ... Kaplan-Meier survival curves comparing patients in node negative disease with fewer than eight lymph nodes (LNs) examined to those with eight or ...

The impact of tumor extent (T stage) and lymph node involvement (N ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2597319/> - 翻译此页

作者: V Zaydfudim - 2008 - 被引用次数: 25 - 相关文章

跳到Results - From this group 4365 patients underwent surgical treatment for gallbladder adenocarcinoma and 4048 patients (93%) had sufficient staging information recorded in the SEER database and survived the ... Only 1298 patients (32%) of the full cohort group had pathologically examined lymph nodes.

Immunohistochemically Demonstrated Lymph Node ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1570614/> - 翻译此页

作者: E Sasaki - 2006 - 被引用次数: 52 - 相关文章

The purpose of this study was to investigate whether immunohistochemically detected lymph node micrometastasis has prognostic significance in patients with advanced gallbladder carcinoma (pT2-4 tumors). For this purpose, a large number of lymph nodes were examined immunohistochemically, and the impact of lymph



The impact of the number of examined lymph nodes on outcomes in pati



全部

图片

新闻

购物

地图

更多

设置

工具

找到约 257,000 条结果 (用时 0.55 秒)

Immunohistochemically Demonstrated Lymph Node ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1570614/> ▼ 翻译此页

作者 : E Sasaki - 2006 - 被引用次数 : 52 - 相关文章

A total of 1476 lymph nodes from 67 patients with gallbladder carcinoma (pN0, n = 40; ... Results: Lymph node micrometastases were detected ... Lymph node micrometastasis has a significant survival impact in patients with ... For this purpose, a large number of lymph nodes were examined immunohistochemically, and the ...

Immunohistochemically Demonstrated Lymph Node ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1522147/> ▼ 翻译此页

作者 : Y Tojima - 2003 - 被引用次数 : 62 - 相关文章

Lymph node micrometastasis has no survival impact in patients with otherwise node-negative hilar cholangiocarcinoma. ... For this purpose a large number of lymph nodes were examined immunohistochemically, ligament was effective for improving outcome in gallbladder cancer patients without overt nodal metastasis.

Radical lymph node dissection and assessment: Impact on ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/23964151> - 翻译此页

作者 : GJ Liu - 2013 - 被引用次数 : 18 - 相关文章

2013年8月21日 - Then the metastatic to examined lymph nodes ratio (LNR) was calculated. Disease-specific survival (DSS) and predictors of outcome were analyzed. ... When node-negative patients were divided according to TLNC, there was ... value of six nodes was identified for the number of TLNC that clearly stratified ...

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 257,000 条结果 (用时 0.45 秒)

Radical lymph node dissection and assessment: Impact on ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/23964151> - 翻译此页

作者: GJ Liu - 2013 - 被引用次数: 19 - 相关文章

2013年8月21日 - Then the metastatic to **examined lymph nodes** ratio (LNR) was calculated. Disease-specific survival (DSS) and predictors of **outcome** were **analyzed**. ... When **node-negative patients** were divided according to TLNC, there was ... value of six nodes was identified for the **number** of TLNC that clearly stratified ...

Immunohistochemically Demonstrated Lymph Node ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1570614/> - 翻译此页

作者: E Sasaki - 2006 - 被引用次数: 52 - 相关文章

A total of 1476 **lymph nodes** from 67 **patients** with **gallbladder carcinoma** (pN0, n = 40; ... **Results:** **Lymph node** micrometastases were detected ... **Lymph node** micrometastasis has a significant survival **impact** in **patients** with ... For this purpose, a large **number** of **lymph nodes** were **examined** immunohistochemically, and the ...

Immunohistochemically Demonstrated Lymph Node ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1522147/> - 翻译此页

作者: Y Tojima - 2003 - 被引用次数: 62 - 相关文章

Lymph node micrometastasis has no survival **impact** in **patients** with otherwise **node-negative** hilar cholangiocarcinoma. ... For this purpose a large **number** of **lymph nodes** were **examined** immunohistochemically, ... ligament was effective for improving **outcome** in **gallbladder cancer patients** without overt nodal metastasis.

Radical lymph node dissection and assessment: Impact on ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3746389/> - 翻译此页

作者: GJ Liu - 2013 - 被引用次数: 19 - 相关文章

跳到**RESULTS** - A total of 465 **lymph nodes** taken from the 78 studied **patients** were ... Topographical distribution of 465 **lymph nodes** evaluated in 78 **patients** with **gallbladder cancer** n (%) ... The postoperative DSS of **node-negative patients** was ... Based on these **results**, the **number** of TLNC was placed into two ...