

找到约 14,100 条结果 (用时 0.47 秒)

Chemotherapy for the conversion of unresectable colorectal cancer ...

[www.croh-online.com/article/S1040-8428\(10\)00198-8/pdf](http://www.croh-online.com/article/S1040-8428(10)00198-8/pdf) - 翻译此页

作者: DG Power - 2011 - 被引用次数: 33 - 相关文章

A sound physiologic rationale exists for the use of **hepatic** arterial infusion (HAI) **therapy**. Studies have shown that HAI with floxuridine combined with systemic chemotherapy increases response rates and liver resection rates in those patients **with initially unresectable** CLM. Toxicity from preoperative chemotherapy, biologic ...

Chemotherapy for the conversion of unresectable colorectal cancer ...

[www.croh-online.com/article/S1040-8428\(10\)00198-8/fulltext](http://www.croh-online.com/article/S1040-8428(10)00198-8/fulltext) - 翻译此页

作者: DG Power - 2011 - 被引用次数: 33 - 相关文章

2010年10月25日 - The logic for considering neoadjuvant chemotherapy is to treat extra-**hepatic** micro-metastatic disease, to make liver resection easier, to assess aggressive ... **Conversion therapy** (the topic of this review) is, by definition, given to those patients **with initially unresectable** or borderline resectable CLM with a ...

Conversion Chemotherapy for Technically Unresectable Colorectal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4902434/> - 翻译此页

作者: M Basso - 2016 - 被引用次数: 2 - 相关文章

2016年5月20日 - Moreover, the role of molecular targeted **therapies** in the **treatment of initially unresectable** liver metastases is still controversial. ... known extra-**hepatic** disease at the **time** of diagnosis, or other cancers (with the exception of nonmelanoma skin cancers and in situ cervical **cancer**) diagnosed within the past 5 ...

Does conversion therapy in colorectal cancer achieve... - ResearchGate

https://www.researchgate.net/.../Does_conversion_therapy_in_colorectal_cancer_achieve...

Prolonged survival of **initially unresectable hepatic** colorectal **cancer** patients treated with **hepatic** arterial infusion of oxaliplatin followed by radical **surgery** of metastases. Ann Surg 2010; 251:686. 3. ... Only FOLFOX or CAPOX can be used as a **conversion** and as a neoadjuvant **therapy** in the same **time**. It means that if the ...

Name of Journal: World Journal of Clinical Cases

Manuscript NO: 39010

Manuscript Type: Systematic Review

For initially unresectable hepatocellular carcinoma: Conversion therapy
and the suitable timing for subsequent salvage surgery

Ze-Feng Zhang, Yu-Jun Luo, Quan Lu, Shi-Xue Dai, Wei-hong Sha

Abstract

AIM

To review the conversion therapy for initially unresectable hepatocellular carcinoma (HCC) patients and the suitable timing for subsequent salvage surgery.

Match Overview

1	Crossref 65 words Wan Yee Lau. "Salvage Surgery Following Downstaging of Unresectable Hepatocellular Carcinoma—A Strateg ...	1%
2	Crossref 40 words S. Ogata. "Sequential arterial and portal vein embolizations before right hepatectomy in patients with cirrhosis a ...	1%
3	Crossref 39 words Venous Embolization of the Liver, 2011.	1%
4	Internet 36 words crawled on 30-Sep-2010 archsurg.ama-assn.org	1%
5	Internet 32 words crawled on 25-Aug-2016 mdanderson.influent.utsystem.edu	<1%
6	Crossref 31 words She, Wong Hoi. "Strategies to increase the resectability of hepatocellular carcinoma", World Journal of Hepatolo	<1%
7	Internet 27 words crawled on 24-Feb-2017 theoncologist.alphamedpress.org	<1%
8	Crossref 26 words Hirai, I. "Evaluation of preoperative portal embolization f	<1%

[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 44,900 条结果 (用时 0.42 秒)

Google 学术: For initially unresectable hepatocellular carcinoma: Conversion therapy and the suitable timing for subsequent salvage surgery

... in unresectable cholangiocarcinoma: initial experience ... - Burger - 被引用次数: 153

... for patients with unresectable hepatic metastases from ... - D'Angelica - 被引用次数: 67

Guidelines for surgical treatment of hepatoblastoma in ... - Czauderna - 被引用次数: 155

Salvage surgery following downstaging of unresectable hepatocellular ...

<https://www.ncbi.nlm.nih.gov/pubmed/17891443> - 翻译此页

作者: WY Lau - 2007 - 被引用次数: 77 - 相关文章

2007年9月22日 - Salvage surgery following downstaging of unresectable ... gives the best chance of a cure for patients with hepatocellular carcinoma (HCC). However, the resectability of HCC at the time of diagnosis is low (10-30%). ... to the initial palliative treatment to downstage HCC to allow salvage surgical resection.

缺少字词: conversion subsequent

Salvage Surgery Following Downstaging of Unresectable ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1356407/> - 翻译此页

作者: W Lau - 2004 - 被引用次数: 144 - 相关文章

Occasionally, there are good responses to treatment so that salvage surgery becomes ... Patients with unresectable hepatocellular carcinoma, from September 1993 to ... The reasons why the HCC was initially unresectable but became resectable were ... The median survival from the time of surgery was 71.5 months (95% ...

缺少字词: conversion subsequent

Conversion Chemotherapy for Technically Unresectable Colorectal ...

找到约 41,200 条结果 (用时 0.46 秒)

Google 学术 : For initially unresectable hepatocellular carcinoma: Conversion therapy and the suitable timing for subsequent salvage surgery

... in unresectable cholangiocarcinoma: initial experience ... - Burger - 被引用次数 : 153

... for patients with unresectable hepatic metastases from ... - D'Angelica - 被引用次数 : 68

Guidelines for surgical treatment of hepatoblastoma in ... - Czauderna - 被引用次数 : 156

Salvage surgery following downstaging of unresectable hepatocellular ...

<https://www.ncbi.nlm.nih.gov/pubmed/17891443> - 翻译此页

作者 : WY Lau - 2007 - 被引用次数 : 77 - 相关文章

2007年9月22日 - Salvage surgery following downstaging of unresectable ... gives the best chance of a cure for patients with hepatocellular carcinoma (HCC). However, the resectability of HCC at the time of diagnosis is low (10-30%). ... to the initial palliative treatment to downstage HCC to allow salvage surgical resection.

缺少字词 : conversion subsequent

Salvage Surgery Following Downstaging of Unresectable ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1356407/> - 翻译此页

作者 : W Lau - 2004 - 被引用次数 : 144 - 相关文章

Occasionally, there are good responses to treatment so that salvage surgery becomes ... Patients with unresectable hepatocellular carcinoma, from September 1993 to The reasons why the HCC was initially unresectable but became resectable were ... The median survival from the time of surgery was 71.5 months (95% ...

缺少字词 : conversion subsequent

Conversion Chemotherapy for Technically Unresectable Colorectal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4902434/> - 翻译此页

作者 : M Basso - 2016 - 被引用次数 : 4 - 相关文章

2016年5月20日 - Our data confirmed that the conversion of initially inoperable disease to ... affected treatment duration, survival, or the possibility of surgery at the time of diagnosis. ... treatment for metastatic disease, known extra-hepatic disease at the time of Rescue surgery for unresectable colorectal liver metastases ...

Conversion Chemotherapy Using Cetuximab plus FOLFIRI Followed ...

<https://academic.oup.com/jjco/article/41/10/1229/1181362> - 翻译此页

作者 : K Uehara - 2011 - 被引用次数 : 9 - 相关文章