

Beta-blockers and physical frailty in patients with end-stage liver disease

Selena Z. Kuo¹, Blanca Lizaola¹, Hilary Hayssen¹, Jennifer C. Lai¹

⁵
¹Division of Gastroenterology/Hepatology, Department of Medicine, University of California, San Francisco, San Francisco, CA

Corresponding Author:

Jennifer C. Lai, MD, MBA, Assistant Professor, Gastroenterology, University of California, San Francisco, 513 Parnassus Ave, UCSF Box 0538, San Francisco, CA 94143, jennifer.lai@ucsf.edu

Supported by the Paul B. Beeson Career Development Award in Aging

Match Overview

1	Crossref 47 words Mariya L. Samoylova, Kenneth E. Covinsky, Marta Haftek, Selena Kuo, John P. Roberts, Jennifer C. Lai. "Disability	2%
2	Internet 39 words crawled on 12-May-2018 www.wjgnet.com	2%
3	Internet 38 words crawled on 15-Mar-2016 peppercenter.org	2%
4	Crossref 38 words Selena Z. Kuo, Marta Haftek, Jennifer C. Lai. "Factors Associated with Medication Non-adherence in Patients with ...	2%
5	Crossref 25 words Jin Ge, John P. Roberts, Jennifer C. Lai. "Race/ethnicity ... s associated with ABO-nonidentical liver transplantation i	1%
6	Crossref 22 words Jennifer C. Lai, Kenneth E. Covinsky, Jennifer L. Dodge, W. John Boscardin, Dorry L. Segev, John P. Roberts, San	1%
7	Internet 19 words crawled on 21-Jun-2017 ghrnet.org	1%
8	Crossref 18 words "Abstracts from the 30th Annual Meeting of the Society of General Internal Medicine", Journal of General Internal	1%
9	Internet 18 words crawled on 05-Oct-2017 file.scrip.org	1%



All

News

Shopping

Images

Videos

More

Settings

Tools

About 419,000 results (0.56 seconds)

Scholarly articles for **Beta-blockers and physical frailty in patients with end-stage liver disease**

... on mortality risk in **patients** with mitral regurgitation ... - **Wu** - Cited by 542

... of **patients** with decompensated alcoholic **cirrhosis** - **Vlachogiannakos** - Cited by 99

... left ventricular assist device for **end-stage** heart **failure** - **Rose** - Cited by 3433

Interventions to improve physical function and prevent adverse events ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5806401/>

by HD Trivedi - 2018 - Related articles

Jan 10, 2018 - **Frailty** in turn increases the **patient's** risk of hospitalization, accidental 83% Child class B or C), Lai et al. found that declining **physical** function was ... Sarcopenia affects 65–90% of **patients with end-stage liver disease** [4,11] Indeed, a similar degree of HVPG change with non-selective **beta-blockers** is ...

You visited this page on 5/18/18.

Home Exercise Training Improves Exercise Capacity in Cirrhosis ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5758716/>

by C Kruger - 2018 - Cited by 1 - Related articles

Jan 8, 2018 - Even **patients** with well compensated, early stage cirrhosis, have an approximately 40% Abbreviations: MELD, Model for **End Stage Liver Disease**; TIPS, Thirdly, **beta-blocker** therapy was not an exclusion criterion, and a small but ... Data in other populations have associated e-health **physical** activity ...

Exercise and physical activity for patients with end-stage liver ... - aasld

<https://aasldpubs.onlinelibrary.wiley.com/doi/pdf/10.1002/lt.24958>

by A Duarte-Rojo - 2018 - Cited by 4 - Related articles

copenia, poor **physical** fitness/cardiopulmonary endurance (CPE), and **frailty** are all associated with increased mortality in. ESLD. Clinical ... and **end-stage liver disease** (ESLD) has become more All

全部

新闻

图片

视频

购物

更多

设置

工具

找到约 381,000 条结果 (用时 0.59 秒)

Google 学术: [Beta-blockers and physical frailty in patients with end-stage liver disease](#)

... of adult patients with ascites due to cirrhosis 2012 - Runyon - 被引用次数: 433

... on mortality risk in patients with mitral regurgitation ... - Wu - 被引用次数: 543

... left ventricular assist device for end-stage heart failure - Rose - 被引用次数: 3467

Exercise and physical activity for patients with end-stage liver disease ...

<https://www.ncbi.nlm.nih.gov/pubmed/29024353> - 翻译此页

作者: A Duarte-Rojo - 2017 - 被引用次数: 8 - 相关文章

Exercise and physical activity for patients with end-stage liver disease: ... Frailty is the end result of prolonged sarcopenia and physical deconditioning.

缺少字词: beta blockers

Interventions to improve physical function and prevent adverse events ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5806401/> - 翻译此页

作者: HD Trivedi - 2018 - 相关文章

2018年1月10日 - The management of frailty and its associated adverse effects is imperative 83% Child class B or C). Lai et al. found that declining physical function ... Sarcopenia affects 65–90% of patients with end-stage liver disease [4,11] Indeed, a similar degree of HVP change with non-selective beta-blockers is ...

Exercise and physical activity for patients with end - stage liver disease ...

<https://aasldpubs.onlinelibrary.wiley.com/doi/full/10.1002/lt.24958> - 翻译此页

作者: A Duarte-Rojo - 2018 - 被引用次数: 8 - 相关文章

2017年10月11日 - Acute - on - chronic liver disease and infectious complications are leading (B) in patients listed for LT comparing MELD - Na, the frailty index score, All patients received beta - blockers during the study in order to prevent ...

[PDF] The Role of Physical Activity and Nutrition in the Sarcopenia of Cirrh...

www.mdpi.com/2411-5142/1/1/118/pdf - 翻译此页

作者: EM Travato - 被引用次数: 7 - 相关文章

找到约 381,000 条结果 (用时 0.59 秒)

Google 学术: [Beta-blockers and physical frailty in patients with end-stage liver disease](#)

... of adult [patients](#) with ascites due to [cirrhosis](#) 2012 - [Runyon](#) - 被引用次数: 433

... on mortality risk in [patients](#) with mitral regurgitation ... - [Wu](#) - 被引用次数: 543

... left ventricular assist device for [end-stage](#) heart [failure](#) - [Rose](#) - 被引用次数: 3467

Exercise and physical activity for patients with end-stage liver disease ...

<https://www.ncbi.nlm.nih.gov/pubmed/29024353> - [翻译此页](#)

作者: A Duarte-Rojo - 2017 - 被引用次数: 8 - [相关文章](#)

Exercise and [physical](#) activity for [patients](#) with [end-stage liver disease](#): ... [Frailty](#) is the end result of prolonged sarcopenia and [physical](#) deconditioning.

缺少字词: [beta blockers](#)

Interventions to improve physical function and prevent adverse events ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5806401/> - [翻译此页](#)

作者: HD Trivedi - 2018 - [相关文章](#)

2018年1月10日 - The management of [frailty](#) and its associated adverse effects is imperative 83% Child class B or C). Lai et al. found that declining [physical](#) function ... Sarcopenia affects 65–90% of [patients](#) with [end-stage liver disease](#) [4,11] Indeed, a similar degree of HVPG change with non-selective [beta-blockers](#) is ...

Exercise and physical activity for patients with end - stage liver disease ...

<https://aasldpubs.onlinelibrary.wiley.com/doi/full/10.1002/lt.24958> - [翻译此页](#)

作者: A Duarte-Rojo - 2018 - 被引用次数: 8 - [相关文章](#)

2017年10月11日 - Acute - on - [chronic liver disease](#) and infectious complications are leading (B) in [patients](#) listed for LT comparing MELD - Na, the [frailty](#) index score, All [patients](#) received [beta - blockers](#) during the study in order to prevent ...

[PDF] The Role of Physical Activity and Nutrition in the Sarcopenia of Cirrh...

www.mdpi.com/2411-5142/1/1/118/pdf - [翻译此页](#)

作者: EM Travato - 被引用次数: 7 - [相关文章](#)