

Name of journal: *World Journal of Clinical Cases*

Manuscript NO: 40034

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Diagnostic value of elevated serum carbohydrate antigen 19-9 level in acute cholangitis secondary to choledocholithiasis

Mei Y *et al.* Diagnostic value of CA19-9 level in secondary acute cholangitis

Yong Mei, Li Chen, Ci-Jun Peng, Jun Wang, Peng-Fei Zeng, Guo-Xing Wang,

Match Overview

1	Internet 30 words crawled on 15-Nov-2017 hub.hku.hk	1%
2	Internet 25 words crawled on 28-Nov-2017 s-space.snu.ac.kr	1%
3	Crossref 24 words "Monday 29 August 2016", European Heart Journal, 2016	1%
4	Internet 14 words crawled on 29-Nov-2016 www.basic.medscimonit.com	1%
5	Crossref 12 words "Abstracts from the Third Chinese Congress on Gerontology and Health Industry", Journal of the American Geriatrics	<1%
6	Internet 11 words crawled on 14-Jun-2018 www.birpublications.org	<1%
7	Crossref 11 words Xiao-Dan Ju, Peng-Fei Li, Peng-Bin Zhang. "Value of serum CA19-9 levels in diagnosis of secondary acute cholangitis"	<1%
8	Internet 11 words crawled on 23-Jun-2017 hentheman.com	<1%

[全部](#)[图片](#)[新闻](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 4,050 条结果 (用时 0.56 秒)

[Elevated serum level of carbohydrate antigen 19-9 in benign biliary ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3992417/> ▼ [翻译此页](#)

作者: MS Lin - 2014 - 被引用次数: 22 - [相关文章](#)

2014年3月15日 - Although carbohydrate antigen (CA19-9) level is frequently upregulated in ... The latter, including inflammatory stricture **secondary to choledocholithiasis**, Mirizzi syndrome, The **diagnostic value** of CA19-9 might be improved by using a more ... **Cholangitis** may result in **high** serum levels of CA19-9 [22].

缺少字词: 499

[Highly elevated serum levels of CA 19-9 in choledocholithiasis: a case ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2740064/> ▼ [翻译此页](#)

作者: G Marcouizos - 2009 - 被引用次数: 21 - [相关文章](#)

2009年7月30日 - However, high levels of serum CA 19-9 is occasionally found in benign disease of ... stone with **acute cholangitis** presenting with extraordinarily **high** serum CA 19-9 levels, ... A CA 19-9 **value** of more than 1.000 U/ml usually indicates a The clinical utility of serum CA 19-9 in the **diagnosis**, prognosis and ...

[Benign Hydronephrosis and Elevated of Serum Levels of ... - NCBI - NIH](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4917065/> ▼ [翻译此页](#)

作者: B Filipovic - 2016 - 被引用次数: 4 - [相关文章](#)

2016年6月11日 - There is no doubt that CA 19-9 level is occasionally elevated in ... in the differential **diagnosis** of elevated concentrations of CA 19-9 marker. ... track such as pancreatitis, **cholangitis**, and **choledocholithiasis** [5-7]. Serum **carbohydrate antigen** 19-9 levels in patients with unilateral hydronephrosis **due** to ...

[An extremely increased CA19-9 level due to common bile duct stone ...](#)

amj.amegroups.com/article/view/3666/4397 ▼ [翻译此页](#)

作者: X Wang - 2017 - [相关文章](#)

2017年2月9日 - Abstract: **Serum carbohydrate antigen** 19-9 (CA19-9), a common tumor ... that CA19-9 level was more than 1,000.0 U/mL (normal **range**: 0-37 U/mL), ... Serum carbohydrate antigen 19-9 (CA19-9) is mainly used for the **diagnosis** and ... **acute cholangitis**, acute pancreatitis, diabetes, and liver cirrhosis (3).

找到约 4,500 条结果 (用时 0.55 秒)

小提示： 仅限搜索简体中文结果。您可以在设置中指定搜索语言

Elevated serum level of carbohydrate antigen 19-9 in benign biliary ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3992417/> ▾ 翻译此页

作者：MS Lin - 2014 - 被引用次数：22 - 相关文章

2014年3月15日 - Although **carbohydrate antigen** (CA19-9) **level** is frequently upregulated in ... The latter, including inflammatory stricture **secondary to choledocholithiasis**, Mirizzi syndrome, The **diagnostic value** of CA19-9 might be improved by using a more ... **Cholangitis** may result in **high serum levels** of CA19-9 [22].

缺少字词：499

Highly elevated serum levels of CA 19-9 in choledocholithiasis: a case ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2740064/> ▾ 翻译此页

作者：G Marcouizos - 2009 - 被引用次数：21 - 相关文章

2009年7月30日 - However, **high levels** of **serum CA** 19-9 is occasionally found in benign disease of ... stone with **acute cholangitis** presenting with extraordinarily **high serum CA 19-9 levels**, ... A **CA 19-9 value** of more than 1.000 U/ml usually indicates a The clinical utility of **serum CA 19-9** in the **diagnosis**, prognosis and ...

Benign Hydronephrosis and Elevated of Serum Levels of ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4917065/> ▾ 翻译此页

作者：B Filipovic - 2016 - 被引用次数：4 - 相关文章

2016年6月11日 - There is no doubt that **CA 19-9 level** is occasionally **elevated** in ... in the differential **diagnosis** of **elevated** concentrations of **CA 19-9** marker. ... track such as pancreatitis, **cholangitis**, and **choledocholithiasis** [5–7]. **Serum carbohydrate antigen 19-9 levels** in patients with unilateral hydronephrosis **due** to ...

找到约 2,970 条结果 (用时 0.42 秒)

小提示：仅限搜索简体中文结果。您可以在设置中指定搜索语言

Elevated serum level of carbohydrate antigen 19-9 in benign biliary ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3992417/> ▼ 翻译此页

作者：MS Lin - 2014 - 被引用次数：23 - 相关文章

2014年3月15日 - Although **carbohydrate antigen** (CA19-9) **level** is frequently upregulated in ... The latter, including inflammatory stricture **secondary to choledocholithiasis**, Mirizzi syndrome, The **diagnostic value** of CA19-9 might be improved by using a more ... **Cholangitis** may result in **high serum levels** of CA19-9 [22].

[PDF] Extraordinarily elevated serum levels of CA 19-9 and rapid decrease ...

www.turkjgastroenterol.org/sayilar/220/buyuk/8641.pdf ▼ 翻译此页

2010年1月15日 - drate antigen 19-9 **value** of >1,000 U/ml usually indicates a di- gestive cancer and has been ... sis and **choledocholithiasis** with **acute cholangitis** who had very **high serum levels** of **carbohydrate antigen** 19-9 (9586 IU/ml). The rapid decrease in ... for the **diagnosis** of pancreatic cancer. A CA 19-9 **value** of ...

Oncotarget | Significant increased CA199 levels in acute pancreatitis ...

www.oncotarget.com/fulltext/23993 ▼ 翻译此页

2018年2月27日 - Background and study aims: **Carbohydrate antigen** 19-9 (CA199) has ... Patients with **high CA199 levels** were older and had predominantly biliary The mean **CA199 value** in 8 patients with pancreatic cancer was 338.2 ... **levels** [25], and this rate was even higher when **acute cholangitis** was included.