

9

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 40483

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Second primary malignancy risk after radiotherapy in rectal cancer survivors

Wang TH *et al.* SPM risk for irradiated rectal cancer

7

Ti-Hao Wang, Chia-Jen Liu, Tze-Fan Chao, Tzeng-Ji Chen, Yu-Wen Hu

Abstract

AIM

To investigate second primary malignancy (SPM) risk after radiotherapy in rectal cancer survivors

Match Overview

1	Crossref 37 words Lee, Yu-Ting, Chia-Jen Liu, Yu-Wen Hu, Chung-Jen Teng, Cheng-Hwai Tzeng, Chiu-Mei Yeh, Tzeng-Ji Chen, Je	1%
2	Internet 33 words crawled on 02-Aug-2018 onlinelibrary.wiley.com	1%
3	Internet 26 words crawled on 06-Apr-2018 www.bjs.co.uk	1%
4	Crossref 26 words Rene Warschkow, Ulrich Güller, Thomas Cerny, Bruno Martin Schmied, Ludwig Plasswilm, Paul Martin Putora. "S	1%
5	Crossref 18 words de Gonzalez, A.B.. "Proportion of second cancers attributable to radiotherapy treatment in adults: a cohort study ...	<1%
6	Internet 17 words crawled on 10-Sep-2017 skemman.is	<1%
7	Internet 15 words crawled on 23-Sep-2017 www.nejm.org	<1%
8	Internet 14 words crawled on 17-Aug-2015 www.science.gov	<1%
9	Internet 14 words crawled on 05-Oct-2017 real.mtak.hu	<1%

找到约 5,500,000 条结果 (用时 0.59 秒)

Incidence of Second Primary Malignancies Following Colorectal Cancer

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4504576/> - 翻译此页

作者: YT Lee - 2015 - 被引用次数: 12 - 相关文章

2015年7月2日 - In addition, the overall survival of patients with advanced CRC was significantly

The risk of subsequent primary cancers after colorectal cancer in Proportion of second cancers attributable to radiotherapy treatment in ...

Abstract · INTRODUCTION · PATIENTS AND METHODS · DISCUSSION

Second primary malignancy risk after radioactive iodine treatment for ...

<https://www.ncbi.nlm.nih.gov/pubmed/19281429> ▼ 翻译此页

作者: AM Sawka - 2009 - 被引用次数: 278 - 相关文章

Second primary malignancy risk after radioactive iodine treatment for thyroid ... therapies is an important concern of thyroid cancer survivors and physicians. ... prior RAI treatment: bladder, breast, central nervous system, colon and rectum, ... Risk Factors; Thyroid Neoplasms/radiotherapy*; Time Factors; Treatment Outcome ...

Second Malignancies Among Elderly Survivors of Cancer

theoncologist.alphamedpress.org/content/16/11/1572.full ▼ 翻译此页

作者: AM VanderWalde - 2011 - 被引用次数: 23 - 相关文章

2011年10月31日 - Although the risks for the development of second malignancies following a first ...

Primary malignancies and association with second cancers [14] ... However, in women aged <45 years at the time of radiation therapy, the RR was For survivors of colorectal cancer, the NCCN recommends colonoscopy at ...

Young patients with colorectal cancer have increased risk of second ...

<https://academic.oup.com/jjco/article/45/11/1029/2385102> - 翻译此页

作者: YH Liang - 2015 - 被引用次数: 4 - 相关文章

2015年9月18日 - Because the number of long-term survivors of colorectal cancer has increased, second ... Radiotherapy, 309, 1.25^a, 1.12–1.40, 1.85^a, 1.14, 1.08, 1.04, 0.97 Increased risks of secondary primary cancers in young patients.

Abstract · Introduction · Results · Discussion

找到约 5,500,000 条结果 (用时 0.59 秒)

Incidence of Second Primary Malignancies Following Colorectal Cancer

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4504576/> - 翻译此页

作者: YT Lee - 2015 - 被引用次数: 12 - 相关文章

2015年7月2日 - In addition, the overall survival of patients with advanced CRC was significantly

The risk of subsequent primary cancers after colorectal cancer in Proportion of second cancers attributable to radiotherapy treatment in ...

Abstract · INTRODUCTION · PATIENTS AND METHODS · DISCUSSION

Second primary malignancy risk after radioactive iodine treatment for ...

<https://www.ncbi.nlm.nih.gov/pubmed/19281429> ▼ 翻译此页

作者: AM Sawka - 2009 - 被引用次数: 278 - 相关文章

Second primary malignancy risk after radioactive iodine treatment for thyroid ... therapies is an important concern of thyroid cancer survivors and physicians. ... prior RAI treatment: bladder, breast, central nervous system, colon and rectum, ... Risk Factors; Thyroid Neoplasms/radiotherapy*; Time Factors; Treatment Outcome ...

Second Malignancies Among Elderly Survivors of Cancer

theoncologist.alphamedpress.org/content/16/11/1572.full ▼ 翻译此页

作者: AM VanderWalde - 2011 - 被引用次数: 23 - 相关文章

2011年10月31日 - Although the risks for the development of second malignancies following a first ...

Primary malignancies and association with second cancers [14] ... However, in women aged <45 years at the time of radiation therapy, the RR was For survivors of colorectal cancer, the NCCN recommends colonoscopy at ...

Young patients with colorectal cancer have increased risk of second ...

<https://academic.oup.com/jjco/article/45/11/1029/2385102> - 翻译此页

作者: YH Liang - 2015 - 被引用次数: 4 - 相关文章

2015年9月18日 - Because the number of long-term survivors of colorectal cancer has increased, second ... Radiotherapy, 309, 1.25^a, 1.12–1.40, 1.85^a, 1.14, 1.08, 1.04, 0.97 Increased risks of secondary primary cancers in young patients.

Abstract · Introduction · Results · Discussion



Second primary malignancy risk after radiotherapy in rectal cancer survi



全部

新闻

图片

视频

购物

更多

设置

工具

找到约 5,150,000 条结果 (用时 0.66 秒)

Google 学术: Second primary malignancy risk after radiotherapy in rectal cancer survivors

..., and leucovorin for metastatic colorectal cancer - Hurwitz - 被引用次数: 10645

Second malignancies in prostate carcinoma patients ... - Brenner - 被引用次数: 590

... radiation therapy, protons, and the risk of second ... - Hall - 被引用次数: 998

Second primary malignancy risk after radioactive iodine treatment for ...

<https://www.ncbi.nlm.nih.gov/pubmed/19281429> ▼ 翻译此页

作者: AM Sawka - 2009 - 被引用次数: 284 - 相关文章

Second primary malignancy risk after radioactive iodine treatment for thyroid ... therapies is an important concern of thyroid cancer survivors and physicians. ... prior RAI treatment: bladder, breast, central nervous system, colon and rectum, ... Risk Factors; Thyroid Neoplasms/radiotherapy*; Time Factors; Treatment Outcome ...

Incidence of Second Primary Malignancies Following Colorectal Cancer

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4504576/> - 翻译此页

作者: YT Lee - 2015 - 被引用次数: 11 - 相关文章

2015年7月2日 - The risk for breast, bladder, kidney, lung, and uterine cancers was significantly higher in patients with colon and rectal cancers than the general population. ... In conclusion, patients with CRC were at increased risk for a second cancer. The pattern of SPMs was distinct between patients with colon and rectal cancer.

Abstract · INTRODUCTION · PATIENTS AND METHODS · DISCUSSION

Estimation of second cancer risk after radiotherapy for rectal cancer ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5103599/> ▼ 翻译此页

作者: DR Zwahlen - 2016 - 被引用次数: 5 - 相关文章

2016年11月10日 - Keywords: Second malignancies, Rectal cancer, Radiotherapy, ... Second cancer risk (SCR) is of concern in long-term radiotherapy (RT) survivors [1]. In their of second primary malignancies occurring after radiotherapy?