

World Journal of *Transplantation*

World J Transplant 2018 October 22; 8(6): 198-236





EDITORIAL

- 198 Surgeon's perspective on short bowel syndrome: Where are we?
Marino IR, Lauro A

REVIEW

- 203 Complement-mediated renal diseases after kidney transplantation - current diagnostic and therapeutic options in *de novo* and recurrent diseases
Abbas F, El Kossi M, Kim JJ, Shaheen IS, Sharma A, Halawa A

SYSTEMATIC REVIEW

- 220 Impact of machine perfusion of the liver on post-transplant biliary complications: A systematic review
Boteon YL, Boteon AP, Attard J, Wallace L, Bhogal RH, Afford SC

CASE REPORT

- 232 Treatment of transplant renal artery pseudoaneurysm using expandable hydrogel coils: A case report and review of literature
Marie Y, Kumar A, Hinchliffe S, Curran S, Brown P, Turner D, Shrestha B

ABOUT COVER

Editorial Board Member of *World Journal of Transplantation*, Yuri L Boteon, MD, PhD, Academic Research, Doctor, Surgeon, The Liver Transplant and Hepatobiliary Surgery Unit, Queen Elizabeth Hospital Birmingham, Birmingham B15 2TT, United Kingdom

AIM AND SCOPE

World Journal of Transplantation (*World J Transplant*, *WJT*, online ISSN 2220-3230, DOI: 10.5500) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJT covers topics concerning organ and tissue donation and preservation; tissue injury, repair, inflammation, and aging; immune recognition, regulation, effector mechanisms, and opportunities for induction of tolerance, thoracic transplantation (heart, lung), abdominal transplantation (kidney, liver, pancreas, islets), transplantation of tissues, cell therapy and islet transplantation, clinical transplantation, experimental transplantation, immunobiology and genomics, and xenotransplantation. The current columns of *WJT* include editorial, frontier, diagnostic advances, therapeutics advances, field of vision, mini-reviews, review, topic highlight, medical ethics, original articles, case report, clinical case conference (Clinicopathological conference), and autobiography.

AIM AND SCOPE

World Journal of Transplantation (*WJT*) is now abstracted and indexed in PubMed, PubMed Central, China National Knowledge Infrastructure (CNKI), and Superstar Journals Database.

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Yan Hnang*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Ying Dou*
Proofing Editorial Office Director: *Jin-Lai Wang*

NAME OF JOURNAL
World Journal of Transplantation

ISSN
ISSN 2220-3230 (online)

LAUNCH DATE
December 24, 2011

EDITOR-IN-CHIEF
Maurizio Salvadori, MD, Professor, Renal Unit, Careggi University Hospital, Florence 50139, Italy

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com/2220-3230/editorialboard.htm>

EDITORIAL OFFICE
Jin-Lai Wang, Director
World Journal of Transplantation
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA

Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
October 22, 2018

COPYRIGHT
© 2018 Baishideng Publishing Group Inc. Articles

published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
<http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.f6publishing.com>

Surgeon's perspective on short bowel syndrome: Where are we?

Ignazio R Marino, Augusto Lauro

Ignazio R Marino, Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia PA 19107, United States

Augusto Lauro, Liver and Multiorgan Transplant Unit, St. Orsola-Malpighi University Hospital, Bologna 40138, Italy

ORCID number: Ignazio R Marino (0000-0002-1519-8385); Augusto Lauro (0000-0002-2292-5595).

Author contributions: Marino IR and Lauro A conceived the study and drafted the manuscript; both authors approved the final version of the article.

Conflict-of-interest statement: The authors have no conflict of interest to declare.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited Manuscript

Correspondence to: Augusto Lauro, MD, PhD, Surgeon, Liver and Multiorgan Transplant Unit, St. Orsola-Malpighi University Hospital, Via Massarenti 9, Bologna 40138, Italy. augustola@yahoo.com
Telephone: +39-051-6363721

Received: July 4, 2018

Peer-review started: July 4, 2018

First decision: August 20, 2018

Revised: August 25, 2018

Accepted: October 8, 2018

Article in press: October 8, 2018

Published online: October 22, 2018

Abstract

Short bowel syndrome (SBS) is due to a massive loss of

small bowel: the reduction of gut function is below the minimum necessary to maintain health (in adults) and growth (in children) so intravenous supplementation is required. Parenteral nutrition represents the milestone of treatment and surgical attempts should be limited only when the residual bowel is sufficient to increase absorption, reducing diarrhea and slowing the transit time of nutrients, water and electrolytes. The surgical techniques lengthen the bowel (tapering it) or reverse a segment of it: developed in children, nowadays are popular also among adults. The issue is mainly represented by the residual length of the small bowel where ileum has shown increased adaptive function than jejunum, but colon should be considered because of its importance in the digestive process. These concepts have been translated also in intestinal transplantation, where a colonic graft is nowadays widely used and the terminal ileum is the selected segment for a living-related donation. The whole replacement by a bowel or multivisceral transplant is still affected by poor long term outcome and must be reserved to a select population of SBS patients, affected by intestinal failure associated with irreversible complications of parenteral nutrition.

Key words: Parenteral nutrition; Bowel rehabilitation; Surgical rescue; Intestinal transplantation; Short bowel syndrome

© **The Author(s) 2018.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Short bowel syndrome represents a surgical dilemma: parenteral nutrition is considered the gold standard of care and any surgical attempt must be limited by the universal principle "first do not harm." The surgical rehabilitation should be pursued when there are enough residual intestines to obtain a better bowel function: lengthening the intestine or reversing a loop of it with different techniques should have the only aim of slowing the transit while increasing the absorptive surface. When intestinal failure is associated to life-threatening parenteral nutrition complications, bowel transplantation should be considered as an option.

Marino IR, Lauro A. Surgeon's perspective on short bowel syndrome: Where are we? *World J Transplant* 2018; 8(6): 198-202 Available from: URL: <http://www.wjgnet.com/2220-3230/full/v8/i6/198.htm> DOI: <http://dx.doi.org/10.5500/wjt.v8.i6.198>

INTRODUCTION

Short bowel syndrome (SBS) results from a reduced length of the small intestine. A "normal small bowel length," measured from the duodeno-jejunal flexure to ileocolic valve, is estimated at 250 cm \pm 40 cm at birth, and the growth is maximal during the first year of life^[1]. In adults, the small bowel length varies from 275 cm to 850 cm, with a mean of 350 cm \pm 60 cm, depending on the method used, radiologic, surgical, or per autopsy^[2]. The massive loss of small bowel represents the most frequent mechanism of intestinal failure, defined by the European Society for Clinical Nutrition and Metabolism as "the reduction of gut function below the minimum necessary for the absorption of macronutrients and/or water and electrolytes, such that intravenous supplementation is required to maintain health and/or growth"^[3]. Among children "the minimum necessary for the absorption" is a residual small bowel length of more than 25% of the expected for gestational age^[3], in adults SBS usually appears when the small bowel length is less than 200 cm (67% of the normal length)^[4]. Malabsorption and diarrhea represent the classical symptoms, associated to deficit of growth in the pediatric population. Wilmore *et al*^[5] first demonstrated long-term survival with parenteral nutrition (PN) in a child affected by SBS. Nowadays home PN represents the standard of care in patients affected by massive loss of small bowel with excellent long term results^[6-12]. PN does not replace physiologically the bowel function because uses the intravenous route to supplement nutrients and it could be affected by several life-threatening complications. Under this perspective, a surgical rehabilitation in case of SBS should be represented by: (1) the possibility to slow the transit and obtain an adequate absorptive surface of the remnant intestine through lengthening procedures and (2) whole replacement of the massive intestinal loss with a bowel transplant. It is worthwhile to analyze briefly the main reported studies on the issue in international literature, in order to develop an updated perspective under the surgical point of view.

OVERVIEW OF THE LITERATURE

SBS is mainly, but not only, a matter of length. In children, the massive resection of the small bowel could lead to a "very short bowel syndrome" (\leq 40 cm)^[13,14], "ultra-short bowel syndrome" (between < 30 and < 10 cm)^[15-17] or "no gut syndrome" (only the duodenum is left)^[18-20]. Adults with less than 200 cm but more than 75 cm of small bowel^[21] have a potentially functional intestine especially if the colon (and specifically the

ileocolic valve) is preserved in continuity. Among SBS patients, the role of the colon in the process of digestion has been demonstrated since the '90s^[22-25]. The presence of remaining colon is associated with a lower dependency on PN^[26,27] and there is agreement that the remaining small bowel after massive intestinal loss is supported by the colon (if in continuity) for completion of the digestion process. On the other hand, jejunum and ileum have different roles in digestion and ileum has probably a greater adaptive potential than jejunum^[28]. A remnant ileum (especially in continuity with the colon) could probably guarantee a faster weaning from PN. Clinical experience shows that patients with a jejuno-colonic anastomosis (SBS type II), even better with a jejuno-ileo-colonic anastomosis (SBS type III), have an improved absorption with time after a period of intestinal rehabilitation, whereas patients with end-terminal jejunostomy without colon (SBS type I) do not show that. When the colon is missing, among adults 115 cm of small bowel with an end enterostomy are considered the limit before SBS.

SURGEON'S PERSPECTIVE

In SBS the remaining small bowel may dilate. This is important for surgeons in order to lengthen the intestine, tapering it. It has been shown that the extent of dilation is associated with the bowel length, and both are related to enteral autonomy^[29]. Two surgical procedures are popular in order to lengthen the bowel: Bianchi and Serial Transverse Enteroplasty Procedure (STEP). The Bianchi procedure, summarized by Bianchi in 1997^[30], is also known as longitudinal lengthening and tailoring (LILT). The small bowel mesentery is separated as two leaves with a GI anastomosis stapler to create a tunnel, and then the two resulting small bowel segments of smaller diameter are connected with an end-to-end anastomosis in an iso-peristaltic fashion. In the STEP, first described by Kim in 2003^[31], the dilated small bowel is narrowed by serial transverse applications of the GI stapler from opposite directions, creating a new lengthened small intestine (zig-zag channel). This procedure does not require an intestinal anastomosis and the mesenteric vascular supply is untouched. Since its first description, STEP has become a widespread procedure, sometimes repeated on the same patient (re-STEP) to obtain a longer intestinal segment. Bianchi and STEP procedures have been performed at first in children and more recently also in adults^[32-35]. Most of the studies are on STEP: while enteral autonomy (median time: 21 mo) is eventually possible in some patients^[36], improved enteral tolerance can be achieved in a majority^[37,38]. STEP can be performed on shorter intestinal segments or intricate segments such as the duodenum, which is technically not feasible for Bianchi procedure, and it seems to have a lower mortality but an overall progression to transplantation^[39]. The spiral intestinal lengthening and tailoring procedure is a new

surgical technique based on a spiral shape incision of the dilated intestine (at 45°-60° to its longitudinal axis), and re-tubularization in a longer but narrower fashion. It does not alter the orientation of the muscle fibers like STEP, offering minimal mesenteric handling compared to Bianchi procedure. It has been reported in a 3-year-old girl^[40] where, 6 mo after the procedure, PN was weaned off. Another manuscript described the technique in a 10-month-old child^[41] showing at 1-year follow-up a growth on the 15-25th centile on 82% oral calories and 18% PN, passing 2-3 daily stools. Three children with "no gut" syndrome and dilated duodenum underwent a novel surgical procedure of "duodenal lengthening" combined with a technical modification of STEP^[18]: duodenal tapering was performed with sequential transverse applications of an endoscopic stapler on the anterior and posterior wall of the duodenum, avoiding bilio-pancreatic injury. Two patients weaned PN off at 12 mo post-surgery and the last one's PN caloric requirements decreased by 60%. The surgical rescue of "no gut" syndrome has been reported in adults as well. Bueno *et al*^[20] demonstrated the feasibility of lengthening a dilated duodenum in a patient where his mega-duodenal stump was tapered by STEP, restoring his digestive continuity through an end-to-side duodeno-colonic anastomosis. After 24 mo of follow-up, the time on daily PN was shortened from 24 to 9 h and the volume and calorie requirements were reduced by half.

Since lengthening procedures slow the bowel transit time, a "reversed anti-peristaltic segmental bowel loop" has been proposed with the same aim: this procedure can be indicated in patients with an adequate remnant bowel length. Median oral autonomy was described up to 100% \pm 38% with a lower amount of parenteral calories, as well as PN dependence^[42]. In another report^[43] 56% of patients improved their enteral autonomy.

The different graft types used in intestinal transplantation are the isolated small bowel, combined liver-intestine, multivisceral and modified multivisceral ones^[44]: liver-containing grafts have shown the longest survivals. Apart from cadaveric donation, living-related intestinal transplantation has been pursued especially in a pediatric setting^[45]: terminal ileum represents the used graft, because of technical feasibility and its greater adaptive potential than jejunum^[28]. Short term results of intestinal transplantation have recently improved in terms of survival and digestive autonomy, due to advances in surgery and immunosuppression. Immunosuppressive therapy has evolved significantly over the past 20 years: the tacrolimus-based therapy as maintenance, preceded by induction with anti-thymocyte globulin or an interleukin-2 blocker, is the main used protocol worldwide. A "secondary" agent like steroids, azathioprine, mycophenolate mofetil or an mTOR inhibitor is recommended after an episode of rejection. Innovative cross match strategies and optimizing organ allocation could improve the long-term outcome, but the main causes of death and graft loss remain sepsis

Table 1 Surgical rehabilitation of short bowel syndrome

SBS surgical rehabilitation
Lengthening procedures
Bianchi
STEP
SILT
Duodenal lengthening
Reversed anti-peristaltic segmental bowel loop
Intestinal transplantation

SBS: Short bowel syndrome; STEP: Serial transverse enteroplasty procedure; SILT: Spiral intestinal lengthening and tailoring.

and rejection. Challenges for long-term results are chronic rejection and immunosuppressant-related complications^[46,47]. According to Intestinal Transplant Registry reports^[44], 1611 children were transplanted worldwide between 1985 and 2013, with an overall patient survival rate of 51%. In the 2014-2016 Scientific Registry of Transplant Recipients^[48], the 6 American centers that in 2016 performed 10 or more intestinal transplants in adults reported a 1-year graft survival from 61% to 83% and a 3-year graft survival from 29% to 73%. In an earlier report from 2008 to 2010, the 1-year graft survival in adults was 71%, illustrating the relatively modest gains achieved^[47]. Intestinal transplantation should be suggested to a very select subset of SBS patients with severe and irreversible complications of PN and no hope of intestinal rehabilitation. In conclusion, among SBS patients the surgical rehabilitation (Table 1) of the remnant bowel must be performed to slow the intestinal transit time increasing at the same time the absorptive surface: only in cases of irreversible intestinal failure with PN life-threatening complications, intestinal transplantation could represent a therapeutic option even if still encumbered by suboptimal long term results.

ACKNOWLEDGMENTS

We wish to thank Ms. Claudia Cirillo for her English language editing of our text.

REFERENCES

- 1 Siebert JR. Small-intestine length in infants and children. *Am J Dis Child* 1980; **134**: 593-595 [PMID: 7386434]
- 2 Gondolessi G, Ramisch D, Padin J, Almau H, Sandi M, Schelotto PB, Fernandez A, Rumbo C, Solar H. What is the normal small bowel length in humans? first donor-based cohort analysis. *Am J Transplant* 2012; **12 Suppl 4**: S49-S54 [PMID: 22702412 DOI: 10.1111/j.1600-6143.2012.04148.x]
- 3 Pironi L. Definitions of intestinal failure and the short bowel syndrome. *Best Pract Res Clin Gastroenterol* 2016; **30**: 173-185 [PMID: 27086884 DOI: 10.1016/j.bpg.2016.02.011]
- 4 Fung JJ. William Hunter Harridge lecture: The changing face of short-gut syndrome management. *Am J Surg* 2017; **213**: 448-451 [PMID: 28159115 DOI: 10.1016/j.amjsurg.2017.01.018]
- 5 Wilmore DW, Dudrick SJ. Growth and development of an infant receiving all nutrients exclusively by vein. *JAMA* 1968; **203**: 860-864 [PMID: 4965871]
- 6 Pironi L, Goulet O, Buchman A, Messing B, Gabe S, Candusso M,

- Bond G, Gupte G, Pertkiewicz M, Steiger E, Forbes A, Van Gossum A, Pinna AD; Home Artificial Nutrition and Chronic Intestinal Failure Working Group of ESPEN. Outcome on home parenteral nutrition for benign intestinal failure: a review of the literature and benchmarking with the European prospective survey of ESPEN. *Clin Nutr* 2012; **31**: 831-845 [PMID: 22658443 DOI: 10.1016/j.clnu.2012.05.004]
- 7 **Scolapio JS**, Fleming CR, Kelly DG, Wick DM, Zinsmeister AR. Survival of home parenteral nutrition-treated patients: 20 years of experience at the Mayo Clinic. *Mayo Clin Proc* 1999; **74**: 217-222 [PMID: 10089988 DOI: 10.4065/74.3.217]
 - 8 **Messing B**, Lémann M, Landais P, Gouttebel MC, Gérard-Boncompain M, Saudin F, Vangossum A, Beau P, Guédon C, Barnoud D. Prognosis of patients with nonmalignant chronic intestinal failure receiving long-term home parenteral nutrition. *Gastroenterology* 1995; **108**: 1005-1010 [PMID: 7698566]
 - 9 **Colomb V**, Dabbas-Tyan M, Taupin P, Talbotec C, Révillon Y, Jan D, De Potter S, Gorski-Colin AM, Lamor M, Herreman K, Corriol O, Landais P, Ricour C, Goulet O. Long-term outcome of children receiving home parenteral nutrition: a 20-year single-center experience in 302 patients. *J Pediatr Gastroenterol Nutr* 2007; **44**: 347-353 [PMID: 17325556 DOI: 10.1097/MPG.0b013e31802c6971]
 - 10 **Pironi L**, Joly F, Forbes A, Colomb V, Lyszkowska M, Baxter J, Gabe S, Hébuterne X, Gambarara M, Gottrand F, Cuerda C, Thul P, Messing B, Goulet O, Staun M, Van Gossum A; Home Artificial Nutrition & Chronic Intestinal Failure Working Group of the European Society for Clinical Nutrition and Metabolism (ESPEN). Long-term follow-up of patients on home parenteral nutrition in Europe: implications for intestinal transplantation. *Gut* 2011; **60**: 17-25 [PMID: 21068130 DOI: 10.1136/gut.2010.223255]
 - 11 **Amiot A**, Messing B, Corcos O, Panis Y, Joly F. Determinants of home parenteral nutrition dependence and survival of 268 patients with non-malignant short bowel syndrome. *Clin Nutr* 2013; **32**: 368-374 [PMID: 22992308 DOI: 10.1016/j.clnu.2012.08.007]
 - 12 **Messing B**, Crenn P, Beau P, BOUTRON-RUAULT MC, RAMBAUD JC, MATUCHANSKY C. Long-term survival and parenteral nutrition dependence in adult patients with the short bowel syndrome. *Gastroenterology* 1999; **117**: 1043-1050 [PMID: 10535866]
 - 13 **Capriati T**, Giorgio D, Fusaro F, Candusso M, Schingo P, Caldaro T, Laureti F, Elia D, Diamanti A. Pediatric Short Bowel Syndrome: Predicting Four-Year Outcome after Massive Neonatal Resection. *Eur J Pediatr Surg* 2018; **28**: 455-463 [PMID: 28719916 DOI: 10.1055/s-0037-1604113]
 - 14 **Norsa L**, Artru S, Lambe C, Talbotec C, Pigneur B, Ruemmele F, Colomb V, Capito C, Chardot C, Lacaille F, Goulet O. Long term outcomes of intestinal rehabilitation in children with neonatal very short bowel syndrome: Parenteral nutrition or intestinal transplantation. *Clin Nutr* 2018; : [PMID: 29478887 DOI: 10.1016/j.clnu.2018.02.004]
 - 15 **Tannuri U**, Barros F, Tannuri AC. Treatment of short bowel syndrome in children. Value of the Intestinal Rehabilitation Program. *Rev Assoc Med Bras (1992)* 2016; **62**: 575-583 [PMID: 27849236 DOI: 10.1590/1806-9282.62.06.575]
 - 16 **Batra A**, Keys SC, Johnson MJ, Wheeler RA, Beattie RM. Epidemiology, management and outcome of ultrashort bowel syndrome in infancy. *Arch Dis Child Fetal Neonatal Ed* 2017; **102**: F551-F556 [PMID: 28866623 DOI: 10.1136/archdischild-2016-311765]
 - 17 **Dore M**, Junco PT, Moreno AA, Cerezo VN, Muñoz MR, Galán AS, Sánchez AV, Prieto G, Ramos E, Hernandez F, Martínez LM, Santamaria ML. Ultrashort Bowel Syndrome Outcome in Children Treated in a Multidisciplinary Intestinal Rehabilitation Unit. *Eur J Pediatr Surg* 2017; **27**: 116-120 [PMID: 28052307 DOI: 10.1055/s-0036-1597812]
 - 18 **Bueno J**, Redecillas S, García L, Lara A, Giné C, Molino JA, Broto J, Segarra O. Duodenal lengthening in short bowel with dilated duodenum. *J Pediatr Surg* 2015; **50**: 493-496 [PMID: 25746715 DOI: 10.1016/j.jpedsurg.2014.11.047]
 - 19 **Jain V**, Huerta S. More on 'No Gut Syndrome': A case report. *Int J Surg Case Rep* 2016; **19**: 35-37 [PMID: 26708947 DOI: 10.1016/j.ijscr.2015.12.014]
 - 20 **Bueno J**, Burgos R, Redecillas S, López M, Balsells J. Duodenal lengthening in an adult with ultra-short bowel syndrome. A case report. *Rev Esp Enferm Dig* 2018; **110**: 59-62 [PMID: 29106286 DOI: 10.17235/reed.2017.5187/2017]
 - 21 **Lauro A**, Cirocchi R, Cautero N, Dazzi A, Pironi D, Di Matteo FM, Santoro A, Pironi L, Pinna AD. Reconnection surgery in adult post-operative short bowel syndrome < 100 cm: is colonic continuity sufficient to achieve enteral autonomy without autologous gastrointestinal reconstruction? Report from a single center and systematic review of literature. *G Chir* 2017; **38**: 163-175 [PMID: 29182898]
 - 22 **Cosnes J**, Gendre JP, Le Quintrec Y. Role of the ileocecal valve and site of intestinal resection in malabsorption after extensive small bowel resection. *Digestion* 1978; **18**: 329-336 [PMID: 750260 DOI: 10.1159/000198220]
 - 23 **Nordgaard I**. What's new in the role of colon as a digestive organ in patients with short bowel syndrome. *Nutrition* 1998; **14**: 468-469 [PMID: 9614315]
 - 24 **Jeppesen PB**, Mortensen PB. Colonic digestion and absorption of energy from carbohydrates and medium-chain fat in small bowel failure. *JPEN J Parenter Enteral Nutr* 1999; **23**: S101-S105 [PMID: 10483907 DOI: 10.1177/014860719902300525]
 - 25 **Smith KH**, Saunders JA, Nugent KP, Jackson AA, Stroud MA. Reduced parenteral nutrition requirements following anastomosis of a short residual colonic segment to a short jejunum. *JPEN J Parenter Enteral Nutr* 2011; **35**: 732-735 [PMID: 22042049 DOI: 10.1177/0148607111406504]
 - 26 **Nguyen BT**, Blatchford GJ, Thompson JS, Bragg LE. Should intestinal continuity be restored after massive intestinal resection? *Am J Surg* 1989; **158**: 577-9; discussion 579-80 [PMID: 2511774]
 - 27 **Thompson JS**. Reoperation in patients with the short bowel syndrome. *Am J Surg* 1992; **164**: 453-6; discussion 456-7 [PMID: 1443368]
 - 28 **Kong W**, Wang J, Ying R, Li Y, Jin H, Mao Q, Yao D, Guo M. A potential anatomic subtype of short bowel syndrome: a matched case-control study. *BMC Gastroenterol* 2016; **16**: 12 [PMID: 26822147 DOI: 10.1186/s12876-016-0425-4]
 - 29 **Ives GC**, Demehri FR, Sanchez R, Barrett M, Gadepalli S, Teitelbaum DH. Small Bowel Diameter in Short Bowel Syndrome as a Predictive Factor for Achieving Enteral Autonomy. *J Pediatr* 2016; **178**: 275-277.e1 [PMID: 27587075 DOI: 10.1016/j.jpeds.2016.08.007]
 - 30 **Bianchi A**. Longitudinal intestinal lengthening and tailoring: results in 20 children. *J R Soc Med* 1997; **90**: 429-432 [PMID: 9306995]
 - 31 **Kim HB**, Fauza D, Garza J, Oh JT, Nurko S, Jaksic T. Serial transverse enteroplasty (STEP): a novel bowel lengthening procedure. *J Pediatr Surg* 2003; **38**: 425-429 [PMID: 12632361 DOI: 10.1053/jpsu.2003.50073]
 - 32 **Fan S**, Li Y, Zhang S, Wang J, Li J. Success of serial transverse enteroplasty in an adult with extreme short bowel syndrome: a case report. *Int Surg* 2015; **100**: 626-631 [PMID: 25875543 DOI: 10.9738/INTSURG-D-14-00177.1]
 - 33 **Yaprak M**, Erdoğan O, Oğus M. The STEP procedure in an adult patient with short bowel syndrome: a case report. *Turk J Gastroenterol* 2011; **22**: 333-336 [PMID: 21805426]
 - 34 **Bellolio R F**, Klaassen L J, Pulgar B D, Molina P ME, Pinedo M G, Zúñiga D A. Serial transverse enteroplasty for short bowel syndrome. Case report. *Rev Med Chil* 2010; **138**: 478-482 [PMID: 20668797]
 - 35 **Yannam GR**, Sudan DL, Grant W, Botha J, Langnas A, Thompson JS. Intestinal lengthening in adult patients with short bowel syndrome. *J Gastrointest Surg* 2010; **14**: 1931-1936 [PMID: 20734155 DOI: 10.1007/s11605-010-1291-y]
 - 36 **Jones BA**, Hull MA, Potanos KM, Zurakowski D, Fitzgibbons SC, Ching YA, Duggan C, Jaksic T, Kim HB; International STEP Data Registry. Report of 111 consecutive patients enrolled in the International Serial Transverse Enteroplasty (STEP) Data Registry: a retrospective observational study. *J Am Coll Surg* 2013; **216**: 438-446 [PMID: 23357726 DOI: 10.1016/j.jamcollsurg.2012.12.018]
 - 37 **Oh PS**, Fingeret AL, Shah MY, Ventura KA, Brodlië S, Ovchinsky

- N, Martinez M, Lobritto SJ, Cowles RA. Improved tolerance for enteral nutrition after serial transverse enteroplasty (STEP) in infants and children with short bowel syndrome--a seven-year single-center experience. *J Pediatr Surg* 2014; **49**: 1589-1592 [PMID: 25475799 DOI: 10.1016/j.jpedsurg.2014.07.019]
- 38 **Mercer DF**, Hobson BD, Gerhardt BK, Grant WJ, Vargas LM, Langnas AN, Quiros-Tejeira RE. Serial transverse enteroplasty allows children with short bowel to wean from parenteral nutrition. *J Pediatr* 2014; **164**: 93-98 [PMID: 24094877 DOI: 10.1016/j.jpeds.2013.08.039]
- 39 **Frangia G**, Kessler M, Weih S, Nickkholgh A, Mehrabi A, Holland-Cunz S. Comparison of LILT and STEP procedures in children with short bowel syndrome -- a systematic review of the literature. *J Pediatr Surg* 2013; **48**: 1794-1805 [PMID: 23932625 DOI: 10.1016/j.jpedsurg.2013.05.018]
- 40 **Cserni T**, Biszku B, Guthy I, Dicso F, Szaloki L, Folaranmi S, Murphy F, Rakoczy G, Bianchi A, Morabito A. The first clinical application of the spiral intestinal lengthening and tailoring (silt) in extreme short bowel syndrome. *J Gastrointest Surg* 2014; **18**: 1852-1857 [PMID: 24957255 DOI: 10.1007/s11605-014-2577-2]
- 41 **Alberti D**, Boroni G, Giannotti G, Parolini F, Armellini A, Morabito A, Bianchi A. "Spiral intestinal lengthening and tailoring (SILT)" for a child with severely short bowel. *Pediatr Surg Int* 2014; **30**: 1169-1172 [PMID: 25119303 DOI: 10.1007/s00383-014-3583-x]
- 42 **Layec S**, Beyer L, Corcos O, Alves A, Dray X, Amiot A, Stefanescu C, Coffin B, Bretagnol F, Bouhnik Y, Messing B, Panis Y, Kapel N, Joly F. Increased intestinal absorption by segmental reversal of the small bowel in adult patients with short-bowel syndrome: a case-control study. *Am J Clin Nutr* 2013; **97**: 100-108 [PMID: 23151533 DOI: 10.3945/ajcn.112.042606]
- 43 **Thompson JS**. Reversed Intestinal Segment Revisited. *Transplant Proc* 2016; **48**: 453-456 [PMID: 27109977 DOI: 10.1016/j.transproc.2015.09.072]
- 44 **Grant D**, Abu-Elmagd K, Mazariegos G, Vianna R, Langnas A, Mangus R, Farmer DG, Lacaille F, Iyer K, Fishbein T; Intestinal Transplant Association. Intestinal transplant registry report: global activity and trends. *Am J Transplant* 2015; **15**: 210-219 [PMID: 25438622 DOI: 10.1111/ajt.12979]
- 45 **Garcia Aroz S**, Tzvetanov I, Hetterman EA, Jeon H, Oberholzer J, Testa G, John E, Benedetti E. Long-term outcomes of living-related small intestinal transplantation in children: A single-center experience. *Pediatr Transplant* 2017; **21**: [PMID: 28295952 DOI: 10.1111/petr.12910]
- 46 **Celik N**, Mazariegos GV, Soltys K, Rudolph JA, Shi Y, Bond GJ, Sindhi R, Ganoza A. Pediatric intestinal transplantation. *Gastroenterol Clin North Am* 2018; **47**: 355-368 [PMID: 29735029 DOI: 10.1016/j.gtc.2018.01.007]
- 47 **Matsumoto CS**, Subramanian S, Fishbein TM. Adult intestinal transplantation. *Gastroenterol Clin North Am* 2018; **47**: 341-354 [PMID: 29735028 DOI: 10.1016/j.gtc.2018.01.011]
- 48 Scientific Registry of Transplant Recipients (SRTR) Program Specific Reports. Available from: <https://www.srtr.org/reports>

P- Reviewer: Nacif LS, Boteon YL **S- Editor:** Dou Y **L- Editor:** A
E- Editor: Huang Y





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

