

World Journal of *Gastroenterology*

World J Gastroenterol 2018 November 28; 24(44): 4959-5056



**EDITORIAL**

- 4959 Hepatitis elimination by 2030: Progress and challenges
Waheed Y, Siddiq M, Jamil Z, Najmi MH

REVIEW

- 4962 Mononuclear phagocyte system in hepatitis C virus infection
Yang Y, Tu ZK, Liu XK, Zhang P

MINIREVIEWS

- 4974 Learning curves in minimally invasive esophagectomy
van Workum F, Fransen L, Luyer MDP, Rosman C
- 4979 Glutathione depleting drugs, antioxidants and intestinal calcium absorption
Moine L, Rivoira M, Díaz de Barboza G, Pérez A, Tolosa de Talamoni N

ORIGINAL ARTICLE**Basic Study**

- 4989 Different distributions of interstitial cells of Cajal and platelet-derived growth factor receptor- α positive cells in colonic smooth muscle cell/interstitial cell of Cajal/platelet-derived growth factor receptor- α positive cell syncytium in mice
Lu C, Huang X, Lu HL, Liu SH, Zang JY, Li YJ, Chen J, Xu WX
- 5005 Development of a novel rat model of heterogeneous hepatic injury by injection with colchicine *via* the splenic vein
Zhang YY, Zhang CX, Li Y, Jiang X, Wang YF, Sun Y, Wang J, Ji WY, Liu Y
- 5013 Involvement of methylation-associated silencing of formin 2 in colorectal carcinogenesis
Li DJ, Feng ZC, Li XR, Hu G

Retrospective Cohort Study

- 5025 Timing of upper gastrointestinal endoscopy does not influence short-term outcomes in patients with acute variceal bleeding
Yoo JJ, Chang Y, Cho EJ, Moon JE, Kim SG, Kim YS, Lee YB, Lee JH, Yu SJ, Kim YJ, Yoon JH
- 5034 Risk factors and prediction score for chronic pancreatitis: A nationwide population-based cohort study
Lin YC, Kor CT, Su WW, Hsu YC

Retrospective Study

- 5046 Prognostic value of fibrinogen and D-dimer-fibrinogen ratio in resectable gastrointestinal stromal tumors
Cai HX, Li XQ, Wang SF

ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Wan-Long Chuang, MD, PhD, Doctor, Professor, Hepatobiliary Division, Department of Internal Medicine, Kaohsiung Medical University Hospital, Kaohsiung Medical University, Kaohsiung 807, Taiwan

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7th, 14th, 21st, and 28th each month. The *WJG* Editorial Board consists of 642 experts in gastroenterology and hepatology from 59 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

World Journal of Gastroenterology (*WJG*) is now indexed in Current Contents[®]/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch[®]), Journal Citation Reports[®], Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2018 edition of Journal Citation Reports[®] cites the 2017 impact factor for *WJG* as 3.300 (5-year impact factor: 3.387), ranking *WJG* as 35th among 80 journals in gastroenterology and hepatology (quartile in category Q2).

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Shu-Yu Yin*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Rao-Yu Ma*
Proofing Editorial Office Director: *Ze-Mao Gong*

NAME OF JOURNAL
World Journal of Gastroenterology

ISSN
ISSN 1007-9327 (print)
ISSN 2219-2840 (online)

LAUNCH DATE
October 1, 1995

FREQUENCY
Weekly

EDITORS-IN-CHIEF
Andrzej S Tarnawski, MD, PhD, DSc (Med),
Professor of Medicine, Chief Gastroenterology, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach, CA 90822, United States

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE
Ze-Mao Gong, Director
World Journal of Gastroenterology
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
November 28, 2018

COPYRIGHT
© 2018 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.f6publishing.com>

Basic Study

Different distributions of interstitial cells of Cajal and platelet-derived growth factor receptor- α positive cells in colonic smooth muscle cell/interstitial cell of Cajal/platelet-derived growth factor receptor- α positive cell syncytium in mice

Chen Lu, Xu Huang, Hong-Li Lu, Shao-Hua Liu, Jing-Yu Zang, Yu-Jia Li, Jie Chen, Wen-Xie Xu

Chen Lu, Xu Huang, Hong-Li Lu, Yu-Jia Li, Wen-Xie Xu, Department of Physiology, Shanghai Jiao Tong University School of Medicine, Shanghai 200240, China

Shao-Hua Liu, Department of Anesthesiology, Ren Ji Hospital Affiliated to Shanghai Jiao Tong University School of Medicine, Shanghai 201112, China

Jing-Yu Zang, Jie Chen, Department of Pediatric Surgery, Xin Hua Hospital Affiliated to Shanghai Jiao Tong University School of Medicine, Shanghai 200092, China

ORCID number: Chen Lu (0000-0001-8020-0501); Xu Huang (0000-0002-8640-5177); Hong-Li Lu (0000-0001-5374-8336); Shao-Hua Liu (0000-0001-8359-3758); Jing-Yu Zang (0000-0002-6646-9341); Yu-Jia Li (0000-0001-8614-2234); Jie Chen (0000-0003-2308-0130); Wen-Xie Xu (0000-0003-3040-4340).

Author contributions: Lu C designed and performed the majority of experiments; Huang X, Lu HL, and Liu SH contributed new reagents and analytic tools; Zang JY, Li YJ, and Chen J analyzed the data; Xu WX wrote and edited the manuscript.

Supported by The National Natural Science Foundation of China, No. 31671192 and No. 31571180; and Foundation of Xin Hua Hospital, No. JZPI201708.

Institutional review board statement: This research was approved by the Ethics Committee of Shanghai Jiao Tong University School of Medicine.

Institutional animal care and use committee statement: The protocol was approved by the Committee on the Ethics of Animal Experiments of Shanghai Jiao Tong University School of Medicine (Permit Number: Hu 686-2009).

Conflict-of-interest statement: To the best of our knowledge, no conflict of interest exists.

Data sharing statement: No additional data is available.

ARRIVE guidelines statement: The authors have read the ARRIVE guidelines and prepared the manuscript accordingly.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Corresponding author to: Wen-Xie Xu, PhD, Professor, Department of Physiology, Shanghai Jiao Tong University School of Medicine, 800 Dongchuan Road, Shanghai 200240, China. wenxiexu@sjtu.edu.cn
Telephone: +86-21-34205639
Fax: +86-21-34205639

Received: August 24, 2018

Peer-review started: August 24, 2018

First decision: October 5, 2018

Revised: November 7, 2018

Accepted: November 7, 2018

Article in press: November 7, 2018

Published online: November 28, 2018

Abstract

AIM

To investigate the distribution and function of interstitial

cells of Cajal (ICCs) and platelet-derived growth factor receptor- α positive (PDGFR α^+) cells in the proximal and distal colon.

METHODS

The comparison of colonic transit in the proximal and distal ends was performed by colonic migrating motor complexes (CMMCs). The tension of the colonic smooth muscle was examined by smooth muscle spontaneous contractile experiments with both ends of the smooth muscle strip tied with a silk thread. Intracellular recordings were used to assess electrical field stimulation (EFS)-induced inhibitory junction potentials (IJP) on the colonic smooth muscle. Western blot analysis was used to examine the expression levels of ICCs and PDGFR α in the colonic smooth muscle.

RESULTS

Treatment with NG-nitro-L-arginine methyl ester hydrochloride (L-NAME) significantly increased the CMMC frequency and spontaneous contractions, especially in the proximal colon, while treatment with MRS2500 increased only distal CMMC activity and smooth muscle contractions. Both CMMCs and spontaneous contractions were markedly inhibited by NPPB, especially in the proximal colon. Accordingly, CyPPA sharply inhibited the distal contraction of both CMMCs and spontaneous contractions. Additionally, the amplitude of stimulation-induced nitric oxide (NO)/ICC-dependent slow IJPs (sIJPs) by intracellular recordings from the smooth muscles in the proximal colon was larger than that in the distal colon, while the amplitude of electric field stimulation-induced purinergic/PDGFR α -dependent fast IJPs (fIJPs) in the distal colon was larger than that in the proximal colon. Consistently, protein expression levels of c-Kit and anoctamin-1 (ANO1) in the proximal colon were much higher, while protein expression levels of PDGFR α and small conductance calcium-activated potassium channel 3 (SK3) in the distal colon were much higher.

CONCLUSION

The ICCs are mainly distributed in the proximal colon and there are more PDGFR α^+ cells in the distal colon, which generates a pressure gradient between the two ends of the colon to propel the feces to the anus.

Key words: Interstitial cells of Cajal; Platelet-derived growth factor receptor- α positive cells; Smooth muscle cell/interstitial cell of Cajal/platelet-derived growth factor receptor- α positive cell syncytium; Nitric oxide; Purine

© The Author(s) 2018. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: The different distributions of interstitial cells of Cajal (ICCs) and platelet-derived growth factor receptor- α positive (PDGFR α^+) cells in the different parts of the colon result in a pressure gradient in the colon that propels the feces to the anus. Nitric oxide (NO) is mainly involved in the regulation of contraction through ICCs, and purine mainly participates in relaxation through PDGFR α^+ cells.

The regulation of the inhibitory transmitter NO and purine through the two kinds of interstitial cells (ICCs and PDGFR α^+ cells) is very important for rhythmic colonic migrating motor complexes, which are the main driving force underlying colon transit.

Lu C, Huang X, Lu HL, Liu SH, Zang JY, Li YJ, Chen J, Xu WX. Different distributions of interstitial cells of Cajal and platelet-derived growth factor receptor- α positive cells in colonic smooth muscle cell/interstitial cell of Cajal/platelet-derived growth factor receptor- α positive cell syncytium in mice. *World J Gastroenterol* 2018; 24(44): 4989-5004 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v24/i44/4989.htm> DOI: <http://dx.doi.org/10.3748/wjg.v24.i44.4989>

INTRODUCTION

In the gastrointestinal (GI) tract, smooth muscle cell/interstitial cell of Cajal/platelet-derived growth factor receptor- α positive cell (SIP) syncytium consists of interstitial cells of Cajal (ICCs), platelet-derived growth factor receptor- α positive cells (PDGFR α^+ cells), and smooth muscle cells (SMCs). ICCs and PDGFR α^+ cells are involved in the smooth muscle contraction. Remodeling or damaging of these cells can result in a variety of motor disorders^[1]. Cells in the SIP syncytium express many kinds of receptors and ion channels, and conductance changes in any type of cell can induce the excitability or relaxation of the smooth muscle. Although there have been several studies of ICCs and PDGFR α^+ cells in the GI tract regarding their location, morphology, function and more, most studies have focused on the small intestine and stomach, with few concerning the colon^[2,3]. Therefore, in the present study, we explored the distributions of the two kinds of interstitial cells along with their ion channels and transmission mechanisms for inducing contraction and diastolic reactions at both ends of the colon.

ICCs were first suggested to be GI pacemaker cells in the late 1970s based on electrophysiological and ultrastructural observations^[4]. Anoctamin-1 (ANO1), a very important functional protein in ICCs, is a calcium-activated chloride channel that produces pacemaker currents^[5]. Another interstitial cell type, PDGFR α^+ cells, referred to as "fibroblast-like" cells, express specific small conductance calcium-activated potassium channel 3 (SK3)^[6]. When the SK3 channel is activated, large amounts of K⁺ flow out, causing hyperpolarization and subsequent relaxation of PDGFR α^+ cells and downstream SMCs^[7]. Consequently, the motility of smooth muscles depends on the balance between excitatory regulation from ICCs and inhibitory regulation from PDGFR α^+ cells.

Colonic motility requires the coordination of the enteric nervous system (ENS) and SIP syncytium. The ENS, consisting of excitatory motor neurons and inhibitory motor neurons (IMNs), including nitric oxide synthase (NOS) and purine neurons, is involved in the regulation of

colonic movement^[8-10]. Mañé *et al.*^[11] have reported that nitric oxide (NO) is responsible for continuous smooth muscle relaxation, while purine neurons are responsible for transient relaxation, which may be dominant in colonic propulsion. In addition, colonic migrating motor complexes (CMMCs) known as the main form of the colonic transmission, have been demonstrated that their generation require two conditions: the activation of cholinergic motor neurons and the release of inhibitory neurotransmitters (mainly NO and purines), both of which can act on the SIP syncytium to regulate colonic transmission^[12].

It has been reported that electrical field stimulation (EFS) can induce changes in the membrane potential mediated by inhibitory neurotransmitters called the inhibitory junction potential (IJP), and subsequent relaxation reaction^[13,14]. IJPs are composed of two components: a fast-transient hyperpolarization (fIJP) and a subsequent slow and sustained hyperpolarization (sIJP)^[6,12]. The release of NO is closely related to ICCs, which can influence the generation of spontaneous contractions and the initiation of sIJPs through ICCs^[15-17]. On the other hand, fIJPs have been shown to be mediated by purines specifically through the P2Y1 receptor on PDGFR α ⁺ cells^[7,14].

Based on these previous reports, we sought to characterize the mechanism of the ENS and SIP in the regulation of colon motility, especially distributions of ICCs and PDGFR α ⁺ cells in the proximal and distal colon according to the response of CMMC, smooth muscle contraction, and intracellular recoding for the blockers of neurotransmitters.

MATERIALS AND METHODS

Ethical statement

All animals were obtained from the Experimental Animal Center of Shanghai Jiao Tong University School of Medicine. This research rigorously complied with the rules of the Guide for the Care and Use of Laboratory Animals of the Science and Technology Commission of China (STCC Publication No. 2, revised 1988). The protocol was approved by the Committee on the Ethics of Animal Experiments of Shanghai Jiao Tong University School of Medicine (Permit Number: Hu 686-2009). All operations were performed under anesthesia using isopentane, and every manipulation of the experimental animals was performed while simultaneously attempting to maximally relieve any suffering.

Animals

Adult male ICR mice were obtained from the Experimental Animal Center of Shanghai Jiao Tong University School of Medicine. The mice, aged 35 d and weighing approximately 30 g, were housed at 22 °C under a 12 h light/dark cycle with free access to water and food.

CMMC experiments

Mice were sacrificed under general anesthesia induced

by inhalant isoflurane overdose followed by cervical dislocation. Then, the abdomen was opened along the ventral midline, and the colon was exposed, removed quickly, and placed into Krebs solution continuously bubbling with a carbonated mixture (5% CO₂ and 95% O₂). The Krebs solution contained the following components (all concentrations in mmol/L): NaCl, 121.9; NaHCO₃, 15.5; KCl, 5.9; MgSO₄, 1.2; KH₂PO₄, 1.2; glucose, 11.5; and CaCl₂, 2.4. The entire colon was fixed in a Sylgard base dish with impalpable steel pins. The mesentery was carefully removed along the boundary line of the enterocoel under a dissecting microscope. All fecal pellets in the colon were artificially expelled with a 1-mL injector; this procedure was repeated to expel every pellet. This procedure must be performed with care to minimize intestinal damage. The empty colon was gently washed with 5 mL of warm Krebs solution, and a glass capillary tube was inserted through the lumen and linked to an artificial fecal pellet. The capillary was attached to the floor of the silica gel plate using U-shaped pins at the oral and anal end. A rectangular organ filled with 20 mL of warm Krebs solution (36.5 ± 0.5 °C) was also constantly inundated with carbon-oxygen gas. Then, the colon specimen was gently perfused with warm Krebs solution and left to stabilize for 30-40 min to secure the recovery of the colonic contraction activity. A silk thread (USP 5/0) was attached to both ends of the colon. The mechanical activity of the CMMCs was recorded using an isometric force transducer (RM6240C, Chengdu Instrument Factory, China) linked to an amplifier device. A tension of 0.1 g was applied to the empty colon, which was equilibrated for at least 40 min before the addition of the experimental drugs.

Preparation of smooth muscle tissue and isometric tension measurement

The entire colon full of fecal pellets was isolated as described above. The colon was cut along the mesentery, which is on the colonic circular axis; the pellets were flushed out with Krebs solution; the colon was pinned to a Sylgard dish with the mucosa facing upwards; and the mucosa and submucosa were removed carefully under a dissecting microscope. Smooth muscle strips (approximately 2 mm × 8 mm) were obtained by cutting along the circular axis from the fresh smooth muscle tissue. A silk thread (USP 5/0) was attached to both ends of the muscle strips and attached along the circular axis into 10 mL organ baths containing warm (37 °C) Krebs solution filled with 95% O₂ and 5% CO₂. The recording device was the same as that for the CMMCs above. A tension of 0.3 g was applied to the muscle strip, and it was equilibrated for at least 40 min before the recovery of its contraction activity.

Western blot analysis

Protein samples were extracted from the colonic smooth muscle tissues and lysed in radioimmunoprecipitation assay (RIPA) buffer (1:10; P0013, Beyotime Chemical Co., Jiangsu, China) and PMSF (1:100) solution. The

suspended material was centrifuged at 12000 rpm for 15 min at 4 °C, mixed with 4 × loading buffer, and then boiled for 5 min in a 100 °C water bath. The protein concentration of the supernatant was calculated using the bicinchoninic acid (BCA) protein assay method (P0010, Beyotime Chemical Co., Jiangsu, China). Protein (30 µg/lane) was subjected to 7.5% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and then transferred from the polyacrylamide gel to polyvinylidene difluoride (PVDF) membranes using an E-Blotter unit (Bio-Rad) for 120 min. Then, the PVDF membranes were blocked in 5% nonfat milk for 2 h and incubated with rabbit c-Kit and PDGFR- α monoclonal antibodies (1:1000; #3074, #3174, Cell Signaling Technology, United States) and with mouse anti-tubulin antibody (1:1000; AT819, Beyotime Chemical Co., Jiangsu, China) overnight at 4 °C. The blots were then washed five times (5 min per wash) with Tris-HCl-buffered saline including 0.1% Tween-20 (TBST) and then incubated with secondary antibodies, including either anti-rabbit IgG HRP-linked antibody (1:1000; 7074; Cell Signaling Technology) or anti-mouse IgG HRP-linked antibody (1:1000; 7076; Cell Signaling Technology) for 2 h at room temperature. Protein signal detection was performed using an enhanced chemiluminescence agent (ECL reagents). The signals of the blots were analyzed using Quantity One software.

Intracellular microelectrode recording and electrical stimulation

Based on the operation above, smooth muscle tissue (approximately 20 mm × 8 mm) was isolated from the empty colon, fixed facing forward and up onto the base of a Sylgard-covered chamber, and constantly filled with warm (37 °C) Krebs solution, 95% O₂, and 5% CO₂. The tissue was equilibrated for approximately 2 h before the recording started. The muscle tissue was maintained at 37 ± 0.5 °C through continuous perfusion with warm Krebs solution. Experimental procedures were carried out in the presence of nifedipine (1 µmol/L) to minimize the muscle contraction and maintain the cellular implements. Circular muscle cells were inserted using glass microelectrodes (80–100 M Ω) filled with KCl. Membrane potential was recorded with a Duo 773 (WPI Inc., Sarasota, FL, United States). EFS was made under a consistent voltage, pulse width, and duration (50 V, 0.5 ms, and 5 s, respectively) by two parallel platinum electrodes using a square-wave stimulator (YC-2 stimulator; Chengdu, China). The sIJs and fIJs of the smooth muscle were recorded in the presence and absence of various drugs, such as receptor antagonists or agonists in Krebs solution.

Drugs

Tetrodotoxin (TTX) was obtained from Absin Biochemical Company. Atropine was purchased from Sigma-Aldrich. Apamin, NG-Nitro-L-arginine methyl ester hydrochloride (L-NAME), NPPB, MRS2500, and CyPPA were obtained from Tocris Bioscience (Ellisville, MO, United States).

Statistical analysis

The data are described as the mean ± SE. The analysis of data differences between groups was performed using one-way analysis of variance (ANOVA), followed by the Bonferroni's post-hoc test or using the Student's unpaired *t*-test when needed. *P*-values less than 0.05 were considered to represent significant differences between groups, and *n*-values correspond to the number of animals that were used in the indicated experiments.

RESULTS

Regulatory effects of the ENS/Ach/NO for ANO1 channels in ICCs on CMMCs and spontaneous contractions

Effects of TTX and atropine on CMMCs and spontaneous contractions: To demonstrate the role of the ENS in colonic transit, we used a Na⁺ channel antagonist, TTX (0.4 µmol/L), to block neurons. TTX abolished the large amplitude contractions of CMMCs, and only the fast oscillating contractions remained. The contractions were decreased from 100% in controls to 30% ± 2.6% in the proximal colon and 21% ± 4.7% in the distal part (^a*P* < 0.05; *n* = 6, meaning the number of animals used and the same below; Figure 1A and C). Interestingly, the basal tone and frequencies of the burst-like contractions in both ends of the colon were markedly increased by TTX, indicating that the generation of CMMCs in the colon was significantly mediated by the ENS, while the increased basal tone and frequencies of the remaining contractions in the presence of TTX are likely generated directly by ICCs because the dominant effects of the ENS on ICCs were limited by TTX. In addition, the effects of TTX were significantly different between the proximal and distal colon. To further confirm the effects of TTX on the colonic contractions, we observed that TTX significantly increased the rate of spontaneous contractions from 100% in controls to 149% ± 4.2% in the proximal colon and 124% ± 5.4% in the distal part (^a*P* < 0.05; *n* = 7; Figure 1B and D). Moreover, there were significant differences in proximal and distal colon with TTX administration, respectively (^c*P* < 0.05; Figure 1A–D).

To explain the differences of TTX above, we further compared the roles of the excitatory neurons and inhibitory neurons in regulating colonic motility. First, atropine was employed to block acetylcholine (Ach), released by cholinergic neurons. We found that, similar to the effects of TTX, atropine (100 µmol/L) abolished the large amplitude contractions of CMMCs, and only the burst-like contractions (e.g., ICC pacemaking activity) remained in both parts of the colon. Overall, CMMCs were significantly suppressed, from 100% in controls to 21% ± 1.9% in the proximal colon and 39% ± 3.7% in the distal part (^a*P* < 0.05; *n* = 5; Figure 1E and G). Similarly, spontaneous contractions were also partially decreased by atropine treatment, from 100% in controls to 57% ± 1.8% in the proximal colon and 74% ± 3.9% in the distal part (^a*P* < 0.05; *n* = 7; Figure 1F and H). Interestingly, significant




Figure 1 Effects of tetrodotoxin and atropine on smooth muscle contractions of the colon. A and C: Inhibitory effects of tetrodotoxin (TTX) treatment on colonic migrating motor complexes (CMMCs) in the proximal and distal colon, and summary data of the area under the curve (AUC) at 300 s. It is worth noting that AUC measurements can demonstrate the sum of both the amplitudes and frequencies. The data are normalized to the control value (before the application of TTX) ($n = 6$; $^aP < 0.05$ vs control; $^cP < 0.05$ vs proximal colon). B and D: Enhanced effects of TTX on spontaneous contractions in the proximal and distal colon of mice and summary of the contractile responses to TTX, as indicated by the AUC at 200 s, in the smooth muscle tissues of the proximal and distal colon of mice. The data are normalized to the control value (before the application of TTX) ($n = 7$; $^aP < 0.05$ vs control; $^cP < 0.05$ vs proximal colon). E and G: Responses of CMMCs to atropine in the proximal and distal colon and summary of data as indicated by the AUC at 300 s. The data are normalized to the control value (before the application of atropine) ($n = 5$; $^aP < 0.05$ vs control; $^cP < 0.05$ vs proximal colon). F and H: Inhibitory effects of atropine on spontaneous contraction in the proximal and distal colon and summary of data showing contractile responses to atropine as indicated by the AUC at 200 s. The data are normalized to the control value (before the application of atropine) ($n = 7$; $^aP < 0.05$ vs control; $^cP < 0.05$ vs proximal colon). TTX: Tetrodotoxin; AUC: Area under the curve.

differences remained between the proximal and distal ends treated with atropine in the CMMC and muscle strip experiments ($^aP < 0.05$; Figure 1E-H). These results suggest that Ach may play a regulatory role for ICCs.

Effects of L-NAME and NPPB on CMMCs and spontaneous contractions: To characterize the effects of NO on colonic motility, L-NAME (100 $\mu\text{mol/L}$), a nonspecific inhibitor of NOS, was employed in the present study. We found that CMMCs, especially their frequency, were significantly enhanced by L-NAME, from 100% in controls to $172\% \pm 4.6\%$ in the proximal colon and $145\% \pm 4.4\%$ in the distal part ($^aP < 0.05$; $n = 7$; Figure 2A and C). Similarly, spontaneous contractions became larger and faster after administration with L-NAME, from 100% in controls to $170\% \pm 11.5\%$ in the proximal colon and $150\% \pm 4.0\%$ in the distal part ($^aP < 0.05$; $n = 7$; Figure 2E-H). Importantly, we also found that the effects of L-NAME on both CMMCs and spontaneous contractions were more significant in the proximal colon than in the distal part, respectively ($^cP < 0.05$; Figure 2E-H). Based on these results, we confirmed that NO contributes substantially to the regulation of CMMC frequency and suggest that there may be more ICCs in the proximal colon than in the distal part.

To further confirm that ICCs have distinct distributions between different parts of the colon, NPPB, a blocker of ANO1 channels, was applied. NPPB treatment (3 $\mu\text{mol/L}$) significantly inhibited CMMCs, from 100% in controls to $22\% \pm 2.7\%$ in the proximal colon and $46 \pm 4.3\%$ in the distal part ($^aP < 0.05$; $n = 5$; Figure 2E and G). The effects of NPPB were much more significant in the proximal part than in the distal part ($^cP < 0.05$; $n = 5$; Figure 2E and G). Furthermore, spontaneous contractions were also decreased by NPPB treatment, from 100% in controls to $19\% \pm 3.4\%$ in the proximal colon and $48\% \pm 3.7\%$ in the distal part ($^aP < 0.05$; $n = 5$; Figure 2F and H). However, the inhibitory effect of NPPB was much more significant in the proximal part than in the distal part of the colon ($^cP < 0.05$; $n = 5$; Figure 2F and H).

Regulatory effects of purine for SK3 channels in PDGFR α^+ cells on CMMCs and spontaneous contractions

Effects of MRS-2500 treatment on CMMCs and spontaneous contractions: In subsequent experiments, to further investigate the role of PDGFR α^+ cells in the regulation of colonic motility. First, MRS2500 was used (1 $\mu\text{mol/L}$; an antagonist of P2Y1) on CMMCs and spontaneous contractions. We found that in the proximal colon, MRS2500 had no significant effect on either CMMCs or spontaneous contractions (Figure 3A and B). However, in the distal colon, both CMMCs and spontaneous contractions were increased by MRS2500, reaching $143\% \pm 2.4\%$ for CMMCs and $130\% \pm 4.2\%$ for spontaneous contractions ($^aP < 0.05$; $n = 8$; Figure 3A-D). MRS2500 had a markedly stronger effect in the distal colon compared to the proximal part ($^cP < 0.05$;

$n = 8$; Figure 3A-D), which indicates that there is an obvious difference between the two ends. Similarly, this effect may result from different distributions of PDGFR α^+ cells in these colonic regions.

Effects of apamin and CyPPA on CMMCs and spontaneous contractions: To test the above hypothesis, we determined the effects of apamin (300 nmol/L; an SK3 channel antagonist) on colonic motility. We can observe that apamin significantly enhanced CMMCs, from 100% in controls to $126\% \pm 0.4\%$ in the proximal colon and $148\% \pm 2.1\%$ in the distal part ($^aP < 0.05$; $n = 7$; Figure 4A and C). Similar results were obtained for spontaneous contractions, which were increased from 100% in controls to $127\% \pm 1.6\%$ in the proximal colon and $159\% \pm 3.4\%$ in the distal part ($^aP < 0.05$; $n = 7$; Figure 4B and D). Similar to MRS2500 treatment, the effects of apamin in the distal colon were stronger than those in the proximal part ($^cP < 0.05$; $n = 7$; Figure 4A-D).

Subsequently, CyPPA (3 $\mu\text{mol/L}$), an SK3 channel agonist, was also used in the present study. We found that the effects of CyPPA were completely opposite of those observed with apamin treatment. CyPPA markedly inhibited CMMCs, from 100% in controls to $39\% \pm 1.7\%$ in the proximal colon and $25\% \pm 2.2\%$ in the distal part ($^aP < 0.05$; $n = 7$; Figure 4E and G). Clearly, the effects of CyPPA were much more significant in the distal part than in the proximal part of the colon ($^cP < 0.05$; $n = 7$; Figure 4E and G). Moreover, spontaneous contractions were also inhibited by CyPPA, from 100% in controls to $54\% \pm 4.1\%$ in the proximal colon and $19\% \pm 3.6\%$ in the distal part ($^aP < 0.05$; $n = 7$; Figure 4F and H). Therefore, CyPPA significantly inhibited both CMMCs and spontaneous contractions in both ends. Moreover, the effects of this drug in the distal colon were markedly stronger than those in the proximal part ($^cP < 0.05$; $n = 8$; Figure 4E-H).

Effects of ICC/ANO1 and PDGFR α^+ /SK3 on colonic membrane potentials of smooth muscle tissue

Effects of ANO1 and SK3 antagonists on resting membrane potentials of colonic smooth muscle tissues: It is well-established that smooth muscle contraction and relaxation result from membrane depolarization and hyperpolarization^[18]. To further evaluate the distributions of the two interstitial cell type, we first detected the average resting membrane potential (RMP) of colonic circular muscle cells in the proximal colon, which was 42.2 ± 1.92 mV, less than that in the distal part, which was 53.2 ± 2.86 mV ($^aP < 0.05$; $n = 12$; Figure 5A). Furthermore, NPPB induced membrane hyperpolarization in both parts of the colon, showing a greater extent in the proximal colon than in the distal part (9.17 ± 1.19 mV in the proximal colon and 4.67 ± 0.56 mV in the distal colon, $^aP < 0.05$; $n = 6$, Figure 5B and D). Nevertheless, apamin induced membrane depolarization, which was more marked in the distal




Figure 2 Effects of L-NAME and NPPB on smooth muscle contractions of the colon. A and C: Increased effects of L-NAME on colonic migrating motor complexes (CMMCs) in the proximal and distal colon and summary data of the area under the curve (AUC) at 300 s. The data are normalized to the control value (before the application of L-NAME) ($n = 7$; $^aP < 0.05$ vs control; $^cP < 0.05$ vs proximal colon). B and D: Enhanced effects of L-NAME on spontaneous contraction in the proximal and distal colon of mice and summary of the contractile responses to L-NAME as indicated by the AUC at 200 s. The data are normalized to the control value (before the application of L-NAME) ($n = 7$; $^aP < 0.05$ vs control; $^cP < 0.05$ vs proximal colon). E and G: Responses of CMMCs to NPPB in the proximal and distal colon and the summary of data as indicated by the AUC at 300 s. The data are normalized to the control value (before the application of NPPB) ($n = 5$; $^aP < 0.05$ vs control; $^cP < 0.05$ vs proximal colon). F and H: Inhibitory effects of NPPB (3 $\mu\text{mol/L}$) on spontaneous contraction in the proximal and distal colon and summary of the data showing contractile responses to NPPB as indicated by the AUC at 200 s. The data are normalized to the control value (before the application of NPPB) ($n = 5$; $^aP < 0.05$ vs control; $^cP < 0.05$ vs proximal colon). AUC: Area under the curve.




Figure 3 Effects of MRS2500 on smooth muscle contractions in the colon. A and C: Effects of MRS2500 on colonic migrating motor complexes (CMMCs) in the proximal and distal colon and summary of the area under the curve (AUC) at 300 s. The data are normalized to the control value (before the application of MRS2500) ($n = 8$; $^aP < 0.05$ vs control; $^cP < 0.05$ vs proximal colon). B and D: Representative traces illustrating the effects of the P2Y1 antagonist MRS2500 on the contractions of colonic smooth muscles in the proximal and distal colon of mice and summary of the contractile responses to MRS2500 as indicated by the AUC at 200 s. The data are normalized to the control value (before the application of MRS2500) ($n = 8$; $^aP < 0.05$ vs control; $^cP < 0.05$ vs proximal colon). AUC: Area under the curve.

colon than in the proximal part (7.10 ± 1.36 mV in the proximal colon and 11.07 ± 1.01 mV in the distal colon, $^aP < 0.05$; $n = 5$; Figure 5C and E).

Postjunctional potentials evoked by EFS in colonic smooth muscle tissues: Consequently, we used EFS to evaluate the response of membrane potentials to

determine whether there were differences between the proximal and distal colon. We observed that EFS (50 V; 3, 6, and 9 Hz; 5 s) induced IJPs including fIJPs and sIJPs in both ends of the colon. The average amplitude of the fIJPs was larger in the distal colon than in the proximal part (6.70 ± 0.39 , 11.8 ± 0.33 , and 18.86 ± 0.53 mV in the proximal colon, $n = 8$; 11.24 ± 0.35 , 14.58 ± 0.34 ,




Figure 4 Effects of small conductance calcium-activated potassium channel 3 channel antagonist and agonist treatment on smooth muscle contractions in the murine colon. A and C: Responses of colonic migrating motor complexes (CMMCs) to small conductance calcium-activated potassium channel 3 (SK3) channel antagonist treatment in the proximal and distal colon and summary of the CMMC responses to apamin as indicated by the area under the curve (AUC) at 300 s. The data are normalized to the control value (before the application of apamin) ($n = 7$; $^aP < 0.05$ vs control; $^cP < 0.05$ vs proximal colon). B and D: Enhanced effects of apamin (300 nmol/L) on spontaneous contractions in the proximal and distal colon and summary of the data showing contractile responses to apamin as indicated by the AUC at 200 s. The data are normalized to control (before the application of apamin) ($n = 7$; $^aP < 0.05$ vs control; $^cP < 0.05$ vs proximal colon). E and G: Responses of CMMCs to SK3 channel agonist treatment in the proximal and distal colon and summary of the CMMC responses to CyPPA as indicated by the AUC at 300 s. The data are normalized to the control value (before the application of CyPPA) ($n = 7$; $^aP < 0.05$ vs control; $^cP < 0.05$ vs proximal colon). F and H: Inhibitory effects of CyPPA on spontaneous contractions in the proximal and distal colon and summary of the data showing contractile responses to CyPPA as indicated by the AUC at 200 s. The data are normalized to the control value (before the application of CyPPA) ($n = 7$; $^aP < 0.05$ vs control; $^cP < 0.05$ vs proximal colon). AUC: Area under the curve.




Figure 5 Effects of anoctamin-1 and small conductance calcium-activated potassium channel 3 channel antagonist treatment on membrane potentials in the murine colon. **A:** Summary of the resting membrane potentials of colonic smooth muscle cells from the proximal and distal colon of mice ($n = 12$; $^aP < 0.05$). **B:** Responses of membrane potentials to anoctamin-1 (ANO1) antagonist treatment (NPPB, 3 $\mu\text{mol/L}$, arrow). **C:** Responses of membrane potentials to SK3 antagonist treatment (apamin, 300 nmol/L, arrow). **D** and **E:** Summary of data showing the average effects of NPPB and apamin on membrane potentials in colonic smooth muscle cells from both ends of the colon (NPPB $n = 6$; apamin $n = 5$; $^aP < 0.05$). RMP: Resting membrane potentials.

and 23.60 ± 0.45 mV in the distal colon, $n = 8$; $^aP < 0.05$, Figure 6A-C), while the amplitude of sIJs was more obvious in the proximal colon than in the distal parts (2.94 ± 0.17 , 7.8 ± 0.25 , and 3.60 ± 0.51 mV in the proximal colon, $n = 8$; 1.34 ± 0.24 , 5.14 ± 0.15 , and 9.98 ± 0.56 mV in the distal part, $n = 8$; $^aP < 0.05$, Figure 6A, B, and D). These results further confirm that there is indeed a

difference in the distribution of ICC and PDGFR α cells in the proximal and distal colon.

Changes in purine-dependent fIJs and NO-dependent sIJs in colonic smooth muscle tissues:

For the purpose of illustrating the distribution of PDGFR α cells at both ends of the colon, L-NAME (100 $\mu\text{mol/L}$) and




Figure 6 Membrane potentials evoked by electrical field stimulation in normal colonic proximal and distal muscles. A: Electrical field stimulation (EFS) from the proximal colon at different frequencies (50 V; 3, 6, and 9 Hz for 5 s; black bars in each panel) elicited biphasic hyperpolarization comprising a peak component followed by sustained hyperpolarization. B: EFS from the proximal colon at different frequencies (50 V; 3, 6, and 9 Hz for 5 s; black bars in each panel). C and D: Summary of data showing the average amplitude of fast inhibitory junction potentials and slow inhibitory junction potentials at both ends of the colon ($n = 8$; $^aP < 0.05$). fIJP: Fast inhibitory junction potential; sIJP: Slow inhibitory junction potential.

atropine (1 $\mu\text{mol/L}$) were used to block NO and Ach to solely observe the EFS response to purine. We found that the amplitude of the sIJP significantly attenuated and almost all that remains were the fIJP. The amplitudes of fIJP in the distal colonic muscles remained stronger than those in the proximal colon (*i.e.*, 6.78 ± 0.27 , 9.20 ± 0.49 , and 18.00 ± 0.71 mV; $^aP < 0.05$; $n = 7$; Figure 7A and C). In the distal colon, the mean amplitudes were separately 12.44 ± 0.47 , 15.80 ± 0.73 , and 24.22 ± 0.54 mV ($^aP < 0.05$; $n = 8$; Figure 7B and C). These results indicate that the distal colon has more PDGFR α^+ cells and stronger fIJP primarily elicited by purine and induces relaxation.

To explore the distribution of ICCs at both ends of the colon, MRS2500 (1 $\mu\text{mol/L}$) and atropine (1 $\mu\text{mol/L}$) were used to block purines and Ach to solely observe

the EFS response to NO. The addition of MRS2500 and atropine almost fully eliminated the fIJP, leaving only the sIJP. The amplitudes of sIJP (with EFS 3, 6, and 9 Hz) in the proximal colon appeared much larger than those in the distal colon (*i.e.*, 3.12 ± 0.33 , 5.54 ± 0.39 , and 10.18 ± 0.40 mV, Figure 7E and F); in the proximal colon, the mean amplitudes of sIJP were 5.54 ± 0.43 , 9.32 ± 0.53 , and 14.8 ± 0.37 mV, respectively ($^aP < 0.05$; $n = 7$; Figure 7D and F). These results confirm that the proximal colon has a greater distribution of ICCs and stronger sIJP elicited by NO, inducing stimulation by ICCs.

Expression levels of c-Kit and PDGFR α in colonic smooth muscle tissues

Finally, we examined the density of ICC, ANO1, PDGFR α ,




Figure 7 Fast inhibitory junction potential and slow inhibitory junction potential elicited by electrical field stimulation in smooth muscles of the proximal and distal colon. A and B: Atropine and L-NAME treatment blocked excitatory junction potentials (EJPs) and decreased slow inhibitory junction potentials (sIJP) but had no effect on fast IJPs (fIJPs) in colonic muscles from both ends. D and E: Atropine and MRS2500 treatment blocked EJPs and decreased fIJPs but had no effect on sIJPs in colonic muscles from both ends. C and F: Summary of data showing the average amplitude of fIJPs and sIJPs (in the presence of atropine and L-NAME, $n = 8$; in the presence of atropine and MRS2500, $n = 7$; $^*P < 0.05$). fIJP: Fast inhibitory junction potential; sIJP: Slow inhibitory junction potential.

and SK3 in the smooth muscle tissues of the colon. We found that the expression levels of c-Kit and ANO1 were much higher in the proximal colon ($97.7\% \pm 10.9\%$ and $35.7\% \pm 4.3\%$, respectively), while those in the distal end were lower ($58.2\% \pm 6.5\%$ and $19\% \pm 3.6\%$, respectively) [$^*P < 0.05$; $n = 8$; Figure 8A and C (c-Kit) and Figure 8B and D (ANO1)]. Additionally, the expression of the PDGFR α protein in the colonic muscle layer was significantly increased, from $54.8\% \pm 5.0\%$ to $77.6\% \pm 3.2\%$ in the proximal and distal ends, respectively; similarly, the expression of SK3 in the colonic muscle layer ranged from $36.2\% \pm 4.1\%$ to $61.4\% \pm 3.1\%$, respectively, in the proximal and distal ends [$^*P < 0.05$; $n = 8$; Figure 8E and F (PDGFR α) and Figure 8E and G (SK3)]. These protein expression data directly show that the proximal colon has more ICCs and that the distal colon has more PDGFR α^+ cells.

DISCUSSION

This study focused on the distribution and function of ICCs and PDGFR α^+ cells at both ends of the colon and the role of NO and purines on colonic dynamic transmission. This finding may have clinical implications in studying and treating colonic dysmotility of inflammatory bowel disease (IBD) and irritable bowel syndrome (IBS). It has previously been established that the large intestine has many different motor patterns, including segmental activity, antiperistaltic and peristaltic waves, and tonic inhibition^[19,20]. Regardless of transmission form, it is necessary to produce pressure gradient to ensure that feces are propelled from the proximal to distal colon. Therefore, we divided the colon into two segments to separately study its transmitting mechanism. Based on previous reports, the speed of fecal pellet propulsion




Figure 8 Expression levels of c-Kit/anoctamin-1 and platelet-derived growth factor receptor- α /small conductance calcium-activated potassium channel 3 in proximal and distal smooth muscles of the colon. A and B: Western blot analysis of interstitial cells of Cajal (ICCs) and anoctamin-1 (ANO1) in both ends of the colon. C and D: Summary of data analyzed using densitometric quantification (ICC, ANO1 $n = 8$, $^aP < 0.05$). E: Western blot analysis of platelet-derived growth factor receptor- α and small conductance calcium-activated potassium channel 3 in both ends of the colon. F and G: Summary of data analyzed using densitometric quantification ($n = 8$, $^aP < 0.05$). ANO1: Anoctamin-1; SK3: Small conductance calcium-activated potassium channel 3.

along the murine colon is the same as the conduction velocity of the CMMC contraction, indicating that CMMCs drive the pellets forward^[21].

In our study, we found that the large amplitude contraction of CMMCs was almost completely inhibited when the ENS or cholinergic neurons were blocked; instead,

irregular contractions with much higher frequencies were recorded. Notably, in the CMMC experiments (treatment with TTX), the distal colon had more obvious drug effect, while in smooth muscle contraction experiment, the drug effect of proximal colon was more prominent in the presence of TTX or atropine (Figure 1). These results suggested that even though CMMC was almost abolished by TTX, the spontaneous contractions of the smooth muscle were significantly increased by TTX treatment. Therefore, we speculate that in the ENS, inhibitory neurons play a dominant role in spontaneous contractions of the colon. Moreover, the small irregular contraction waves in atropine may result from the pacemaker activity of ICCs and CMMCs are initiated by excitatory cholinergic neurons through ICCs. In the past several years, pacemaker function of ICCs and the ENS network have generally been considered as separate systems in colonic transit. However, new findings have reported that the pacemaker network in the colon consists of both ICC-MY (interstitial cells of Cajal of myenteric plexus) and ICC-IM (intramuscular interstitial cells of Cajal) and that this network is activated *via* cholinergic KV7.5 channel inhibition in ICC-IM^[22]. Even though colonic motor transit has been confirmed by the cooperation of ICC networks with the ENS^[23], information regarding the transit mechanism by which ICCs sequentially function remain unclear. Therefore, we primarily focused on the regulation of murine colonic contractile activity by nitrergic and purinergic neurons.

Based on the finding of nitrergic signaling through ICCs to regulate colonic spontaneous contraction and CMMCs (described above), we used L-NAME and NPPB to block NO and ANO1 in ICCs, respectively, to observe whether there are functional differences between the two ends of the colon (Figure 2). Our results showed that the AUC (equaling the frequencies and amplitudes) of CMMCs and spontaneous contractions in the smooth muscle were significantly increased in the presence of L-NAME or NPPB, indicating that NO is involved in regulating the frequency of CMMC through ICCs to maintain normal colonic transit. Moreover, the proximal colon has larger amplitude and higher frequencies in the presence of these two drugs, indicating that there are more ICCs in this part of the colon.

In addition to NO, purine is also considered to be involved in inhibitory regulation of smooth muscle contractions through another kind of interstitial cells-PDGFR α ⁺ cells^[24]. Burnstock *et al*^[7] first proposed that adenosine triphosphate (ATP) or purine was a main inhibitory neurotransmitter in the GI tract in 1970. Additionally, it may promote the release of glutamate (an inhibitory neurotransmitter) by binding to presynaptic metabotropic P2Y receptors^[25]. Recent studies have proven that purinergic receptors (P2Y1) and SK3 channels are expressed in PDGFR α ⁺ cells responsible for nerve-mediated relaxation in the GI tract^[24]. Compared to other classical neurotransmitters, the role of purinergic signaling in colonic transmission is not well understood or appreciated. Therefore, our results (Figures 3 and 4)

proved that the amplitudes and frequencies of CMMCs and spontaneous contractions in the distal colon were markedly changed when P2Y1 receptors and SK3 channels were blocked by MRS2500 and apamin, respectively, or SK3 was activated by CyPPA, illustrating that purines help to maintain regular colonic transmission by inhibiting contractions. Moreover, the distal colon showing stronger contractile activity indicates that there may be more PDGFR α ⁺ cells, resulting from the inhibitory role of purine on PDGFR α ⁺ cells being eliminated by MRS2500 treatment.

Based on CMMC and smooth muscle contraction experiments, we speculated that ICCs and PDGFR α ⁺ cells may have different distributions at different ends of the colon. To confirm this hypothesis, we used intracellular recordings to detect whether there exists a difference in the electrophysiological properties between the two ends of the colon. Previous studies have shown that GI motility is regulated by excitatory and IMNs, among which IMNs evoke IJPs, causing muscle relaxation. In the colon, neurotransmitters released by the ENS could activate the conductance of interstitial cells (*i.e.*, ANO1 in ICCs or SK3 channels in PDGFR α ⁺ cells) to regulate the excitability of SMCs through gap junctions to mediate colonic motility^[23,26]. ANO1 regulates sensory signal transduction and smooth muscle contractile activity to generate rhythmic contraction, which is supported by the absence of slow waves in *ANO1* knockout mice^[27-29]. Additionally, SK3 channels in PDGFR α ⁺ cells are activated by purines and produce apamin-sensitive outward currents^[30]. These SK3 channels are believed to be involved in inducing membrane hyperpolarization and IJPs. In GI smooth muscle tissues, IJPs have bipolar phases and consist of two main components: fIJP, induced by the release of purines in PDGFR α ⁺ cells, and sIJP arising from NO through ICCs^[31,32], which was also proved in this study (Figures 5-7). More importantly, we also found that the amplitudes of fIJPs induced by purinergic neurons through PDGFR α ⁺ cells were much higher in the distal colon than in the proximal segment. Furthermore, the amplitude of sIJPs induced mainly by NO through ICCs was much higher in the proximal colon than in the distal part. These results provide concrete functional evidence that ICCs and PDGFR α ⁺ cells have a different distribution at both ends of the colon.

Finally, we confirmed that more ICCs were distributed in the proximal colon, whereas more PDGFR α ⁺ cells were distributed in the distal colon using Western blot analysis (Figure 8). We speculate that this distribution difference is responsible for the stronger contractions in the proximal colon and the relatively weaker contractions in the distal part, forming a pressure gradient from the proximal colon to the distal part to propel feces smoothly forward to the anus.

In summary, our findings in the present study provide clear evidence that the ENS regulates colonic motility through cholinergic and nitrergic neurons regulating the spontaneous rhythmic pacemaker activity of ICCs and through purinergic neurons mediating inhibitory ef-

fects on smooth muscle contractions by acting on P2Y1 receptors in PDGFR α ⁺ cells to suppress colonic transit. We further found that inhibitory neuromodulation has a leading role in colonic transmission, which may be associated with the formation of fecal pellets in the colon and full absorption of water and nutrients. Furthermore, the distribution characteristics of more ICCs in the proximal and more PDGFR α ⁺ cells in the distal colon contribute to the formation of a pressure gradient from the oral to anal ends of the colon, which is associated with the roles of ENS-NO/ICCs and purine/PDGFR α ⁺ cells in the regulation of colonic motility. Consequently, an in-depth understanding of the mechanism of colon motility may provide a new direction and target for the study and treatment of diarrhea or constipation in colon motility disorder, such as IBD and IBS.

ARTICLE HIGHLIGHTS

Research background

Interstitial cells of Cajal (ICCs) and PDGFR α ⁺ cells are mainly located in the smooth muscle layer of the colon, which is widely believed to play a critical role in the generation of colonic transit motility. Large numbers of studies have shown that ICCs are mainly responsible for the contraction response of the smooth muscle, while PDGFR α ⁺ cells mainly regulate the relaxation of the smooth muscle. Inhibitory neuronal transmitters [nitric oxide (NO) and purine] can respectively act on ICCs and PDGFR α ⁺ cells to mediate colonic transit, thus propelling feces into the rectum and the anus. However, the distributions of ICC and PDGFR α ⁺ in the colon remain unclear. Therefore, it is an important scientific problem to study the transit mechanism of the proximal and distal colon.

Research motivation

In recent years, the incidence of colonic dysmotility diseases has been increasing year by year, such as inflammatory bowel disease (IBD), irritable bowel syndrome (IBS), and complications of some diseases (e.g., diabetes-induced slow transit constipation), whose main clinical symptoms are diarrhea or constipation. The smooth muscle layer, as the main source of colon power, consists of ICCs, PDGFR α ⁺ cells, and smooth muscle cells (SMCs) contacted by the gap junction. Therefore, it is very significant to study the differential distributions of ICCs and PDGFR α ⁺ cells in the two ends of the colon.

Research objectives

In this study, we for the first time researched the distributions and functions of ICCs and PDGFR α ⁺ cells in the proximal and distal colon. Then, we studied the roles of inhibitory neurotransmitters NO for ICCs and purine for P2Y1 receptor on PDGFR α ⁺ cells in colonic motility transit. This study may represent a future strategy for therapeutic intervention in disorders of colonic motility, such as IBD and IBS, by understanding the distributions and function of NO-ICCs and purine-PDGFR α ⁺ cells at both ends of the colon.

Research methods

First, we compared the isolated colonic transit differences between the proximal and distal colon using colonic migrating motor complexes (CMMCs). Then, we used the smooth muscle contraction experiment, which is the major source of colon motility, to compare the drug differences between the proximal and distal colon by adding the blockers and agonists of anoctamin-1 (ANO1) channels on ICCs and small conductance calcium-activated potassium channel 3 (SK3) channels on PDGFR α ⁺ cells. Subsequently, we compared the membrane potentials of the proximal and distal colon by intracellular recordings. Later, Western blot analysis was used to detect the protein expression of c-Kit, ANO1, PDGFR α , and SK3 in the colon. Finally, we added immunofluorescence methods to visually describe the distributions of ICC and PDGFR α ⁺ cells in the

proximal and distal colon.

Research results

Treatment with tetrodotoxin (TTX) to block the enteric nervous system (ENS) in the CMMC experiment almost completely blocked colonic transit. However, in the smooth muscle contraction experiment, when the ENS was blocked, the contraction of the colon was enhanced, suggesting that inhibitory nerve regulation plays critical roles in the transmission of the colon. In addition, when the ANO1 channel on ICCs was blocked by NPPB, the proximal colon showed a more obvious inhibitory role. While the SK3 channels on PDGFR α ⁺ cells were blocked by apamin, there was a more obvious drug effect in the distal colon, indicating that the proximal colon might distribute more ICCs, and the distal colon has more PDGFR α ⁺ cells. Intracellular electrical recording experiments indicated that slow inhibitory junction potentials (sIJP) mediated by the NO-ICC-ANO1 signal pathway was more obvious in the proximal colon, while fast inhibitory junction potentials (fIJP) mediated by purine-PDGFR α ⁺-SK3 was more prominent in the distal colon, indicating that there are more ICCs in the proximal colon and more PDGFR α ⁺ cells in the distal colon from the membrane potential level.

Research conclusions

In this study, we demonstrated that NO has a more obvious effect on the ICCs in the proximal colon, while purine has a more prominent effect on the distal PDGFR α ⁺ cells, indicating that there are more ICC cells in the proximal colon and more PDGFR α ⁺ cells in the distal colon. In addition, NO and purine acting on the SMC/ICC/PDGFR α ⁺ cell (SIP) syncytium (consisting of SMCs, ICCs, and PDGFR α ⁺ cells) are both inhibitory neurotransmitters, suggesting that colonic transit is mainly dominated by inhibitory neuromodulation. Although the stomach and small intestine are dominated by excitatory neuromodulation, this feature of the colon may contribute to the adequate absorption of nutrients and the formation of feces.

Research perspectives

This is the first study to report that the differential distributions of ICCs and PDGFR α ⁺ cells at the two ends of the colon. Our findings have highlighted the effects of inhibitory neuromodulation NO-ICC-ANO1 and purine-PDGFR α ⁺ cells-SK3 on colonic transit. In order to maintain regular colonic transit, the perfect cooperation of NO-ICC and purine-PDGFR α ⁺ cells is required. Therefore, in the future study of colon dynamic disorder such as IBD or IBS, we can start from whether the distribution and function of ICC and PDGFR α ⁺ cells have changed, and then extend to the effect of the ENS, especially inhibitory neuromodulation, on colonic transmission disorder to find the pathogenesis of colonic transit dysfunction. These findings in the present study may provide new insights and strategies for the diagnosis and treatment of gastrointestinal motility disorders.

REFERENCES

- 1 Sanders KM, Ward SM, Koh SD. Interstitial cells: regulators of smooth muscle function. *Physiol Rev* 2014; **94**: 859-907 [PMID: 24987007 DOI: 10.1152/physrev.00037.2013]
- 2 Sanders KM, Koh SD, Ward SM. Interstitial cells of cajal as pacemakers in the gastrointestinal tract. *Annu Rev Physiol* 2006; **68**: 307-343 [PMID: 16460275 DOI: 10.1146/annurev.physiol.68.040504.094718]
- 3 Huizinga JD, Chen JH. Interstitial cells of Cajal: update on basic and clinical science. *Curr Gastroenterol Rep* 2014; **16**: 363 [PMID: 24408748 DOI: 10.1007/s11894-013-0363-z]
- 4 Faussone Pellegrini MS, Cortesini C, Romagnoli P. [Ultrastructure of the tunica muscularis of the cardiac portion of the human esophagus and stomach, with special reference to the so-called Cajal's interstitial cells]. *Arch Ital Anat Embriol* 1977; **82**: 157-177 [PMID: 613989]
- 5 Bernstein K, Vink JY, Fu XW, Wakita H, Danielsson J, Wapner R, Gallos G. Calcium-activated chloride channels anoctamin 1 and 2 promote murine uterine smooth muscle contractility. *Am J Obstet Gynecol* 2014; **211**: 688.e1-688.10 [PMID: 24928056 DOI: 10.1016/

- j.ajog.2014.06.018]
- 6 **Song NN**, Lu HL, Lu C, Tong L, Huang SQ, Huang X, Chen J, Kim YC, Xu WX. Diabetes-induced colonic slow transit mediated by the up-regulation of PDGFR α ⁺ cells/SK3 in streptozotocin-induced diabetic mice. *Neurogastroenterol Motil* 2018 Epub ahead of print [PMID: 29521017 DOI: 10.1111/nmo.13326]
- 7 **Jiménez M**. Platelet-derived growth factor receptor- α -positive cells: new players in nerve-mediated purinergic responses in the colon. *J Physiol* 2015; **593**: 1765-1766 [PMID: 25871557 DOI: 10.1113/JP270259]
- 8 **Qu ZD**, Thacker M, Castellucci P, Bagyánszki M, Epstein ML, Furness JB. Immunohistochemical analysis of neuron types in the mouse small intestine. *Cell Tissue Res* 2008; **334**: 147-161 [PMID: 18855018 DOI: 10.1007/s00441-008-0684-7]
- 9 **Foong JP**, Hirst CS, Hao MM, McKeown SJ, Boesmans W, Young HM, Bornstein JC, Vanden Berghe P. Changes in Nicotinic Neurotransmission during Enteric Nervous System Development. *J Neurosci* 2015; **35**: 7106-7115 [PMID: 25948261 DOI: 10.1523/JNEUROSCI.4175-14.2015]
- 10 **Lu HL**, Huang X, Wu YS, Zhang CM, Meng XM, Liu DH, Kim YC, Xu WX. Gastric nNOS reduction accompanied by natriuretic peptides signaling pathway upregulation in diabetic mice. *World J Gastroenterol* 2014; **20**: 4626-4635 [PMID: 24782615 DOI: 10.3748/wjg.v20.i16.4626]
- 11 **Mañé N**, Gil V, Martínez-Cutillas M, Clavé P, Gallego D, Jiménez M. Differential functional role of purinergic and nitrergic inhibitory cotransmitters in human colonic relaxation. *Acta Physiol (Oxf)* 2014; **212**: 293-305 [PMID: 25327170 DOI: 10.1111/apha.12408]
- 12 **Durnin L**, Lees A, Manzoor S, Sasse KC, Sanders KM, Mutafova-Yambolieva VN. Loss of nitric oxide-mediated inhibition of purine neurotransmitter release in the colon in the absence of interstitial cells of Cajal. *Am J Physiol Gastrointest Liver Physiol* 2017; **313**: G419-G433 [PMID: 28705804 DOI: 10.1152/ajpgi.00045.2017]
- 13 **Hibberd TJ**, Costa M, Travis L, Brookes SJH, Wattoo DA, Feng J, Hu H, Spencer NJ. Neurogenic and myogenic patterns of electrical activity in isolated intact mouse colon. *Neurogastroenterol Motil* 2017; **29**: 1-12 [PMID: 28418103 DOI: 10.1111/nmo.13089]
- 14 **Zhang Y**, Lomax AE, Paterson WG. P2Y1 receptors mediate apamin-sensitive and -insensitive inhibitory junction potentials in murine colonic circular smooth muscle. *J Pharmacol Exp Ther* 2010; **333**: 602-611 [PMID: 20103587 DOI: 10.1124/jpet.109.160978]
- 15 **Lies B**, Beck K, Keppler J, Saur D, Groneberg D, Friebe A. Nitrergic signalling via interstitial cells of Cajal regulates motor activity in murine colon. *J Physiol* 2015; **593**: 4589-4601 [PMID: 26227063 DOI: 10.1113/JP270511]
- 16 **Kaji N**, Horiguchi K, Iino S, Nakayama S, Ohwada T, Otani Y, Firman, Murata T, Sanders KM, Ozaki H, Hori M. Nitric oxide-induced oxidative stress impairs pacemaker function of murine interstitial cells of Cajal during inflammation. *Pharmacol Res* 2016; **111**: 838-848 [PMID: 27468647 DOI: 10.1016/j.phrs.2016.07.030]
- 17 **Smith TK**, Koh SD. A model of the enteric neural circuitry underlying the generation of rhythmic motor patterns in the colon: the role of serotonin. *Am J Physiol Gastrointest Liver Physiol* 2017; **312**: G1-G14 [PMID: 27789457 DOI: 10.1152/ajpgi.00337.2016]
- 18 **Blair PJ**, Rhee PL, Sanders KM, Ward SM. The significance of interstitial cells in neurogastroenterology. *J Neurogastroenterol Motil* 2014; **20**: 294-317 [PMID: 24948131 DOI: 10.5056/jnm14060]
- 19 **Costa M**, Dodds KN, Wiklendt L, Spencer NJ, Brookes SJ, Dinning PG. Neurogenic and myogenic motor activity in the colon of the guinea pig, mouse, rabbit, and rat. *Am J Physiol Gastrointest Liver Physiol* 2013; **305**: G749-G759 [PMID: 24052530 DOI: 10.1152/ajpgi.00227.2013]
- 20 **Costa M**, Wiklendt L, Simpson P, Spencer NJ, Brookes SJ, Dinning PG. Neuromechanical factors involved in the formation and propulsion of fecal pellets in the guinea-pig colon. *Neurogastroenterol Motil* 2015; **27**: 1466-1477 [PMID: 26251321 DOI: 10.1111/nmo.12646]
- 21 **Heredia DJ**, Dickson EJ, Bayguinov PO, Hennig GW, Smith TK. Localized release of serotonin (5-hydroxytryptamine) by a fecal pellet regulates migrating motor complexes in murine colon. *Gastroenterology* 2009; **136**: 1328-1338 [PMID: 19138686 DOI: 10.1053/j.gastro.2008.12.010]
- 22 **Wright GW**, Parsons SP, Loera-Valencia R, Wang XY, Barajas-López C, Huizinga JD. Cholinergic signalling-regulated KV7.5 currents are expressed in colonic ICC-IM but not ICC-MP. *Pflugers Arch* 2014; **466**: 1805-1818 [PMID: 24375291 DOI: 10.1007/s00424-013-1425-7]
- 23 **Huizinga JD**, Martz S, Gil V, Wang XY, Jimenez M, Parsons S. Two independent networks of interstitial cells of cajal work cooperatively with the enteric nervous system to create colonic motor patterns. *Front Neurosci* 2011; **5**: 93 [PMID: 21833164 DOI: 10.3389/fnins.2011.00093]
- 24 **Lee H**, Koh BH, Peri LE, Sanders KM, Koh SD. Purinergic inhibitory regulation of murine detrusor muscles mediated by PDGFR α ⁺ interstitial cells. *J Physiol* 2014; **592**: 1283-1293 [PMID: 24396055 DOI: 10.1113/jphysiol.2013.267989]
- 25 **Patti L**, Raiteri L, Grilli M, Parodi M, Raiteri M, Marchi M. P2X(7) receptors exert a permissive role on the activation of release-enhancing presynaptic α 7 nicotinic receptors co-existing on rat neocortex glutamatergic terminals. *Neuropharmacology* 2006; **50**: 705-713 [PMID: 16427662 DOI: 10.1016/j.neuropharm.2005.11.016]
- 26 **Furness JB**. The enteric nervous system and neurogastroenterology. *Nat Rev Gastroenterol Hepatol* 2012; **9**: 286-294 [PMID: 22392290 DOI: 10.1038/nrgastro.2012.32]
- 27 **Hwang SJ**, Blair PJ, Britton FC, O'Driscoll KE, Hennig G, Bayguinov YR, Rock JR, Harfe BD, Sanders KM, Ward SM. Expression of anoctamin 1/TMEM16A by interstitial cells of Cajal is fundamental for slow wave activity in gastrointestinal muscles. *J Physiol* 2009; **587**: 4887-4904 [PMID: 19687122 DOI: 10.1113/jphysiol.2009.176198]
- 28 **Cipriani G**, Serboiu CS, Gherghiceanu M, Fausone-Pellegrini MS, Vannucchi MG. NK receptors, Substance P, Anol expression and ultrastructural features of the muscle coat in Cav-1(-/-) mouse ileum. *J Cell Mol Med* 2011; **15**: 2411-2420 [PMID: 21535398 DOI: 10.1111/j.1582-4934.2011.01333.x]
- 29 **Huang F**, Rock JR, Harfe BD, Cheng T, Huang X, Jan YN, Jan LY. Studies on expression and function of the TMEM16A calcium-activated chloride channel. *Proc Natl Acad Sci USA* 2009; **106**: 21413-21418 [PMID: 19965375 DOI: 10.1073/pnas.0911935106]
- 30 **Iino S**, Nojyo Y. Immunohistochemical demonstration of c-Kit-negative fibroblast-like cells in murine gastrointestinal musculature. *Arch Histol Cytol* 2009; **72**: 107-115 [PMID: 20009347 DOI: 10.1679/aohc.72.107]
- 31 **Lang RJ**. Do 'fibroblast-like cells' intercede during enteric inhibitory motor neurotransmission in gastrointestinal smooth muscles? *J Physiol* 2011; **589**: 453-454 [PMID: 21285025 DOI: 10.1113/jphysiol.2010.204016]
- 32 **Yeoh JW**, Corrias A, Buist ML. A mechanistic model of a PDGFR α (+) cell. *J Theor Biol* 2016; **408**: 127-136 [PMID: 27521526 DOI: 10.1016/j.jtbi.2016.08.004]

P- Reviewer: Liu S, Rolle U S- Editor: Ma RY
L- Editor: Wang TQ E- Editor: Yin SY





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>



ISSN 1007-9327

