

找到约 121,000 条结果 (用时 0.68 秒)

Can Ultrasound of the Tibial Nerve Detect Diabetic Peripheral ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3507587/> - 翻译此页

作者: S Riaz - 2012 - 被引用次数: 62 - 相关文章

2012年11月14日 - A masked expert sonographer **measured** the **cross-sectional area** ... The mean CSA of the PTN in DSP compared **with** control **subjects** at distances of 1 (23.03 vs. ... age, height, or weight **between** the **presence** of **diabetic neuropathy** and **Sonographic** evaluation of the **median nerve** in **diabetic** patients: ...

Does Diabetes Mellitus Influence Carpal Tunnel Syndrome?

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5532320/> - 翻译此页

作者: YH Kim - 2017 - 被引用次数: 1 - 相关文章

2017年6月30日 - We investigated the **association between DM** and CTS using both ... US **cross-sectional area** (CSA) and NCS **measurements** at the median ... to differ between **DM** and non-DM patients regardless of the **presence** or absence of CTS. **Median nerve ultrasound** in **diabetic peripheral neuropathy** with and ...

PDF | Can Ultrasound of the Tibial Nerve Detect Diabetic Peripheral ...

https://www.researchgate.net/.../231864096_Can_Ultrasound_of_the_Tibial_... - 翻译此页

We aimed to examine the **association between** the size. ... Can **Ultrasound** of the Tibial **Nerve** Detect **Diabetic Peripheral Neuropathy**? ... A masked expert sonographer **measured** the **cross-sectional area** (CSA) of the PTN at 1, 3, ... The mean CSA of the PTN in DSP compared with control **subjects** at distances of 1 (23.03 vs.

Sonographic assessment of carpal tunnel syndrome in diabetic ...

https://journals.lww.com/.../Sonographic_assessment_of_carpal_tunnel_synd... - 翻译此页

作者: MA Kotb - 2018 - 相关文章

The study included 42 healthy **subjects**, 46 patients with idiopathic CTS, and 76 patients ... CTS was diagnosed in patients with **diabetic neuropathy** if they met the following ... The **cross-sectional area** of the **median nerve** was **measured** by ... et al, who reported no significant **correlation between BMI** and **median nerve CSA**.

Match Overview

1	Crossref 38 words Babalola Ishmael Afolabi, Bolanle Olubunmi Ibitoye, Rosemary Temidayo Ikem, Adeleye Dorcas Omisore et al.	1%
2	Crossref 34 words Tsuneo Watanabe, Hiroyasu Ito, Ayako Morita, Yuriko Uno et al. "Sonographic Evaluation of the Median Nerve in ..."	1%
3	Internet 20 words crawled on 14-Jul-2018 mdpi.com	1%
4	Crossref 19 words Kim, Li-Na, Hee-Kyu Kwon, Hyun-Im Moon, Sung-Bum Park, and Hang-Jae Lee. "Sonography of the Median Nerve ..."	<1%
5	Crossref 18 words Yüksel, Hasan, Ali Riza Odabaşı, Selda Demircan, Kutsi Köseoğlu, Kadir Kizilkaya, and Ergün Onur. "Effects of ..."	<1%
6	Internet 14 words crawled on 03-Aug-2017 medcraveonline.com	<1%

Name of Journal: World Journal of Diabetes

Manuscript type: Case-control study

Relationship between sonographically measured median nerve cross-sectional area and presence of peripheral neuropathy in diabetic subjects

Fredrick Andrew Attah, Christiana Mopelola Asaleye, Adeleye Dorcas Omisore, Babatope Ayodeji Kolawole, Adeniyi Sunday Aderibigbe, Mathew Alo

Abstract

BACKGROUND

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 256,000 条结果 (用时 0.67 秒)

[PDF] Comparison of Cross Sectional Areas of Median Nerve in Type 2 ...

<https://www.banglajol.info/index.php/BIRDEM/article/view/35039/23630> - 翻译此页

作者: MT Rahman - 相关文章

2017年10月31日 - Healthy Adult Subjects Measured by 2 D-B Mode High ... median nerve of diabetic subjects with peripheral neuropathy and control ... Conclusion: This study showed that there is a significant difference between the cross sectional areas of median ... used for diagnosis, high resolution diagnostic ultrasound.

缺少字词: presence

High resolution ultrasonography of the tibial nerve in diabetic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5769664/> - 翻译此页

作者: K Singh - 2017 - 被引用次数: 2 - 相关文章

2017年12月29日 - The cross sectional area and maximum thickness of nerve fascicles of the ... The diabetic patients with no signs of peripheral neuropathy had a larger ... of the tibial nerve (0.40 mm) than healthy non-diabetic subjects (12.42 +/- 1.01 ... the relationship between diabetic neuropathy and cross-sectional area of ...

Sonographic assessment of carpal tunnel syndrome in diabetic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6024071/> - 翻译此页

作者: MA Kotb - 2018 - 被引用次数: 1 - 相关文章

2018年6月15日 - The study included 42 healthy subjects, 46 patients with idiopathic CTS, and 76 ...

找到约 209,000 条结果 (用时 0.62 秒)

Sonographic Evaluation of the Peripheral Nerve in Diabetic Patients ...

https://www.researchgate.net/.../43351800_Sonographic_Evaluation_of_the... - 翻译此页

The **cross-sectional area** (CSA) and echo intensity of the peripheral **nerve** were **Correlation between** markers of peripheral **nerve** function and structure in type 1 ... to affect the **sonographic** visualization of **nerves** is the **presence** of **diabetes**. of **median** in **diabetic** patients with **peripheral neuropathy** and healthy **subjects** ...

Median nerve ultrasound in diabetic peripheral neuropathy with and ...

https://www.researchgate.net/.../235523578_Median_nerve_ultrasound_in_d... - 翻译此页

Introduction: **Median nerve ultrasound** shows increased **cross-sectional area** (CSA) in carpal tunnel syndrome (CTS) and **diabetic peripheral neuropathy** (PN). ... NCS were correlated **with** blinded **median nerve CSA ultrasound measurements**. ... to ulnar interosseous **difference**) were significantly prolonged in CTS cases.

High resolution ultrasonography of the tibial nerve in diabetic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5769664/> - 翻译此页

作者: K Singh - 2017 - 被引用次数: 2 - 相关文章

2017年12月29日 - The **cross sectional area** and maximum thickness of **nerve** fascicles of the ... The **diabetic** patients with no signs of **peripheral neuropathy** had a larger ... of the tibial **nerve** (0.40 mm) than healthy non-diabetic **subjects** (12.42 +/- 1.01 ... the **relationship between diabetic** neuropathy and **cross-sectional area** of ...

Can Ultrasound of the Tibial Nerve Detect Diabetic Peripheral ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3507587/> - 翻译此页

作者: S Riazi - 2012 - 被引用次数: 67 - 相关文章

We aimed to examine the **association between** the size of the posterior tibial nerve ... A masked expert sonographer **measured** the **cross-sectional area** (CSA) of the ... between the **presence** of **diabetic neuropathy** and a greater cross-sectional **Sonographic** evaluation of the **median nerve** in **diabetic** patients: comparison ...

Can Ultrasound of the Tibial Nerve Detect Diabetic Peripheral ...

care.diabetesjournals.org/content/diacare/35/12/2575.full.pdf - 翻译此页

作者: S Riazi - 2012 - 被引用次数: 67 - 相关文章

Diabetic Peripheral Neuropathy? ... We aimed to examine the **association between** the size of the posterior tibial **nerve** (PTN) and the **presence** and severity of DSP. ... consecutive **diabetic** patients classified by NCS as **subjects** with DSP or control ... **measured** the **cross-sectional area** (CSA) of the PTN at 1, 2, and 5 cm.