

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 41898

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Warm ischemia time and elevated serum uric acid are associated with metabolic syndrome after liver transplantation with donation after cardiac death

Hu LS *et al.* PTMS after DCD LT

Liang-Shuo Hu, Yi-Chao Chai, Jie Zheng, Jian-Hua Shi, Chun Zhang, Min Tian, Yi Lv, Bo Wang, Ai Jia

Abstract

AIM

To describe the prevalence of posttransplant metabolic syndrome (PTMS)

Match Overview

1	Internet 19 words crawled on 05-Sep-2016 tpp.uah.edu	1%
2	Crossref 15 words P. Ramirez, D. Ferreras, B. Febrero, M. Royo et al. "Outcome s of Liver Transplantation Using Older Donors After Circula' ...	<1%
3	Internet 15 words crawled on 24-Oct-2014 onlinelibrary.wiley.com	<1%
4	Internet 15 words crawled on 31-Oct-2017 repub.eur.nl	<1%
5	Crossref 13 words "Donor and recipient effects on graft and patient survival", Cl inical Liver Disease, 2013.	<1%

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 92,900 条结果 (用时 0.61 秒)

Donation after cardiac death liver transplantation: predictors of outcome.

<https://www.ncbi.nlm.nih.gov/pubmed/20977642> - 翻译此页

作者: AK Mathur - 2010 - 被引用次数: 141 - 相关文章

Donation after cardiac death liver transplantation: predictors of outcome. ... We aimed to identify recipient, donor and transplant risk factors associated with graft ... metabolic liver disorder, transplant MELD \geq 35, hospitalization at transplant and ... Donor warm ischemia time \geq 35 min significantly increased graft failure rates ...

缺少字词: serum uric acid syndrome

Ischemic cholangiopathy after controlled donation after cardiac death ...

<https://www.ncbi.nlm.nih.gov/pubmed/21245668> - 翻译此页

作者: CL Jay - 2011 - 被引用次数: 171 - 相关文章

Ischemic cholangiopathy after controlled donation after cardiac death liver ... (1)Comprehensive Transplant Center, Feinberg School of Medicine, Northwestern ... of biliary complications (95% CI= 1.8–3.4) and a 10.8 times increased odds of IC ...

缺少字词: warm serum uric acid metabolic syndrome

Successful donation after cardiac death liver transplants with ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/27783454> - 翻译此页

作者: R De Carlis - 2017 - 被引用次数: 15 - 相关文章

The role of donation after cardiac death (DCD) in expanding the donor pool is ... by the incidence of primary nonfunction (PNF) and ischemia-related complications. ... Acceptance of the graft was based on the trend of serum transaminase and ... Time Factors; Tissue Donors; Tissue and Organ Harvesting/ methods*; Warm ...

缺少字词: elevated uric acid metabolic

^[PDF] Extending Donor Pool with Donation after Cardiac Death in ... - ORBi

<https://orbi.uliege.be/bitstream/2268/139275/1/these%20orbi2013.pdf>

作者: H Le Dinh - 2012 - 被引用次数: 1 - 相关文章

(DCD), or donation after circulatory determination of death* (DCDD). 3 ... Warm ischemia time (WIT) is usually unpredictable and longest in categories 1 and 2, but shorter and ... min. CIT < 4-6 hr liver. < 18 hr kidney. Cardiac arrest. Transplantation. Time and lipids, HMGB1, uric acid, and calcium pyrophosphate crystals.

Hypothermic temperature effects on organ survival and restoration ...

<https://www.nature.com/scientific-reports/articles> - 翻译此页

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 101,000 条结果 (用时 0.56 秒)

The Impact of Combined Warm Ischemia Time on Development of ...

<https://www.ncbi.nlm.nih.gov/pubmed/29329188> - 翻译此页

作者: M Kalisvaart - 2018 - 被引用次数: 1 - 相关文章

The Impact of Combined **Warm Ischemia Time** on Development of Acute Kidney Injury in **Donation After Circulatory Death Liver Transplantation**: Stay Within ... when **high**-risk grafts, such as **donation after circulatory death** (DCD) grafts are used. ... Length of combined WIT is **associated** with severity of postoperative AKI and ...

缺少字词: serum uric acid metabolic syndrome

Donation after cardiac death liver transplantation: predictors of outcome.

<https://www.ncbi.nlm.nih.gov/pubmed/20977642> - 翻译此页

作者: AK Mathur - 2010 - 被引用次数: 143 - 相关文章

Donation after cardiac death liver transplantation: predictors of outcome. ... We aimed to identify recipient, **donor** and transplant risk factors **associated** with graft ... **metabolic** liver disorder, transplant MELD ≥ 35 , hospitalization at transplant and ... **Donor warm ischemia time** ≥ 35 min significantly **increased** graft failure rates ...

缺少字词: serum uric acid syndrome

Transplantation - clinical I | Nephrology Dialysis Transplantation ...

https://academic.oup.com/ndt/article/27/suppl_2/ii304/1935120 - 翻译此页

作者: M Bonani - 2012 - 被引用次数: 1 - 相关文章

2012年5月1日 - HR of **death following transplant** failure, by PRD combined **liver** and kidney **transplantation** and eculizumab have been It is hoped that the recent specific recommendations by the Organ **Donation** Taskforce focussing **upon** the had serious AEs (1 possibly drug-**related**: transient **ischemic** attack).

Cardiovascular Morbidity and Mortality After Liver Transplantation: The

...

<https://aasldpubs.onlinelibrary.wiley.com/doi/pdf/10.1002/lt.24738> - 翻译此页

作者: D D'Avola - 2017 - 被引用次数: 11 - 相关文章

Cardiovascular (CV) diseases are recognized longterm causes of **death after** ... diabetes, **metabolic**

找到约 162,000 条结果 (用时 0.54 秒)

The Impact of Combined Warm Ischemia Time on Development of ...

<https://www.ncbi.nlm.nih.gov/pubmed/29329188> - 翻译此页

作者: M Kalisvaart - 2018 - 被引用次数: 1 - 相关文章

The Impact of Combined **Warm Ischemia Time** on Development of Acute Kidney Injury in **Donation After Circulatory Death Liver Transplantation**: Stay Within ... when **high-risk** grafts, such as **donation after circulatory death** (DCD) grafts are used. ... Length of combined WIT is **associated** with severity of postoperative AKI and ...

缺少字词: `serum uric acid metabolic syndrome`

Donation after cardiac death liver transplantation: predictors of outcome.

<https://www.ncbi.nlm.nih.gov/pubmed/20977642> - 翻译此页

作者: AK Mathur - 2010 - 被引用次数: 143 - 相关文章

Donation after cardiac death liver transplantation: predictors of outcome. ... We aimed to identify recipient, **donor** and transplant risk factors **associated** with graft ... **metabolic** liver disorder, transplant MELD ≥ 35 , hospitalization at transplant and ... **Donor warm ischemia time** ≥ 35 min significantly **increased** graft failure rates ...

缺少字词: `serum uric acid syndrome`

Transplantation - clinical | Nephrology Dialysis Transplantation ...

https://academic.oup.com/ndt/article/27/suppl_2/ii304/1935120 - 翻译此页

作者: M Bonani - 2012 - 被引用次数: 1 - 相关文章

2012年5月1日 - HR of **death following transplant** failure, by PRD combined **liver** and kidney **transplantation** and eculizumab have been It is hoped that the recent specific recommendations by the Organ **Donation** Taskforce focussing **upon** the had serious AEs (1 possibly **drug-related**: transient **ischemic** attack).

Hypothermic temperature effects on organ survival and restoration ...

<https://www.nature.com/scientific-reports/articles> - 翻译此页

作者: J Ishikawa - 2015 - 被引用次数: 7 - 相关文章

2015年4月22日 - **Donor organ transplantation** using low-temperature storage is used as ... the organ **preservation time** by storing organs at sub-zero temperatures. ... criteria donors (ECDs) and **donation**