

World Journal of *Gastroenterology*

World J Gastroenterol 2018 December 28; 24(48): 5415-5536



**EDITORIAL**

- 5415 Role of cenicriviroc in the management of nonalcoholic fatty liver disease
Neokosmidis G, Tziomalos K

REVIEW

- 5418 Colorectal cancer vaccines: Tumor-associated antigens vs neoantigens
Wagner S, Mullins CS, Linnebacher M

MINIREVIEWS

- 5433 Checkpoint inhibitors: What gastroenterologists need to know
Ahmed M
- 5439 Virtual reality simulation in endoscopy training: Current evidence and future directions
Mahmood T, Scaffidi MA, Khan R, Grover SC
- 5446 Quality of life in patients with minimal hepatic encephalopathy
Ridola L, Nardelli S, Gioia S, Riggio O
- 5454 Post-translational modifications of prostaglandin-endoperoxide synthase 2 in colorectal cancer: An update
Jaén RI, Prieto P, Casado M, Martín-Sanz P, Bosca L

ORIGINAL ARTICLE**Basic Study**

- 5462 Counteraction of perforated cecum lesions in rats: Effects of pentadecapeptide BPC 157, L-NAME and L-arginine
Drmic D, Samara M, Vidovic T, Malekinusic D, Antunovic M, Vrdoljak B, Ruzman J, Milkovic Perisa M, Horvat Pavlov K, Jeyakumar J, Seiwert S, Sikiric P
- 5477 Mismatched effects of receptor interacting protein kinase-3 on hepatic steatosis and inflammation in non-alcoholic fatty liver disease
Saeed WK, Jun DW, Jang K, Ahn SB, Oh JH, Chae YJ, Lee JS, Kang HT
- 5491 Near-infrared photoimmunotherapy of pancreatic cancer using an indocyanine green-labeled anti-tissue factor antibody
Aung W, Tsuji AB, Sugyo A, Takashima H, Yasunaga M, Matsumura Y, Higashi T
- 5505 Integrated metabolomic profiling for analysis of antilipidemic effects of *Polygonatum kingianum* extract on dyslipidemia in rats
Yang XX, Wei JD, Mu JK, Liu X, Dong JC, Zeng LX, Gu W, Li JP, Yu J

**Retrospective Study**

- 5288** Safety of hepatitis B virus core antigen-positive grafts in liver transplantation: A single-center experience in China

Lei M, Yan LN, Yang JY, Wen TF, Li B, Wang WT, Wu H, Xu MQ, Chen ZY, Wei YG

ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Harald Peter Hoensch, MD, Emeritus Professor, Marien Hospital Medical Department, Private Practice in Internal Medicine and Gastroenterology, Darmstadt D-64285, Germany

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7th, 14th, 21st, and 28th each month. The *WJG* Editorial Board consists of 642 experts in gastroenterology and hepatology from 59 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

World Journal of Gastroenterology (*WJG*) is now indexed in Current Contents[®]/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch[®]), Journal Citation Reports[®], Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2018 edition of Journal Citation Reports[®] cites the 2017 impact factor for *WJG* as 3.300 (5-year impact factor: 3.387), ranking *WJG* as 35th among 80 journals in gastroenterology and hepatology (quartile in category Q2).

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Shu-Yu Yin*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Rao-Yu Ma*
Proofing Editorial Office Director: *Ze-Mao Gong*

NAME OF JOURNAL
World Journal of Gastroenterology

ISSN
ISSN 1007-9327 (print)
ISSN 2219-2840 (online)

LAUNCH DATE
October 1, 1995

FREQUENCY
Weekly

EDITORS-IN-CHIEF
Andrzej S Tarnawski, MD, PhD, DSc (Med),
Professor of Medicine, Chief Gastroenterology, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach, CA 90822, United States

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <https://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE
Ze-Mao Gong, Director
World Journal of Gastroenterology
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <https://www.f0publishing.com/helpdesk>
<https://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f0publishing.com/helpdesk>
<https://www.wjgnet.com>

PUBLICATION DATE
December 28, 2018

COPYRIGHT
© 2018 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
Full instructions are available online at <https://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<https://www.f0publishing.com>

Virtual reality simulation in endoscopy training: Current evidence and future directions

Tahrin Mahmood, Michael Anthony Scaffidi, Rishad Khan, Samir Chandra Grover

Tahrin Mahmood, Michael Anthony Scaffidi, Rishad Khan, Samir Chandra Grover, Division of Gastroenterology, St. Michael's Hospital, University of Toronto, Toronto M5B 1W8, Canada

ORCID number: Tahrin Mahmood (0000-0001-9013-8166); Michael Anthony Scaffidi (0000-0003-2068-6655); Rishad Khan (0000-0002-5090-7685); Samir Chandra Grover (0000-0003-3392-1220).

Author contributions: Mahmood T, Scaffidi MA, Khan R, and Grover SC drafted the manuscript, critically appraised it for important intellectual content, and approved the final version of this manuscript.

Conflict-of-interest statement: No authors have any conflicts of interest to report of relevance to the subject matter of this manuscript.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Corresponding author: Samir Chandra Grover, MD, MEd, FRCPC, Assistant Professor, Division of Gastroenterology, St. Michael's Hospital, University of Toronto, 30 Bond Street, Toronto M5B 1W8, Canada. samir.grover@utoronto.ca
Telephone: +1-416-864-5628
Fax: +1-416-864-5882

Received: September 29, 2018
Peer-review started: September 29, 2018
First decision: October 26, 2018
Revised: November 22, 2018
Accepted: November 30, 2018
Article in press: December 1, 2018
Published online: December 28, 2018

Abstract

Virtual reality simulation is becoming the standard when beginning endoscopic training. It offers various benefits including learning in a low-stakes environment, improvement of patient safety and optimization of valuable endoscopy time. This is a review of the evidence surrounding virtual reality simulation and its efficacy in teaching endoscopic techniques. There have been 21 randomized controlled trials (RCTs) that have investigated virtual reality simulation as a teaching tool in endoscopy. 10 RCTs studied virtual reality in colonoscopy, 3 in flexible sigmoidoscopy, 5 in esophagogastroduodenoscopy, and 3 in endoscopic retrograde cholangiopancreatography. RCTs reported many outcomes including distance advanced in colonoscopy, comprehensive assessment of technical and non-technical skills, and patient comfort. Generally, these RCTs reveal that trainees with virtual reality simulation based learning improve in all of these areas in the beginning of the learning process. Virtual reality simulation was not effective as a replacement of conventional teaching methods. Additionally, feedback was shown to be an essential part of the learning process. Overall, virtual reality endoscopic simulation is emerging as a necessary augment to conventional learning given the ever increasing importance of patient safety and increasingly valuable endoscopy time; although work is still needed to study the nuances surrounding its integration into curriculum.

Key words: Endoscopy; Gastrointestinal/education; Endoscopy; Gastrointestinal/standards; Simulation; Educational measurement; Clinical competence/standards; Competency-based medical education

© The Author(s) 2018. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: There is substantial evidence to support that virtual reality simulation is an excellent augment to the traditional apprenticeship model in learning endoscopic

procedures. Further work is still needed to study the nuances surrounding its integration into curriculum.

Mahmood T, Scaffidi MA, Khan R, Grover SC. Virtual reality simulation in endoscopy training: Current evidence and future directions. *World J Gastroenterol* 2018; 24(48): 5439-5445
URL: <https://www.wjgnet.com/1007-9327/full/v24/i48/5439.htm>
DOI: <https://dx.doi.org/10.3748/wjg.v24.i48.5439>

INTRODUCTION

Educators in gastroenterology are increasingly integrating simulation-based training (SBT) into endoscopy curricula. SBT allows for a learner-centered environment in which novices can engage in deliberate practice without the fear of making mistakes or harming a patient^[1-4]. The application of SBT to endoscopy has manifested as a variety of model systems, including mechanical models, animal models, live *ex-vivo* models, and virtual reality (VR) computer simulators. Among these, VR simulators have become the most commonly tested model system^[5,6]. In this narrative review, we explore the current state of evidence for the use of VR SBT in endoscopic training.

ENDOSCOPY TRAINING

Apprenticeship model

The apprenticeship model of learning is based on an expert endoscopist (coach) teaching a trainee (apprentice) endoscopic skills. The apprenticeship helps the trainee progress from observation to participation, and finally to independence with progressive responsibilities^[7,8]. The apprenticeship model is based in situated learning theory, which holds that a skill must be learned in the authentic context where it will be applied^[9]. During training, the trainee's responsibility and independence increases as he/she accumulates experience and skill (Figure 1)^[3,5]. Experienced endoscopists intervene if they deem that the trainee will be unable to complete the procedure safely, and trainees are expected to demonstrate certain milestones before progressing to the next steps.

There are, however, several important disadvantages of the apprenticeship model. First, there may be an increased risk of adverse events for patients. For example, one study found that the number of complications of endoscopy are significantly increased in July or August, when the training programs start with their new trainees^[10]. Similarly, a study by Matharoo *et al.*^[11] showed that trainee related factors resulted in as many patient safety incidents as sedation with no oxygen saturation monitor. Second, the staff endoscopist has to completely give up the control of the endoscope in order for the trainee to learn^[5,12]. As the trainee does not have the experience to identify and appropriately manage the findings, the pathological findings, which are intermittent and unpredictable, maybe mismanaged^[5,12]. Third, training endoscopies take longer to complete, adding to the

already strained availability of the endoscopy suite^[5]. Fourth, the feedback given to each trainee depends on the staff endoscopist, and therefore may vary^[13,14]. Finally, there is a lack of continuity as the trainees are expected to work with different staff endoscopists, who may not understand the trainee's level of competence as a whole^[8].

Simulation based training

Given these drawbacks, the endoscopy educators increasingly use SBT prior to first patient contact. SBT provides the trainee with a simulated model of reality to help the trainee understand the skills required for clinical practice^[6,15]. The simulated model can be made physically (*i.e.*, bench top) or with the help of technology using VR^[6,13,15]. SBT is based on the constructivist learning model, which holds that learning is constructed by the learner, rather than transferred to the learner; this renders the context of the situation not as important^[16]. In endoscopic training, SBT has gained attention because it provides a way of training without the risk of adverse events to patients, and allows for standardization of feedback through the simulation^[5,17,18]. Virtual reality simulators can model endoscopy by using an endoscope that is inserted into a computer-based module which displays the gastrointestinal lumen on a screen, and provides visual and tactile feedback related to the procedure^[13].

CURRENT STATE OF EVIDENCE IN

VIRTUAL-REALITY SBT IN ENDOSCOPY

Previous systematic reviews and meta-analyses have effectively analyzed the results of previous studies, including a report of internal validity of the existing literature^[5,17,18]. Briefly, to date there have been 21 major RCTs investigating VR simulation and its role in training (Table 1). 10 of these have studies investigated VR in colonoscopy, 5 in organic gastrointestinal disease (OGD), 3 in flexible sigmoidoscopy, and 3 in endoscopic retrograde cholangiopancreatography ERCP (Table 1)^[19-39].

SBT vs apprenticeship model

18 studies investigated SBT and compared it to either no training or conventional model of learning. 16 of the RCTs reported positive results for SBT, *i.e.*, improved outcomes. Of note is that in the 16 RCTs that reported positive results, SBT was either used in conjunction with conventional teaching, or the control arm did not receive any conventional teaching before the assessment period. A closer look at the two negative trials by Gerson and Sedlack reveal several reasons that can explain the results^[19,20,33]. In the trial by Gerson and Dam, the SBT group had unlimited access for two weeks to didactic modules and six simulated OGD cases on a VR simulator, with no external feedback or observation, while the conventional training group completed 10 OGDs with supervision from staff endoscopists^[20]. As a result, the SBT group completed fewer procedures independently




Figure 1 Progression of learning in the apprenticeship model. The apprenticeship model relies on the trainee modelling after an expert, followed by experimentation and approximation that eventually lead to increased autonomy, adaptation to new problems and eventual mastery.

compared to the conventional training group. In the trial by Sedlack, there is no clear explanation for why the SBT group did worse, however the small sample size of 8 trainees and lack of feedback must be taken into consideration^[33]. In other trials by Sedlack, trainees who received VR training in combination with conventional training achieved better overall ratings of competency and patient comfort^[21,22]. These studies suggest that SBT may have a role in supplementing early endoscopy training. There is, however, no strong evidence to suggest the use of SBT as a replacement for conventional training, giving the limitation of a model in mimicking real life variables and complexities. Furthermore, there is no literature comparing SBT to apprenticeship training for ERCP or other advanced procedures.

SBT with feedback vs without feedback

Feedback is essential in SBT. Recently, Mahmood and Darzi showed that without feedback on performance, SBT does not augment learning^[40]. Specifically, the delivery of feedback should be given with an awareness of the trainee's cognitive load^[41]. Cognitive load is the effort used in working memory and has three types: intrinsic, extraneous, and germane^[41,42]. Intrinsic load relates to essential components of the learning task, *i.e.*, endoscopy, while extrinsic load relates to non-essential tasks, such as distractions related to other staff members^[42]. Germane load relates to forming learning schemas to consolidate the learning, *i.e.*, compartmentalizing concepts to gain mastery^[42]. Learning is thought to occur when trainees form their own cognitive schemas that they can readily access^[42]. In this setting, feedback can help the trainee through negotiating goals for the current session and setting goals for future sessions. Feedback delivered during endoscopic procedures should be minimal to reduce a trainee's cognitive load and should be directive. For example, the expert endoscopist may tell or show the trainee how to change their hand position while navigating the endoscope during the session^[41]. This allows the trainee to incorporate that feedback during that endoscopy and learn from the mistake. After the session, facilitative feedback, which elicits trainee's thoughts should be used to help the trainee engage in reflective learning and develop problem solving skills^[41]. This is supported by previous work suggesting that delayed feedback, when compared with immediate feedback, may be advantageous in the development of cognitive schemas^[41,43].

Integration of SBT in curriculum

Another area of interest in endoscopic training is whether SBT should be embedded in structured curriculum or

self-directed curriculum. The benefit of self-directed curriculum is that it allows learners to set their own learning goals and pace their learning^[44]. However, this is usually best suited for knowledge sharing that is meant to be sustained for a lifetime or for knowledge that is meant to be applied in varied situations after being acquired in one setting - for example, medical students learning pathophysiology that they will be expected to apply on clinical rotations^[45]. Structured learning has a set goal for the trainee, and a trainer to help accomplish that goal^[44]. This goal can be in a form of a certain skill or concept. Structured learning works best for teaching cognitive concepts and technical skills. Grover *et al.*^[30] investigated SBT as part of a structured curriculum compared to self-regulated curriculum, and found that structured curriculum led to better acquisition of technical and non-technical skills that transferred to the clinical setting. The authors concluded that when using SBT to augment the conventional teaching model, it should be through a structured curriculum that encompasses the cognitive, technical, and integrative skills needed for endoscopy.

Structured curricula can be further enhanced by incorporating educational theory-based interventions. For example, two groups applied progressive learning to endoscopy curricula, which involved challenging the learner by increasing the task difficulty and/or complexity as the learner's abilities improves^[29,31]. In both studies, the authors found that training regimens in which trainees encountered progressively more difficult cases led to improved technical skills in the trainees.

FUTURE DIRECTIONS

Simulation based training through virtual-reality modalities has a role in training the novice endoscopist with no or minimal prior experience in endoscopy. SBT offers the opportunity to practice endoscopy in a risk-free environment prior to first patient contact. It is, however, necessary for trainees to undergo patient-based training with an expert endoscopist. Despite the high fidelity of simulators, it difficult to replicate all the variables that a trainee would encounter during a real patient experience, including the non-technical aspects of endoscopy that need to be mastered. As medical education moves towards a competency-based framework, one area that requires elucidation is level of expertise or competency required in the simulated setting before moving onto a real patient. Currently, this distinction is unclear and merits further evaluation. Moreover, the use of educational constructs in SBT should be further explored. Furthermore, there is no data in the literature to date on the cost-effectiveness of VR simulation in endoscopy,

Table 1 Summary of randomized controlled trials

Ref.	Sample size	Intervention	Outcomes
Flexible Sigmoidoscopy			
Tuggy <i>et al</i> ^[19]	10	Group 1: VR simulation Group 2: No simulation	Group 1 performed better: Faster mean completion time (323 s <i>vs</i> 654 s), lower directional errors (1.6% <i>vs</i> 8.6%), higher % of colon visualized (79% <i>vs</i> 45%)
Gerson <i>et al</i> ^[20]	16	Group 1: VR simulation Group 2: Conventional teaching	Group 1 performed worse: Lower mean score (2.9 <i>vs</i> 3.8 out of 5), lower cases completed independently (29% <i>vs</i> 72%), lower retroflexion completed (56% <i>vs</i> 84%); average time, patient satisfaction did not differ
Sedlack <i>et al</i> ^[21]	38	Group 1: VR simulation Group 2: Conventional teaching	Group 1 performed better: Higher patient comfort; procedural skills (independence, identifying pathology, landmarks, performing biopsies, adequate visualization) did not differ
Colonoscopy			
Sedlack <i>et al</i> ^[22]	8	Group 1: VR simulation Group 2: No simulation	Group 1 performed better in first 30 procedures: High depth of unassisted insertion, higher % of procedures completed independently (64.1% <i>vs</i> 56.3%), high scores on other measures such as ability to insert in a safe manner, adequacy visualize mucosa, identify landmarks; mean time to reach maximum insertion did not differ
Ahlberg <i>et al</i> ^[23]	12	Group 1: VR simulation Group 2: No simulation	Group 1 performed better: Higher rates of insertion to cecum (52% <i>vs</i> 19%), shorter procedure time (30 min <i>vs</i> 40 min), less patient discomfort
Cohen <i>et al</i> ^[24]	45	Group 1: VR simulation Group 2: No simulation	Group 1 performed better: Higher competence scores as judged by ability to reach the transverse colon and cecum without assistance (92.7% <i>vs</i> 90.9% by Session 10); patient comfort did not differ
Park <i>et al</i> ^[25]	24	Group 1: VR simulation Group 2: No simulation	Group 1 performed better: Higher global ratings (17.9 <i>vs</i> 14.8 out of 35) based on technique, use of controls, flow of procedures, advancement.
Yi <i>et al</i> ^[26]	11	Group 1: VR simulation Group 2: No simulation	Group 1 performed better: Higher scores during colonoscopy. Higher number of procedures completed independently (76% <i>vs</i> 43%), higher patient comfort; no difference in time or visualization of mucosa
Haycock <i>et al</i> ^[27]	36	Group 1: VR simulation Group 2: Conventional teaching	Group 1 performed better: Higher completion rates (95% <i>vs</i> 70%) and shorter times (407 s <i>vs</i> 743 s), higher patient comfort, higher use of correction abdominal pressure (79% <i>vs</i> 52%), lower insertion force; other variables such as number of transverse loops, correct use of variable stiffness did not differ
McIntosh <i>et al</i> ^[28]	18	Group 1: VR simulation Group 2: No simulation	Group 1 performed better: Less instances of requiring assistance (1.94 <i>vs</i> 3.43), greater unassisted insertion depth (43 cm <i>vs</i> 24 cm), greater rate of cecal intubation (26% <i>vs</i> 10%), high overall competence scores; patient comfort did not differ
Gomez <i>et al</i> ^[29]	27	Group 1: VR simulation + benchtop simulation Group 2: VR simulation Group 3: Benchtop simulation	Group 1 and 2 improved: Performed better on post-test compared to pre-test through Global Assessment of Gastrointestinal Endoscopic Skills tool (navigation, strategies, clear lumen and quality of examination)
Grover <i>et al</i> ^[30]	33	Group 1: Self-regulated learning with VR simulation Group 2: Structured curriculum with VR simulation	Group 1 and 2 improved; Group 2 performed better: Both groups improved on colonoscopy-specific performance; Group 2 performed better based on Joint Advisory Group on GI Endoscopy's Direct Observation of Procedural Skills Tool (JAG DOPS), had better communication rating, and better integrated global rating
Grover <i>et al</i> ^[31]	37	Group 1: Progressive learning with VR simulation Group 2: Non-progressive learning with benchtop simulator	Group 1 performed better: Higher JAG DOPS score, communication and integrated global rating
Esophagogastroduodenoscopy			
Di Giulio <i>et al</i> ^[32]	22	Group 1: VR simulation Group 2: No simulation	Group 1 performed better: Higher number of completed procedures (87.8% <i>vs</i> 70%), required less assistance (41.3% <i>vs</i> 97.9%), overall performance was better; length of time was not significantly different
Sedlack <i>et al</i> ^[33]	8	Group 1: VR simulation Group 2: No simulation	Group 1 performed worse: Lower patient comfort (5 <i>vs</i> 6), independence and competence scores
Shirai <i>et al</i> ^[34]	20	Group 1: VR simulation + Conventional teaching Group 2: Conventional teaching	Group 1 performed better: Required less direct assistance (8.6% <i>vs</i> 25.9%), higher score on 11 items scored during the procedure; no significant difference in completion time
Ferlitsch <i>et al</i> ^[35]	28	Group 1: VR simulation Group 2: No simulation	Group 1 performed better: Decreased total time to reach duodenum (239 s <i>vs</i> 310 s); higher technical accuracy; diagnostic accuracy did not differ
Ende <i>et al</i> ^[36]	29	Group 1: VR simulation + Conventional teaching Group 2: Conventional teaching Group 3: VR simulation alone	Group 1 and 2 improved: Improvement in time within group (195 s <i>vs</i> 119 s; 261 s <i>vs</i> 150 s); no significant difference in between groups All groups showed improvement in post-intervention manual skills test score. None of the other outcomes reached statistical significance, such as time to intubate esophagus
Endoscopic retrograde cholangiopancreatography (ERCP)			
Lim <i>et al</i> ^[37]	16	Group 1: Mechanical simulator Group 2: No simulator	Group 1 performed better: Improved cannulation rates (47.1% <i>vs</i> 69.6%), decreased total time (4.7 <i>vs</i> 10.3 mins); overall performance score not significantly different.

Liao <i>et al</i> ^[38]	16	Group 1: Mechanical simulator Group 2: No simulator	Group 1 performed better: Improved cannulation rates (73.25% <i>vs</i> 47.35%) and improve overall performance; benefit of single <i>vs</i> multiple simulator practices was not statistically significant.
Meng <i>et al</i> ^[39]	5	Group 1: Mechanical simulator Group 2: No simulator	Group 1 performed better: Improved cannulation rates (79.4% <i>vs</i> 61.5%), lower total time (19.38 min <i>vs</i> 26.31 min), and improved overall performance score.

Summary of 21 randomized controlled trials investigating endoscopic techniques, including flexible sigmoidoscopy, colonoscopy, organic gastrointestinal disease and endoscopic retrograde cholangiopancreatography, including sample size, groups in the study and summary of results. Conventional training refers to the apprenticeship model.

which can be expensive in terms of capital costs and costs associated with physician-trainers.

Feedback

During SBT, feedback is often provided by an expert that provides through verbal cues and instructions, and/or by performance metrics from the VR simulator. There are several ways to enhance the delivery of feedback. First, before each simulation session, trainees can be encouraged to document and share their objectives with the expert endoscopist, helping them form a habit of being conscious of their goals during each endoscopic procedure. This enhances learning as it provides the trainer and trainee with a specific goal to comment upon^[41]. Moreover, taking the feedback from one session, the trainees can be encouraged to reflect upon what they will change in their next session, allowing them to understand how to integrate feedback into their training^[41]. SBT in procedural settings allows trainees to observe how incorporating feedback into their next session can result in better outcomes, as other extraneous variables will be controlled. Feedback can be incorporated into endoscopy curricula through gamification, where similar to a game, incremental increase in difficulty and ability to achieve certain goalposts will indicate the level of skill in a trainee^[46]. Second, SBT can be used to help identify trainees that are struggling early in their training and allow programs to supplement their learning. Finally, SBT allows for video-recording of procedures and for trainees to watch videos of both their own performance and expert performance. In endoscopy, watching these videos is associated with improved trainee self-assessment skills over time^[47].

Non-technical skills

Non-technical components of the endoscopic procedure include communication and teamwork, judgement and decision making, leadership, and situational awareness^[48]. Non-technical skills are difficult to teach, especially in a setting where the patient may be partially sedated or even awake. Soft skills in medicine are often taught through role modeling, *i.e.*, apprenticeship^[49]. Similarly, non-technical skills in endoscopy are often taught in the endoscopy suite. However, using high fidelity simulation with standardized patients and actors playing the role of nursing staff to simulate integration scenarios can be used help prime trainees to the non-technical skills of endoscopy^[30].

Emergencies during endoscopy

Dealing with emergencies during endoscopy can be a very stressful experience for trainees. Currently, SBT is limited to polypectomies and other routine procedures, though it may be expanded to include more emergent cases. Similar to Code Blue training, Kiesslich *et al*^[50] showed that simulation training was associated with better endoscopic performance and crisis management in endoscopic emergencies. Additionally, randomized trials in Advanced Cardiac Life Support simulation revealed improved performance in the simulator-trained groups^[51]. SBT allows for trainees to practice the shared mental model and crisis resource management^[50]. Further studies randomized trials can explore these concepts.

Ergonomics

Recently, the importance of ergonomics in endoscopy has become evident. Studies have indicated a high prevalence of musculoskeletal disorders ranging from 37%-89% among endoscopists^[52]. These injuries are thought to be due to repeated pinching or gripping, pushing, pulling and torquing of the endoscope in potentially awkward postures^[53]. Currently, there are no guidelines for optimal ergonomic position due to individual differences in anthropometry. Rather than strict guidelines about positioning during procedures and endoscope handling, some general guidelines may be more valuable. For instance, for computer workstations, guidelines give users recommendations such as keeping the elbow at a right angle with the forearm, which should be parallel to the ground^[54]. In endoscopy, posture while doing the procedure should be neutral without excessive bending. Trainees often tend to bend their body unconsciously when trying to get a better view on the screen. Other tips include to use the endoscope as a lever to reduce right hand torqueing, using a neutral thumb grip position, using gauze with right-hand grip to reduce pressure, and using the left-hand pinky grip technique to reduce right-hand strain in difficult or tight endoscope positions^[55]. SBT can be used to teach these skills to the trainees by helping them practice posturing with a model that mimics the physical aspects of the endoscope workstation. Studies assessing the effectiveness of SBT to teach ergonomics are lacking.

CONCLUSION

SBT is an important supplement to conventional training

to help facilitate learning of endoscopy with no risk to patients. There is, however, insufficient evidence to advocate for SBT as a replacement for conventional training due to its limitation in mimicking real life. Current simulation training curricula may be enhanced with additional focus on the delivery of feedback and the integration of educational theory based strategies. Future studies should broaden the context in which SBT is studied or utilized.

REFERENCES

1. Ziv A, Ben-David S, Ziv M. Simulation based medical education: an opportunity to learn from errors. *Med Teach* 2005; **27**: 193-199 [PMID: 16011941 DOI: 10.1080/01421590500126718]
2. Simulation-Based Learning. Available from: URL: https://www.learning-theories.org/doku.php?id=instructional_design:simulation-based_learning
3. Al-Elq AH. Simulation-based medical teaching and learning. *J Family Community Med* 2010; **17**: 35-40 [PMID: 22022669 DOI: 10.4103/1319-1683.68787]
4. Cook DA, Hatala R, Brydges R, Zendejas B, Szostek JH, Wang AT, Erwin PJ, Hamstra SJ. Technology-enhanced simulation for health professions education: a systematic review and meta-analysis. *JAMA* 2011; **306**: 978-988 [PMID: 21900138 DOI: 10.1001/jama.2011.1234]
5. Walsh CM, Sherlock ME, Ling SC, Carnahan H. Virtual reality simulation training for health professions trainees in gastrointestinal endoscopy. *Cochrane Database Syst Rev* 2012; CD008237 [PMID: 22696375 DOI: 10.1002/14651858.CD008237.pub2]
6. Triantafyllou K, Lazaridis LD, Dimitriadis GD. Virtual reality simulators for gastrointestinal endoscopy training. *World J Gastrointest Endosc* 2014; **6**: 6-12 [PMID: 24527175 DOI: 10.4253/wjge.v6.i1.6]
7. Cognitive Apprenticeship. Available from: URL: https://www.learning-theories.org/doku.php?id=instructional_design:cognitive_apprenticeship
8. Rassie K. The apprenticeship model of clinical medical education: time for structural change. *N Z Med J* 2017; **130**: 66-72 [PMID: 28859068]
9. Ataizi M. Situated Learning. In: Seel NM. Encyclopedia of the Sciences of Learning. Boston, Springer 2012
10. Bini EJ, Firoozi B, Choung RJ, Ali EM, Osman M, Weinshel EH. Systematic evaluation of complications related to endoscopy in a training setting: A prospective 30-day outcomes study. *Gastrointest Endosc* 2003; **57**: 8-16 [PMID: 12518123 DOI: 10.1067/mge.2003.15]
11. Matharoo M, Haycock A, Sevdalis N, Thomas-Gibson S. A prospective study of patient safety incidents in gastrointestinal endoscopy. *Endosc Int Open* 2017; **5**: E83-E89 [PMID: 28191498 DOI: 10.1055/s-0042-117219]
12. Dunkin BJ. Flexible endoscopy simulators. *Semin Laparosc Surg* 2003; **10**: 29-35 [PMID: 12695807]
13. Sedlack RE. The state of simulation in endoscopy education: continuing to advance toward our goals. *Gastroenterology* 2013; **144**: 9-12 [PMID: 23149221 DOI: 10.1053/j.gastro.2012.11.007]
14. Kim JS, Kim BW. Training in Endoscopy: Esophagogastroduodenoscopy. *Clin Endosc* 2017; **50**: 318-321 [PMID: 28783922 DOI: 10.5946/ce.2017.096]
15. Desilets DJ, Banerjee S, Barth BA, Kaul V, Kethu SR, Pedrosa MC, Pfau PR, Tokar JL, Varadarajulu S, Wang A, Wong Kee Song LM, Rodriguez SA; ASGE Technology Committee. Endoscopic simulators. *Gastrointest Endosc* 2011; **73**: 861-867 [PMID: 21521562 DOI: 10.1016/j.gie.2011.01.063]
16. Riegler A. Constructivism. In: L'Abate L. Paradigms in theory construction. New York: Springer, 2012: 235-256
17. Singh S, Sedlack RE, Cook DA. Effects of simulation-based training in gastrointestinal endoscopy: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol* 2014; **12**: 1611-1623.e4 [PMID: 24509241 DOI: 10.1016/j.cgh.2014.01.037]
18. Khan R, Plahouras J, Johnston BC, Scaffidi MA, Grover SC, Walsh CM. Virtual reality simulation training for health professions trainees in gastrointestinal endoscopy. *Cochrane Database Syst Rev* 2018; **8**: CD008237 [PMID: 30117156 DOI: 10.1002/14651858.CD008237.pub3]
19. Tuggy ML. Virtual reality flexible sigmoidoscopy simulator training: impact on resident performance. *J Am Board Fam Pract* 1998; **11**: 426-433 [PMID: 9875997]
20. Gerson LB, Van Dam J. A prospective randomized trial comparing a virtual reality simulator to bedside teaching for training in sigmoidoscopy. *Endoscopy* 2003; **35**: 569-575 [PMID: 12822091 DOI: 10.1055/s-2003-40243]
21. Sedlack RE, Kolars JC, Alexander JA. Computer simulation training enhances patient comfort during endoscopy. *Clin Gastroenterol Hepatol* 2004; **2**: 348-352 [PMID: 15067632]
22. Sedlack RE, Kolars JC. Computer simulator training enhances the competency of gastroenterology fellows at colonoscopy: results of a pilot study. *Am J Gastroenterol* 2004; **99**: 33-37 [PMID: 14687137]
23. Ahlberg G, Hultcrantz R, Jaramillo E, Lindblom A, Arvidsson D. Virtual reality colonoscopy simulation: A compulsory practice for the future colonoscopist? *Endoscopy* 2005; **37**: 1198-1204
24. Cohen J, Cohen SA, Vora KC, Xue X, Burdick JS, Bank S, Bini EJ, Bodenheimer H, Cerulli M, Gerdes H, Greenwald D, Gress F, Grosman I, Hawes R, Mullin G, Schnoll-Sussman F, Starnoli A, Stevens P, Tenner S, Villanueva G. Multicenter, randomized, controlled trial of virtual-reality simulator training in acquisition of competency in colonoscopy. *Gastrointest Endosc* 2006; **64**: 361-368 [PMID: 16923483 DOI: 10.1016/j.gie.2005.11.062]
25. Park J, MacRae H, Musselman LJ, Rossos P, Hamstra SJ, Wolman S, Reznick RK. Randomized controlled trial of virtual reality simulator training: transfer to live patients. *Am J Surg* 2007; **194**: 205-211 [PMID: 17618805 DOI: 10.1016/j.amjsurg.2006.11.032]
26. Yi SY, Ryu KH, Na YJ, Woo HS, Ahn W, Kim WS, Lee DY. Improvement of colonoscopy skills through simulation-based training. *Stud Health Technol Inform* 2008; **132**: 565-567 [PMID: 18391369]
27. Haycock A, Koch AD, Familiari P, van Delft F, Dekker E, Petruzzello L, Haringsma J, Thomas-Gibson S. Training and transfer of colonoscopy skills: a multinational, randomized, blinded, controlled trial of simulator versus bedside training. *Gastrointest Endosc* 2010; **71**: 298-307 [PMID: 19889408 DOI: 10.1016/j.gie.2009.07.017]
28. McIntosh KS, Gregor JC, Khanna NV. Computer-based virtual reality colonoscopy simulation improves patient-based colonoscopy performance. *Can J Gastroenterol Hepatol* 2014; **28**: 203-206 [PMID: 24729994]
29. Gomez PP, Willis RE, Van Sickle K. Evaluation of two flexible colonoscopy simulators and transfer of skills into clinical practice. *J Surg Educ* 2015; **72**: 220-227 [PMID: 25239553 DOI: 10.1016/j.jsurg.2014.08.010]
30. Grover SC, Garg A, Scaffidi MA, Yu JJ, Plener IS, Yong E, Cino M, Grantcharov TP, Walsh CM. Impact of a simulation training curriculum on technical and nontechnical skills in colonoscopy: a randomized trial. *Gastrointest Endosc* 2015; **82**: 1072-1079 [PMID: 26007221 DOI: 10.1016/j.gie.2015.04.008]
31. Grover SC, Scaffidi MA, Khan R, Garg A, Al-Mazroui A, Alomani T, Yu JJ, Plener IS, Al-Awamy M, Yong EL, Cino M, Ravindran NC, Zasowski M, Grantcharov TP, Walsh CM. Progressive learning in endoscopy simulation training improves clinical performance: a blinded randomized trial. *Gastrointest Endosc* 2017; **86**: 881-889 [PMID: 28366440 DOI: 10.1016/j.gie.2017.03.1529]
32. Di Giulio E, Fregonese D, Casetti T, Cestari R, Chilovi F, D'Ambra G, Di Matteo G, Ficano L, Delle Fave G. Training with a computer-based simulator achieves basic manual skills required for upper endoscopy: a randomized controlled trial. *Gastrointest Endosc* 2004; **60**: 196-200 [PMID: 15278044]
33. Sedlack RE. Validation of computer simulation training for esophagogastroduodenoscopy: Pilot study. *J Gastroenterol Hepatol* 2007; **22**: 1214-1219 [PMID: 17559386 DOI: 10.1111/j.1440-1746.2007.04841.x]
34. Shirai Y, Yoshida T, Shiraiishi R, Okamoto T, Nakamura H, Harada T, Nishikawa J, Sakaida I. Prospective randomized study on the use of a computer-based endoscopic simulator for training in esophagogastroduodenoscopy. *J Gastroenterol Hepatol* 2008; **23**: 1046-1050 [PMID: 18554236 DOI: 10.1111/j.1440-1746.2008.05457.

- x]
- 35 **Ferlitsch A**, Schoeff R, Puespoek A, Miehsler W, Schoeniger-Hekele M, Hofer H, Gangl A, Homoncik M. Effect of virtual endoscopy simulator training on performance of upper gastrointestinal endoscopy in patients: a randomized controlled trial. *Endoscopy* 2010; **42**: 1049-1056 [PMID: 20972956 DOI: 10.1055/s-0030-1255818]
 - 36 **Ende A**, Zopf Y, Konturek P, Naegel A, Hahn EG, Matthes K, Maiss J. Strategies for training in diagnostic upper endoscopy: a prospective, randomized trial. *Gastrointest Endosc* 2012; **75**: 254-260 [PMID: 22153875 DOI: 10.1016/j.gie.2011.07.063]
 - 37 **Lim BS**, Leung JW, Lee J, Yen D, Beckett L, Tancredi D, Leung FW. Effect of ERCP mechanical simulator (EMS) practice on trainees' ERCP performance in the early learning period: US multicenter randomized controlled trial. *Am J Gastroenterol* 2011; **106**: 300-306 [PMID: 20978485 DOI: 10.1038/ajg.2010.411]
 - 38 **Liao WC**, Leung JW, Wang HP, Chang WH, Chu CH, Lin JT, Wilson RE, Lim BS, Leung FW. Coached practice using ERCP mechanical simulator improves trainees' ERCP performance: a randomized controlled trial. *Endoscopy* 2013; **45**: 799-805 [PMID: 23897401 DOI: 10.1055/s-0033-1344224]
 - 39 **Meng W**, Leung JW, Yue P, Wang Z, Wang X, Wang H. Practice with ercp mechanical simulator (EMS) improves basic ercp skills of novice surgical trainees. *Gastrointest Endosc* 2016; **83**: AB267-AB268 [DOI: 10.1016/j.gie.2016.03.397]
 - 40 **Mahmood T**, Darzi A. The learning curve for a colonoscopy simulator in the absence of any feedback: no feedback, no learning. *Surg Endosc* 2004; **18**: 1224-1230 [PMID: 15457382 DOI: 10.1007/s00464-003-9143-4]
 - 41 **Dilly CK**, Sewell JL. How to Give Feedback During Endoscopy Training. *Gastroenterology* 2017; **153**: 632-636 [PMID: 28757268 DOI: 10.1053/j.gastro.2017.07.023]
 - 42 **Leppink J**, van den Heuvel A. The evolution of cognitive load theory and its application to medical education. *Perspect Med Educ* 2015; **4**: 119-127 [PMID: 26016429 DOI: 10.1007/s40037-015-0192-x]
 - 43 **Walsh CM**, Ling SC, Wang CS, Carnahan H. Concurrent versus terminal feedback: it may be better to wait. *Acad Med* 2009; **84**: S54-S57 [PMID: 19907387 DOI: 10.1097/ACM.0b013e3181b38daf]
 - 44 **Structured Learning vs Self-Directed Learning**. Available from: URL: <https://eimf.eu/structured-learning-vs-self-directed-learning/>
 - 45 **Hung W**, Jonassen DH, Liu R. Problem-Based Learning. *Handb Res Educ Commun Technol* 2008; **3**: 485-506
 - 46 **Burke B**. Gamification 2020: What Is the Future of Gamification? Available from: URL: <https://www.gartner.com/doc/2226015/gamification--future-gamification>
 - 47 **Khan R**, Scaffidi MA, Parker C, Al-Mazroui A, Tsui C, Iqbal S. Sa1089 The Influence of Video-Based Feedback on Self-Assessment Accuracy of Procedural Skills. *Gastrointest Endosc* 2017; **85**: AB186 [DOI: 10.1016/j.gie.2017.03.408]
 - 48 **Haycock A**. Framework for observing and rating Endoscopic Non-Technical Skills. Available from: URL: https://www.thejag.org.uk/downloads/JAG_training_information/ENTS_handbook_v2.pdf
 - 49 **Dyche L**. Interpersonal skill in medicine: the essential partner of verbal communication. *J Gen Intern Med* 2007; **22**: 1035-1039 [PMID: 17437144 DOI: 10.1007/s11606-007-0153-0]
 - 50 **Kiesslich R**, Moenk S, Reinhardt K, Kanzler S, Schilling D, Jakobs R, Denzer U, Neumann M, Vollmer J, Schütz M, Heinrichs W, Neurath MF, Galle PR. [Combined simulation training: a new concept and workshop is useful for crisis management in gastrointestinal endoscopy]. *Z Gastroenterol* 2005; **43**: 1031-1039 [PMID: 16142611 DOI: 10.1055/s-2005-858542]
 - 51 **Sahu S**, Lata I. Simulation in resuscitation teaching and training, an evidence based practice review. *J Emerg Trauma Shock* 2010; **3**: 378-384 [PMID: 21063561 DOI: 10.4103/0974-2700.70758]
 - 52 **Shergill AK**, McQuaid KR, Rempel D. Ergonomics and GI endoscopy. *Gastrointest Endosc* 2009; **70**: 145-153 [PMID: 19559836 DOI: 10.1016/j.gie.2008.12.235]
 - 53 **ASGE Technology Committee**, Pedrosa MC, Farraye FA, Shergill AK, Banerjee S, Desilets D, Diehl DL, Kaul V, Kwon RS, Mamula P, Rodriguez SA, Varadarajulu S, Song LM, Tierney WM. Minimizing occupational hazards in endoscopy: personal protective equipment, radiation safety, and ergonomics. *Gastrointest Endosc* 2010; **72**: 227-235 [PMID: 20537638 DOI: 10.1016/j.gie.2010.01.071]
 - 54 **Wahlström J**. Ergonomics, musculoskeletal disorders and computer work. *Occup Med (Lond)* 2005; **55**: 168-176 [PMID: 15857896 DOI: 10.1093/occmed/kqi083]
 - 55 **Chang MA**, Mitchell J, Abbas Fehmi SM. Optimizing ergonomics during endoscopy. *VideoGIE* 2017; **2**: 170 [PMID: 29905300 DOI: 10.1016/j.vgie.2017.03.005]

P- Reviewer: Eleftheriadis NP, Hosoe N **S- Editor:** Wang XJ
L- Editor: A **E- Editor:** Yin SY





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>



ISSN 1007-9327



9 771007 932045