

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 2,190,000 条结果 (用时 0.75 秒)

The 1-Year Mortality of Patients Treated in a Hip Fracture Program for ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3597289/> ▼ [翻译此页](#)

作者: S Schnell - 2010 - 被引用次数: 108 - [相关文章](#)

In usual care, the reported 1-year mortality after sustaining a hip fracture has population, comparison data were derived from a thorough literature review. However, this improved mortality was only a trend in the Kaplan-Meier The aftermath of hip fracture: discharge placement, functional status change, and mortality.

Thirty-day mortality after hip fractures: has anything changed?

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4856719/> ▼ [翻译此页](#)

作者: D Giannoulis - 2016 - 被引用次数: 27 - [相关文章](#)

2016年3月4日 - A literature review of relevant publications was carried out for the past ... 30-day mortality rate after hip fractures also demonstrated this trend (Table 3). in the 1-year mortality rate after delayed surgery for a patient with a hip ...

hip fracture in a developing country: a picture in need of change

www.jarcp.com/603-hip-fracture-in-a-developing-country-a-picture-in-need... ▼ [翻译此页](#)

Mortality rates 30 days, 90 days and one year after hospital admission were 3.4%, trends in incidence of hip fractures: a comprehensive literature review.

Length of hospital stay after hip fracture and risk of early mortality after ...

<https://www.bmj.com/content/351/bmj.h6246> - [翻译此页](#)

作者: LE Nikkel - 2015 - 被引用次数: 34 - [相关文章](#)

2015年12月10日 - Mortality rates after hip fracture have changed little over the past 30 The average length of hospital stay was 8.1 days (range 1-995 days; ... Mortality rates did not change between the first six years and the last six years of the study. each year in 2000-11 demonstrated a consistent trend of increasing ...

Similar mortality rates in hip fracture patients over the past 31 years: A ...

<https://www.tandfonline.com/doi/full/10.3109/17453674.2013.878831> - [翻译此页](#)

作者: S Mundi - 2014 - 被引用次数: 73 - [相关文章](#)

2014年2月25日 - Overall, the mean 1-year mortality rate changed from 24% in the 1980s It remains unclear whether mortality after hip fractures has seen similar improvements; few analyses of trends in ... We systematically reviewed the literature to identify clinical trials evaluating the surgical

Name of Journal: *World Journal of Orthopedics*

Manuscript NO: 42794

Manuscript Type: SYSTEMATIC REVIEW

Changing trends in the mortality rate at 1-year post hip fracture - a systematic review

Downey C *et al.* One-year mortality rate post hip fracture

Colum Downey, Martin Kelly, John F Quinlan

Abstract

BACKGROUND

Match Overview

1	Internet 16 words crawled on 20-Jun-2018 journals.sagepub.com	1%
2	Crossref 14 words Simona Scarparo. "CLINICAL AUDIT, GUIDELINES AND STANDARDS: A PRODUCTIVE RELATION FOR MANAGING	<1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 4,090,000 条结果 (用时 0.60 秒)

The 1-Year Mortality of Patients Treated in a Hip Fracture Program for ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3597289/> - [翻译此页](#)

作者 : S Schnell - 2010 - [被引用次数 : 120](#) - [相关文章](#)

In usual care, the reported 1-year mortality after sustaining a hip fracture has been population, comparison data were derived from a thorough literature review. this improved mortality was only a trend in the Kaplan-Meier analysis and not The aftermath of hip fracture: discharge placement, functional status change, ...

Trend in Hip Fracture Incidence and Mortality in Korea: A Prospective ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4366971/> - [翻译此页](#)

作者 : YC Ha - 2015 - [被引用次数 : 30](#) - [相关文章](#)

2015年3月19日 - The crude incidence of hip fractures in the Jeju population ≥50 yr of age increased ... and December 2011 were reviewed to identify patients who had hip fractures. ... A systemic search for death certificates at the National Statistical Office was They reported no change in post-fracture mortality trends.

Role of Comorbidity in Mortality After Hip Fracture: A Nationwide ...

<https://academic.oup.com/aje/article/188/2/398/5166372> - [翻译此页](#)

作者 : A Lunde - 2018 - [相关文章](#)

We investigated whether the combination of comorbidity and hip fracture could ... One-year mortality after hip fracture is more than doubled, with the absolute excess in mortality as a function of time since hip fracture, we used survival analysis. Missing a code for surgery did not change study results substantially, ...

Similar mortality rates in hip fracture patients over the past 31 years: A ...

<https://www.tandfonline.com/doi/full/10.3109/17453674.2013.878831> - [翻译此页](#)

作者 : S Mundi - 2014 - [被引用次数 : 81](#) - [相关文章](#)

2014年2月25日 - Overall, the mean 1-year mortality rate changed from 24% in the 1980s It remains unclear whether mortality after hip fractures has seen similar improvements; few analyses of trends in ... We systematically reviewed the literature to identify clinical trials evaluating the surgical management of hip fractures.

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 3,710,000 条结果 (用时 0.64 秒)

Trend in Hip Fracture Incidence and Mortality in Korea: A Prospective ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4366971/> - 翻译此页

作者: YC Ha - 2015 - 被引用次数: 30 - 相关文章

2015年3月19日 - The crude incidence of hip fractures in the Jeju population ≥ 50 yr of age increased ... We previously performed a hip fracture survey in the Jeju cohort and found that ... A systemic search for death certificates at the National Statistical Office was They reported no change in post-fracture mortality trends.

The 1-Year Mortality of Patients Treated in a Hip Fracture Program for ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3597289/> - 翻译此页

作者: S Schnell - 2010 - 被引用次数: 120 - 相关文章

In usual care, the reported 1-year mortality after sustaining a hip fracture has been population, comparison data were derived from a thorough literature review. this improved mortality was only a trend in the Kaplan-Meier analysis and not The aftermath of hip fracture: discharge placement, functional status change, ...

Meta-analysis: Excess Mortality After Hip Fracture Among Older ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3010729/> - 翻译此页

作者: P Haentjens - 2010 - 被引用次数: 797 - 相关文章

Men with a hip fracture at age 80 years have excess annual mortality of 18%, 22%, 26%, and 20% at 1, 2, 5, and 10 years after injury, respectively. injury and because we wished to establish uniform periods for comparing trends across studies. Regardless of limitations, we believe ours is the first systematic review to ...

Role of Comorbidity in Mortality After Hip Fracture: A Nationwide ...

<https://academic.oup.com/aje/article/188/2/398/5166372> - 翻译此页

作者: A Lunde - 2018 - 相关文章

We investigated whether the combination of comorbidity and hip fracture could ... One-year mortality after hip fracture is more than doubled, with the absolute excess in mortality as a function of time since hip fracture, we used survival analysis. Missing a code for surgery did not change study results substantially, ...

Similar mortality rates in hip fracture patients over the past 31 years: A ...

<https://www.tandfonline.com/doi/full/10.3109/17453674.2013.878831> - 翻译此页

作者: S Mundi - 2014 - 被引用次数: 81 - 相关文章

2014年2月25日 - Overall, the mean 1-year mortality rate changed from 24% in the 1980s It remains