

World Journal of *Cardiology*

World J Cardiol 2019 February 26; 11(2): 47-93



**EDITORIAL**

- 47 Not all arrestins are created equal: Therapeutic implications of the functional diversity of the β -arrestins in the heart

Lymperopoulos A, Wertz SL, Pollard CM, Desimine VL, Maning J, McCrink KA

REVIEW

- 57 Percutaneous devices for left atrial appendage occlusion: A contemporary review

Moussa Pacha H, Al-khadra Y, Soud M, Darmoch F, Moussa Pacha A, Alraies MC

MINIREVIEWS

- 71 Heart valve disease in elderly

Rostagno C

ORIGINAL ARTICLE**Retrospective Study**

- 84 Prevalence and clinical characteristics associated with left atrial thrombus detection: Apixaban

Whiteside HL, Nagabandi A, Brown K, Ayyala DN, Sharma GK

ABOUT COVER

Editor-in-Chief of *World Journal of Cardiology*, Ramdas G Pai, FACC, FRCP (Hon), MD, Doctor, Professor, Director Cardiovascular Fellowship Program, Chair of Clinical Sciences, University of California Riverside School of Medicine, Riverside, CA 92507, United States

AIMS AND SCOPE

World Journal of Cardiology (*World J Cardiol*, *WJC*, online ISSN 1949-8462, DOI: 10.4330) is a peer-reviewed open access journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJC covers topics concerning arrhythmia, heart failure, vascular disease, stroke, hypertension, prevention and epidemiology, dyslipidemia and metabolic disorders, cardiac imaging, pediatrics, etc. Priority publication will be given to articles concerning diagnosis and treatment of cardiology diseases. The following aspects are covered: Clinical diagnosis, laboratory diagnosis, differential diagnosis, imaging tests, pathological diagnosis, molecular biological diagnosis, immunological diagnosis, genetic diagnosis, etc. We encourage authors to submit their manuscripts to *WJC*.

We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

INDEXING/ABSTRACTING

The *WJC* is now abstracted and indexed in Emerging Sources Citation Index (Web of Science), PubMed, PubMed Central, Scopus, China National Knowledge Infrastructure (CNKI), China Science and Technology Journal Database (CSTJ), and Superstar Journals Database.

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: Yun-Xiaojian Wu Proofing Editorial Office Director: Jin-Lei Wang

NAME OF JOURNAL

World Journal of Cardiology

ISSN

ISSN 1949-8462 (online)

LAUNCH DATE

December 31, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Marco Matteo Ciccone, Ramdas G Pai, Dimitrios Tousoulis

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/1949-8462/editorialboard.htm>

EDITORIAL OFFICE

Jin-Lei Wang, Director

PUBLICATION DATE

February 26, 2019

COPYRIGHT

© 2019 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Prevalence and clinical characteristics associated with left atrial thrombus detection: Apixaban

Hoyle L Whiteside, Arun Nagabandi, Kristen Brown, Deepak N Ayyala, Gyanendra K Sharma

ORCID number: Hoyle L Whiteside (0000-0002-7181-2482); Arun Nagabandi (<https://orcid.org/0000-0002-1668-6970>); Kristen Brown (<https://orcid.org/0000-0002-5589-7172>); Deepak N Ayyala (<https://orcid.org/0000-0002-7251-580X>); Gyanendra K Sharma (<https://orcid.org/0000-0001-9027-4440>).

Author contributions: Whiteside HL wrote the manuscript; Whiteside HL, Nagabandi A and Sharma GK reviewed and revised the final version of the manuscript; Whiteside HL, Nagabandi A and Sharma GK designed the study; Whiteside HL, Nagabandi A and Brown K coordinated and provided the data collection; Ayyala DN provided statistical consultation and analysis.

Institutional review board statement: The Institutional Review Board of Augusta University reviewed and approved this study on February 8, 2017.

Informed consent statement: This study was a retrospective analysis utilizing data readily available within the electronic medical record system. The study received a waiver of the consent process upon review by the Institutional Review Board and the dataset was stored electronically on an institutional research drive.

Conflict-of-interest statement: The authors report no financial relationships or conflicts of interest regarding the content.

Open-Access: This article is an

Hoyle L Whiteside, Kristen Brown, Division of Internal Medicine, Medical College of Georgia at Augusta University, Augusta, GA 30912, United States

Arun Nagabandi, Gyanendra K Sharma, Division of Cardiology, Medical College of Georgia at Augusta University, Augusta, GA 30912, United States

Deepak N Ayyala, Division of Biostatistics and Data Science, Department of Population Health Sciences, Medical College of Georgia at Augusta University, Augusta, GA 30912, United States

Corresponding author: Hoyle L Whiteside, MD, Doctor, Division of Internal Medicine, Medical College of Georgia at Augusta University, 1120 15th St., Augusta, GA30909, United States. hwhiteside@augusta.edu

Telephone: +1-706-7212423

Fax: +1-706-7211059

Abstract

BACKGROUND

The prevalence of left atrial appendage (LAA) thrombus detection by transesophageal echocardiogram (TEE) in patients with non-valvular atrial fibrillation (AF) anticoagulated with apixaban is not well defined and identification of additional risk factors may help guide the selection process for pre-procedural TEE. The purpose of our study was to retrospectively analyze the prevalence of LAA thrombus detection by TEE in patients continuously anticoagulated with apixaban for ≥ 4 wk and evaluate for any cardiac risk factors or echocardiographic characteristics which may serve as predictors of thrombus formation.

AIM

To retrospectively analyze the prevalence of LAA thrombus detection by TEE in patients continuously anticoagulated with apixaban.

METHODS

Clinical and echocardiographic data for 820 consecutive patients with AF undergoing TEE at Augusta University Medical Center over a four-year period were retrospectively analyzed. All patients (apixaban: 226) with non-valvular AF and documented compliance with apixaban for ≥ 4 wk prior to index TEE were included.

RESULTS

Following ≥ 4 wk of continuous anticoagulation with apixaban, the prevalence of

open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Received: October 24, 2018

Peer-review started: October 24, 2018

First decision: December 10, 2018

Revised: December 18, 2018

Accepted: January 8, 2019

Article in press: January 9, 2019

Published online: February 26, 2019

LAA thrombus and LAA thrombus/dense spontaneous echocardiographic contrast was 3.1% and 6.6%, respectively. Persistent AF, left ventricular ejection fraction < 30%, severe LA dilation, and reduced LAA velocity were associated with thrombus formation. Following multivariate logistic regression, persistent AF (OR: 7.427; 95%CI: 1.02 to 53.92; $P = 0.0474$), and reduced LAA velocity (OR: 1.086; 95%CI: 1.010 to 1.187; $P = 0.0489$) were identified as independent predictors of LAA thrombus. No Thrombi were detected in patients with a CHA₂DS₂-VASc score ≤ 1 .

CONCLUSION

Among patients with non-valvular AF and ≥ 4 wk of anticoagulation with apixaban, the prevalence of LAA thrombus detected by TEE was 3.1%. This suggests that continuous therapy with apixaban does not completely eliminate the risk of LAA thrombus and that TEE prior to cardioversion or catheter ablation may be of benefit in patients with multiple risk factors.

Key words: Atrial fibrillation; Anticoagulation; Left atrial appendage thrombus; Transesophageal echocardiography

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: The prevalence of left atrial appendage (LAA) thrombus detection by transesophageal echocardiogram (TEE) in patients with non-valvular atrial fibrillation (AF) anticoagulated with apixaban is not well defined and identification of additional risk factors may help guide the selection process for pre-procedural TEE. At our institution, the prevalence of thrombus detection in patients compliant with apixaban was 3.1%. Persistent AF, left ventricular ejection fraction < 30%, severe LA dilation, and reduced LAA velocity were associated with thrombus formation. Following multivariate logistic regression, persistent AF and reduced LAA velocity were identified as independent predictors of thrombus detection.

Citation: Whiteside HL, Nagabandi A, Brown K, Ayyala DN, Sharma GK. Prevalence and clinical characteristics associated with left atrial thrombus detection: Apixaban. *World J Cardiol* 2019; 11(2): 84-93

URL: <https://www.wjgnet.com/1949-8462/full/v11/i2/84.htm>

DOI: <https://dx.doi.org/10.4330/wjc.v11.i2.84>

INTRODUCTION

Atrial fibrillation (AF) is the most common sustained cardiac arrhythmia and is associated with increased risk of systemic thromboembolism due to the development of left atrial (LA) and LA appendage (LAA) thrombi^[1,2]. Among patients with non-valvular AF, 90% of atrial thrombi are seen within the LAA^[3]. Transesophageal echocardiography (TEE) is the most sensitive and specific imaging modality for the detection LA thrombi and is routinely utilized in patients undergoing elective cardioversion or catheter ablation to reduce the risk of thromboembolic events^[1,2,4-7]. Risk factors such as structural heart disease, left atrial size, reduced left ventricular ejection fraction (LVEF), persistent AF, an CHA₂DS₂-VASc score have been reported as independent predictors of LAA thrombus detection by TEE^[8-13].

Several randomized trials have demonstrated the efficacy of non-vitamin K dependent oral anticoagulants (NOACs) to reduce rates of stroke and systemic thromboembolism compared to warfarin, however, their impact on the detection of LA thrombi by TEE is less well established^[14,15]. In patients receiving ≥ 3 wk of continuous anticoagulation with warfarin, the prevalence of LA thrombus detection is reported to be between 1.55% and 7.7%^[8,12,13,16-20]. Recent data has helped to elucidate the prevalence of LA thrombi in patients anticoagulated with NOACs, particularly in patients prescribed dabigatran or rivaroxaban prior to catheter ablation^[18-20]. With regards to apixaban, data remains limited^[18-22]. With at least one study reporting a decline in utilization of TEE prior to catheter ablation and prescription rates for NOACs increasing on a yearly basis, further analysis of patients prescribed apixaban could have clinically meaningful implications^[22,23]. Identification of additional risk

factors which may predict LAA thrombus detection in patients prescribed apixaban is of particular interest as it could help identify a population which would be at increased risk of adverse outcomes should intervention be performed without TEE.

MATERIALS AND METHODS

Study population

Following institutional review board approval, we retrospectively identified 820 consecutive patients with a diagnosis of AF undergoing TEE at Augusta University Medical Center between January 1, 2014 and September 30, 2017 (Figure 1). We excluded 146 patients who were not on any anticoagulation, 183 patients anticoagulated with other NOACs (Rivaroxaban: 122, Dabigatran: 60, Edoxaban: 1), 221 patients anticoagulated with Warfarin, and 15 patients who were determined to be incorrectly coded as AF and in whom TEE was performed for an alternative indication. Two hundred fifty-five patients were anticoagulated with apixaban. Within this cohort, 13 patients were excluded due to documented non-compliance with continuous oral anticoagulation in the 4 wk preceding index TEE, 8 for LAA ligation, 2 for valvular AF, and 6 patients with incomplete echocardiographic data. The final study population included 226 patients anticoagulated with apixaban.

Data extraction and baseline assessment

A detailed chart review was conducted in accordance with the study protocol targeting cardiac risk factors, anticoagulant therapy, and echocardiographic data. A CHA₂DS₂-VASc score was calculated for each patient in accordance with Lip *et al*^[24] AF lasting ≤ 7 d or > 7 d was defined as paroxysmal and persistent, respectively^[1,2]. All physician notes in the four weeks preceding TEE were reviewed. Any patient with documented medication noncompliance was excluded from the study.

Cardiac imaging

TEE Imaging was performed using Phillips EPIQ 7 ultrasound machine and Phillips IE33 ultrasound transducer (Andover, Massachusetts). Standard TEE images were acquired including focused imaging of the LA and LAA. Technique routinely used at our institution involves acquisition of at least two orthogonal views of the LAA. All TEEs were reviewed by at least one of two echocardiographers with strong agreement between observers (Cohen's kappa: 0.89). A thrombus was reported if a well-circumscribed, echo-reflective mass distinct from the LA endocardium or pectinate muscles was present in the appendage or body of the LA^[4]. Spontaneous echo contrast (SEC) was classified as dense, clearing, or absent correlating with 3-4+, 1-2+, or 0 as graded by Fatkin *et al*^[25] SEC was classified as dense if a dense swirling pattern was observed in the LAA and was detectable throughout the cardiac cycle (with variable intensity). SEC was classified as clearing if minimal echodensity was observed in the LAA and was detectable transiently during the cardiac cycle. LAA velocities were determined based on peak velocities averaged over a minimum of two full cardiac cycles in the view which was most parallel to the LAA ostium. LA size was assessed semi-quantitatively and documented as normal, mildly, moderately, or severely dilated.

Follow up

For any patient with LA or LAA thrombi identified on TEE, data was collected for 180 d following index study. If a subsequent TEE was performed between 30 and 180 d following diagnosis of LA or LAA thrombi, then follow up data was collected including changes in oral anticoagulation, antiplatelet therapy, and resolution of thrombus.

Statistical analysis

All statistical computations and hypothesis tests were performed using R 3.4.3 (<https://www.r-project.org/>). To compare demographics and echocardiographic variables between groups, Fisher's exact test was used for categorical variables and Welch's *t*-test was used for the continuous variables. All hypothesis tests were performed at 5% significance level. Univariate logistic regression model was used to associate the odds of occurrence of an event (LAA thrombus formation, dense SEC) to demographic and echocardiographic factors. For scenarios where the odds of the event are zero, pseudo-data points were added to obtain valid estimates and test statistics. Significance of the factors was determined using a two-sided *z*-test. For each univariate model, a two-sided 95% confidence interval (CI) is reported. Using the variables deemed as significant, a multivariate logistic regression was fit to study independence of the variables.




Figure 1 Study population. AC: Anticoagulation; AF: Atrial fibrillation; LAA: Left atrial appendage; NOAC: Non-vitamin K oral anticoagulant.

RESULTS

Population demographics and echocardiographic characteristics

Patient demographics, cardiac risk factors and echocardiographic characteristics for patients with and without thrombus detection are reported in Table 1. The presence of end-stage renal disease, persistent AF, reduced LAA velocities, severe LA dilation, and dense SEC were more common in patients found to have LAA thrombus. Overall, the study population was at meaningful risk of thromboembolic events as 181 (80.1%) patients had CHA₂DS₂-VASc score ≥ 2 .

Prevalence of LAA thrombus and SEC

In patients compliant with apixaban, the prevalence of LAA thrombus and LAA thrombus/Dense SEC was 3.1% and 6.6%, respectively. Full data for the study population based on prevalence of thrombus is provided in Table 1. A similar table, based on the prevalence of thrombus/dense SEC, is available in the online materials (Supplementary Table 1). Among patients with LVEF $< 30\%$ and $\geq 50\%$, thrombus was detected in 9.1% and 2.0% ($P = 0.074$), respectively. The prevalence of LAA thrombus based on CHA₂DS₂-VASc score is summarized in Supplementary Table 2. Notably, no thrombi were identified in the 45 (19.9%) patients with a CHA₂DS₂-VASc score ≤ 1 .

Univariate and multivariate predictors of LAA thrombus

In patients anticoagulated with apixaban, persistent AF, LVEF $< 30\%$, severe LA dilation, and reduced LAA velocity were identified as univariate predictors of LAA thrombus detection (Table 2). On multivariate logistic regression, persistent AF (OR: 7.427; 95%CI: 1.02 to 53.92; $P = 0.0474$), and reduced LAA velocity (OR: 1.086; 95%CI: 1.010 to 1.187; $P = 0.0489$) remained independent predictors of LAA thrombus detection. On further analysis, the OR for each 10% decrease in LVEF was 1.517 (95%CI: 0.971 to 2.369; $P = 0.067$). For the combined endpoint of LAA thrombus detection/dense SEC, reduced LAA velocity (OR: 1.131; 95%CI: 1.031 to 1.235; $p = 0.0061$), was a significant independent predictor on multivariate logistic regression with persistent AF (OR: 4.665; 95%CI: 0.81 to 27.0; $P = 0.0856$) and severe LA dilation (OR: 5.915; 95%CI: 0.74 to 46.98; $P = 0.0927$) approaching significance (Supplementary Table 3).

Subsequent cardiac imaging and thrombus resolution

Six patients had a subsequent TEE performed 30-180 d following the diagnosis of LAA thrombus and documented compliance with continuous anticoagulation. Anticoagulation and echocardiographic data, as well as, CHA₂DS₂-VASc score for each patient is provided in Table 3. These patients were anticoagulated for a mean of

Table 1 Cardiac risk factors and echocardiographic characteristics by presence of thrombus

	Study population (n = 226)	Apixaban-thrombus (n = 7)	Apixaban - no thrombus (n = 219)	P-value ¹
Age	65.8 ± 11.9	68.1 ± 8.0	65.7 ± 11.9	0.458
Race				
White	190 (84.1)	4 (57.1)	186 (84.9)	0.0825
Black	34 (15.0)	3 (42.9)	31 (14.2)	0.0712
Hispanic	0 (0)	0 (0)	0 (0)	-
Asian	2 (0.9)	0 (0)	2 (0.9)	1
Most recent Cr	1.01 ± 0.33	1.18 ± 0.82	1.01 ± 0.32	0.662
ESRD	6 (2.7)	2 (28.5)	4 (1.8)	0.012
Clopidogrel	7 (3.1)	0 (0)	7 (3.2)	1
Aspirin	83 (36.7)	2 (28.5)	81 (37.0)	1
CHA ₂ DS ₂ -VASc	2.83 ± 1.62	3.43 ± 1.40	2.81 ± 1.62	0.354
CHF	88 (34.5)	3 (42.9)	75 (34.2)	0.695
Hypertension	184 (81.4)	7 (100)	177 (80.8)	0.353
Age > 75	50 (22.1)	1 (14.3)	49 (22.4)	1
Diabetes	50 (22.1)	2 (28.5)	48 (21.9)	0.652
Stroke	11 (4.9)	0 (0)	11 (5.0)	1
Vascular disease	26 (11.5)	1 (14.3)	25 (11.4)	0.580
Age > 65	145 (64.2)	6 (85.7)	139 (63.5)	0.426
Female	87 (38.5)	2 (28.5)	85 (38.8)	0.710
Persistent AF	59 (26.1)	5 (71.4)	54 (24.7)	0.014
Echocardiographic data				
LVEF	47.8 ± 14.3	38.6 ± 19.3	48.1 ± 14.1	0.241
LVEF < 30	33 (14.6)	3 (42.9)	30 (13.7)	0.066
LVEF 30-49	44 (19.5)	1 (14.3)	43 (19.6)	1
LVEF ≥ 50	149 (65.9)	3 (42.9)	146 (66.7)	0.233
LAA velocity	48.4 ± 18.1	27.8 ± 10.5	49.0 ± 18.0	0.001
LAA velocity < 40 cm/s	71 (31.4)	6 (85.7)	65 (29.7)	0.004
SEC classification				
None	206 (91.2)	1 (14.3)	205 (93.6)	<0.001
Clearing	7 (3.1)	1 (14.3)	6 (2.7)	0.200
Dense	13 (5.8)	5 (71.4)	8 (3.7)	<0.001
LA dilation				
Moderate	66 (29.2)	1 (14.3)	65 (29.7)	0.677
Severe	59 (26.1)	6 (85.7)	53 (24.2)	0.001
Mitral Regurgitation				
Moderate	53 (23.5)	1 (14.3)	52 (23.7)	1
Severe	5 (2.2)	1 (14.3)	4 (1.8)	0.147

¹P-values denote differences between cohorts with and without thrombus. AF: Atrial fibrillation; CHF: Congestive heart failure; Cr: Creatinine; ESRD: End-stage renal disease; LA: Left atria; LAA: Left atrial appendage; LVEF: Left ventricular ejection fraction; SEC: Spontaneous echo contrast.

94.0 d and thrombus resolution occurred in 83.3%.

DISCUSSION

The purpose of our study was to retrospectively analyze the prevalence of LAA thrombus in patients continuously anticoagulated with apixaban for ≥ 4 wk and evaluate for any cardiac risk factors or echocardiographic characteristics which may serve as predictors of thrombus formation.

Prevalence of LAA thrombus detection

To date, there remains limited data on the prevalence of LAA thrombus detection by TEE amongst patients with non-valvular AF on apixaban therapy. Multiple recent retrospective analyses have helped to elucidate the prevalence of LAA thrombus in

Table 2 Univariate and multivariate predictors of left atrial appendage thrombus

Apixaban						
Variable	Unadjusted (univariate analysis)			Adjusted (multivariate analysis)		
	OR	95%CI	P-value	OR	95%CI	P-value
CHF	1.440	0.31-6.60	0.6388			
Hypertension	1.933	0.24-15.87	0.5396			
Age > 65	3.453	0.41-29.20	0.2552			
Age > 75	0.578	0.07-4.92	0.6160			
Diabetes	1.425	0.27-7.58	0.6778			
Stroke	2.177	0.25-18.85	0.4800			
Vascular disease	1.293	0.15-11.19	0.8152			
CHA ₂ DS ₂ -VASc	1.261	0.80-2.00	0.3240			
Gender (female)	0.631	0.12-3.32	0.5866			
Persistent AF	7.639	1.44-40.51	0.0169 ¹	7.427	1.02-53.92	0.0474
LVEF < 30%	4.725	1.01-22.17	0.0489 ¹	0.726	0.10-5.12	0.7480
LVEF < 50% ²	2.667	0.58-12.23	0.2069			
Severe left atrial dilation ¹	8.877	1.27-61.85	0.0275 ¹	5.901	0.69-50.62	0.1054
LAA velocity (decrease) ¹	1.110	1.031-1.19	0.0032 ¹	1.086	1.010-1.187	0.0489
LAA velocity < 40 cm/s ²	14.215	1.68-120.40	0.0149 ²			

¹Significant factor in the univariate model which is used in the multivariate model;

²Significant factor in the univariate model but not considered in the multivariate model. AF: Atrial fibrillation; CHF: Congestive heart failure; CI: Confidence interval; LAA: Left atrial appendage; LVEF: Left ventricular ejection fraction; OR: Odds ratio.

patients treated with NOACs, in particular, patients undergoing catheter ablation^[18-20,22]. However, apixaban is often the least represented oral anticoagulant in these studies with a reported prevalence of 0% to 2.9%^[21]. In our study population, the prevalence of LAA thrombus was 3.1% despite ≥ 4 wk of continuous anticoagulation which is consistent with previously published data from smaller cohorts when risk factors are considered. The cohort was at considerable risk given mean CHA₂DS₂-VASc 2.83 ± 1.62 and 80.1% of patients with CHA₂DS₂-VASc ≥ 2 .

Predictors of LAA thrombus detection

The presence of persistent AF, reduced LVEF, severe LA dilation, and reduced LAA velocity were identified as univariate predictors of LA thrombus detection in the apixaban cohort. Following evaluation with multivariate logistic regression, persistent AF and reduced LAA velocity were identified as independent predictors of LA thrombus detection. Commonly identified independent predictors of thrombus formation in recent studies include CHF, persistent AF, reduced LVEF, and elevated CHA₂DS₂-VASc score^[18-20]. Of note, apixaban was often the least represented NOAC in these studies and made minimal contribution to the population with thrombus. Finally, these analyses pooled vitamin K antagonist and NOAC data in order to perform multivariate analysis with one exception, in which the authors describe only 1 independent predictor and report small sample size as a limitation^[18-20]. CHA₂DS₂-VASc score was not identified as a univariate predictor which is likely a result of the relatively small number of low-risk patients in our study population as only 19.9% of patients had CHA₂DS₂-VASc score < 2. Of note, reduced LVEF < 30% was identified as a significant univariate predictor in both analyses, however was not determined to be a significant independent predictor. We believe that reduced LVEF is a significant predictor of thrombus formation as identified in similar studies with variable anticoagulation strategies and rates of compliance^[9,13,19,20]. However, our result likely reflects a limitation of sample size as well as an inherent relationship between advanced cardiomyopathy and clinical/echocardiographic findings most prevalent in high-risk patients with AF.

Rate of LAA thrombus resolution

Data regarding thrombus resolution in patients prescribed apixaban has thus far been limited to case reports and small cohorts^[19,20,26,27]. While data regarding thrombus resolution in patients prescribed warfarin is more prevalent, rates of resolution range

Table 3 Clinical characteristics and thrombus resolution

Study ID	AC after thrombus identification	Thrombus resolution	Duration of AC (d)	P2Y12 inhibitor	Aspirin	LVEF (%)	LA dilation	Spontaneous contrast	LAA velocity (cm/s)	Duration of A-fib	CHADS-VASc score
Apixaban following index TEE (<i>n</i> = 6)											
LA-011	Apixaban	Yes	143	No	No	55	Moderate	None	30.4	Paroxysmal	3
LA-016	Apixaban	Yes	38	Yes	Yes	15	Severe	Severe	14.4	Paroxysmal	2
LA-017	Apixaban	Yes	175	No	Yes	15	Severe	Mild	34.0	Paroxysmal	4
LA-019	Apixaban	Yes	40	No	No	30	Severe	None	26.5	Paroxysmal	2
LA-020	Apixaban	Yes	56	No	Yes	25	Severe	Moderate	28.0	Persistent	3
LA-005	Apixaban	No	112	No	No	65	Moderate	Moderate	49.2	Persistent	4
		5/6 (83.3%)	Mean 94.0 Median 84.0								

AC: Anticoagulation; A-fib: Atrial fibrillation; LA: Left atrium; LAA: Left atrial appendage; LVEF: Left ventricular ejection fraction; TEE: Transesophageal echocardiogram.

from 55% to 82%^[16,17,28]. In our limited cohort, we identified thrombus resolution in 83.3% of patients anticoagulated with apixaban (*n* = 6). Although all patients were confirmed to be compliant with continuous anticoagulation throughout the follow up period, there was significant heterogeneity in the duration of therapy prior to repeat cardiac imaging which limits our ability to draw conclusions regarding the optimal duration of anticoagulation. Regardless, apixaban appears to be a reasonable anticoagulation strategy in this population and warrants further investigation in prospective trial.

Application of study findings

A recent expert consensus statement recommends that current anticoagulation guidelines as they pertain to cardioversion of AF should be observed for patients presenting with AF prior to catheter ablation and that TEE is reasonable despite ≥ 3 wk of continuous anticoagulation^[1,2,7]. Two recent surveys, one including 16 Canadian centers and the other including 521 ablation centers in 24 countries, report that > 70% of ablation centers routinely utilized pre-procedure TEE in all patients^[29,30]. One cost-effectiveness analysis reports an incremental cost-effectiveness ratio of \$226608 per quality-adjusted life year for routine use of TEE in an unselected population prior to pulmonary vein isolation modeled with a 4% prevalence of thrombus. While the prevalence is likely overestimated, this analysis highlights the need to better identify patients with a high pretest probability of LAA thrombus despite continuous anticoagulation in order to improve the cost-benefit ratio of the procedure^[31]. A trend toward more conservative use of pre-procedural TEE appears to be underway as one large ablation center reports a significant decline in the routine utilization of TEE from 86% to 42% over a 5 year period^[22]. While another recent study completely eliminated pre-procedural TEE in favor of intracardiac echocardiography prior to AF ablation. Despite adequate imaging of the LAA in only 71% of patients, the authors report excellent outcomes^[32]. This study is retrospective and meant to explore variables which could be predictors of thrombus formation in patients treated with apixaban. A prospective randomized trial would be needed to conclusively determine and validate a scoring system and/or various cutoffs. However, this may not be practical given the low event rate in this population. Nonetheless, we hope that our work can provide evidence to help guide the selection of patients for pre-procedural TEE.

Study limitations

Our study is limited by the retrospective nature of the data collected. In addition, we cannot objectively confirm 100% compliance with apixaban therapy as quantitative assays are not routinely used in clinical practice. Although we took great effort to exclude any patients with documented non-compliance, our ability to do so would be limited by the history provided and documentation of health care professionals.

In patients with non-valvular AF and a minimum of 4 wk continuous oral

anticoagulation with apixaban, the prevalence of LAA thrombus and LAA thrombus/dense SEC detected by TEE was 3.1% and 6.6%, respectively. Both persistent AF and reduced LAA velocity were identified as independent predictors of LA thrombus detection in patients anticoagulated with apixaban. In addition, LVEF < 30% and severe LA dilation were identified as univariate predictors. We hope that the presence or absence of these clinical findings in addition to established risk factors can help guide the selection process for utilization of pre-procedural TEE in future patients with non-valvular AF anticoagulated with apixaban.

ARTICLE HIGHLIGHTS

Research background

The prevalence of left atrial appendage (LAA) thrombus detection by transesophageal echocardiogram (TEE) in patients anticoagulated for ≥ 4 wk with apixaban is not well defined and predictors of LAA thrombus detection are not completely understood. Furthermore, the efficacy of apixaban to resolve pre-existing LAA thrombi is not well documented.

Research motivation

Prescriptions rates for non-vitamin K dependent oral anticoagulants are increasing on a yearly basis and further analysis of patients prescribed apixaban could have clinically meaningful implications. We aimed to identify significant predictors of LAA thrombus detection on TEE to aid in the selection process for screening in future patients undergoing direct current cardioversion or catheter ablation.

Research objectives

The purpose of our study was to retrospectively analyze the prevalence of LAA thrombus detection by TEE in patients continuously anticoagulated with apixaban for ≥ 4 wk and evaluate for any cardiac risk factors or echocardiographic characteristics which may serve as predictors of thrombus formation.

Research methods

Clinical and echocardiographic data for 820 consecutive patients with atrial fibrillation (AF) undergoing TEE at Augusta University Medical Center over a four-year period were retrospectively analyzed. All patients (apixaban: 226) with non-valvular AF and documented compliance with apixaban for ≥ 4 wk prior to index TEE were included.

Research results

Following ≥ 4 wk of continuous anticoagulation with apixaban, the prevalence of LAA thrombus and LAA thrombus/dense spontaneous echocardiographic contrast was 3.1% and 6.6%, respectively. Persistent AF, left ventricular ejection fraction < 30%, severe LA dilation, and reduced LAA velocity were associated with thrombus formation. Following multivariate logistic regression, persistent AF (OR: 7.427; 95%CI: 1.02 to 53.92; $P = 0.0474$), and reduced LAA velocity (OR: 1.086; 95%CI: 1.010 to 1.187; $P = 0.0489$) were identified as independent predictors of LAA thrombus. No Thrombi were detected in patients with a CHA₂DS₂-VASc score ≤ 1 .

Research conclusions

Among patients with non-valvular AF and ≥ 4 wk of anticoagulation with apixaban, the prevalence of LAA thrombus detected by TEE was 3.1%. This suggests that continuous therapy with apixaban does not completely eliminate the risk of LAA thrombus and that TEE prior to cardioversion or catheter ablation may be of benefit in patients with multiple risk factors.

Research perspectives

Compliance with non-vitamin K oral anticoagulants reduces but does not eliminate the prevalence of thrombus detection by TEE. However, available cost-effectiveness analysis reports that pre-procedural TEE is unlikely to be cost-effective in an unselected population. Therefore, there is a need to better identify patients with increased pretest probability of LAA thrombus in order to improve the cost-benefit ratio of the procedure. It is our hope that identification of additional clinical and echocardiographic characteristics; in addition to established risk factors, can help guide the selection process for utilization of pre-procedural TEE

REFERENCES

- 1 January CT, Wann LS, Alpert JS, Calkins H, Cigarroa JE, Cleveland JC, Conti JB, Ellnor PT, Ezekowitz MD, Field ME, Murray KT, Sacco RL, Stevenson WG, Tchou PJ, Tracy CM, Yancy CW; ACC/AHA Task Force Members. 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines and the Heart Rhythm Society. *Circulation* 2014; **130**: e199-e267 [PMID: 24682347 DOI: 10.1161/CIR.0000000000000041]
- 2 Kirchhof P, Benussi S, Kotecha D, Ahlsson A, Atar D, Casadei B, Castella M, Diener HC, Heidbuchel H, Hendriks J, Hindricks G, Manolis AS, Oldgren J, Popescu BA, Schotten U, Van Putte B, Vardas P; ESC Scientific Document Group. 2016 ESC Guidelines for the management of atrial fibrillation developed in

- collaboration with EACTS. *Eur Heart J* 2016; **37**: 2893-2962 [PMID: [27567408](#) DOI: [10.1093/eurheartj/ehw210](#)]
- 3 **Al-Saady NM**, Obel OA, Camm AJ. Left atrial appendage: structure, function, and role in thromboembolism. *Heart* 1999; **82**: 547-554 [PMID: [10525506](#) DOI: [10.1136/hrt.82.5.547](#)]
 - 4 **Manning WJ**, Weintraub RM, Waksmonski CA, Haering JM, Rooney PS, Maslow AD, Johnson RG, Douglas PS. Accuracy of transesophageal echocardiography for identifying left atrial thrombi. A prospective, intraoperative study. *Ann Intern Med* 1995; **123**: 817-822 [PMID: [7486462](#) DOI: [10.7326/0003-4819-123-11-199512010-00001](#)]
 - 5 **Manning WJ**, Silverman DI, Gordon SP, Krumholz HM, Douglas PS. Cardioversion from atrial fibrillation without prolonged anticoagulation with use of transesophageal echocardiography to exclude the presence of atrial thrombi. *N Engl J Med* 1993; **328**: 750-755 [PMID: [8437595](#) DOI: [10.1056/NEJM199303183281102](#)]
 - 6 **Klein AL**, Grimm RA, Murray RD, Apperson-Hansen C, Asinger RW, Black IW, Davidoff R, Erbel R, Halperin JL, Orsinelli DA, Porter TR, Stoddard MF; Assessment of Cardioversion Using Transesophageal Echocardiography Investigators. Use of transesophageal echocardiography to guide cardioversion in patients with atrial fibrillation. *N Engl J Med* 2001; **344**: 1411-1420 [PMID: [11346805](#) DOI: [10.1056/NEJM200105103441901](#)]
 - 7 **Calkins H**, Hindricks G, Cappato R, Kim YH, Saad EB, Aguinaga L, Akar JG, Badhwar V, Brugada J, Camm J, Chen PS, Chen SA, Chung MK, Nielsen JC, Curtis AB, Davies DW, Day JD, d'Avila A, de Groot NMSN, Di Biase L, Duytschaever M, Edgerton JR, Ellenbogen KA, Ellinor PT, Ernst S, Fenelon G, Gerstenfeld EP, Haines DE, Haissaguerre M, Helm RH, Hylek E, Jackman WM, Jalife J, Kalman JM, Kautzner J, Kottkamp H, Kuck KH, Kumagai K, Lee R, Lewalter T, Lindsay BD, Macle L, Mansour M, Marchlinski FE, Michaud GF, Nakagawa H, Natale A, Nattel S, Okumura K, Packer D, Pokushalov E, Reynolds MR, Sanders P, Scanavacca M, Schilling R, Tondo C, Tsao HM, Verma A, Wilber DJ, Yamane T. 2017 HRS/EHRA/ECAS/APHRS/SOLAECE expert consensus statement on catheter and surgical ablation of atrial fibrillation. *Heart Rhythm* 2017; **14**: e275-e444 [PMID: [28506916](#) DOI: [10.1016/j.hrthm.2017.05.012](#)]
 - 8 **Wallace TW**, Atwater BD, Daubert JP, Voora D, Crowley AL, Bahnson TD, Hranitzky PM. Prevalence and clinical characteristics associated with left atrial appendage thrombus in fully anticoagulated patients undergoing catheter-directed atrial fibrillation ablation. *J Cardiovasc Electrophysiol* 2010; **21**: 849-852 [PMID: [20158561](#) DOI: [10.1111/j.1540-8167.2010.01729.x](#)]
 - 9 **Puwanant S**, Varr BC, Shrestha K, Hussain SK, Tang WH, Gabriel RS, Wazni OM, Bhargava M, Saliba WI, Thomas JD, Lindsay BD, Klein AL. Role of the CHADS2 score in the evaluation of thromboembolic risk in patients with atrial fibrillation undergoing transesophageal echocardiography before pulmonary vein isolation. *J Am Coll Cardiol* 2009; **54**: 2032-2039 [PMID: [19926009](#) DOI: [10.1016/j.jacc.2009.07.037](#)]
 - 10 **Yamamoto M**, Seo Y, Kawamatsu N, Sato K, Sugano A, Machino-Ohtsuka T, Kawamura R, Nakajima H, Igarashi M, Sekiguchi Y, Ishizu T, Aonuma K. Complex left atrial appendage morphology and left atrial appendage thrombus formation in patients with atrial fibrillation. *Circ Cardiovasc Imaging* 2014; **7**: 337-343 [PMID: [24523417](#) DOI: [10.1161/CIRCIMAGING.113.001317](#)]
 - 11 **Calvo N**, Mont L, Vidal B, Nadal M, Montserrat S, Andreu D, Tamborero D, Pare C, Azqueta M, Berrueto A, Brugada J, Sitges M. Usefulness of transoesophageal echocardiography before circumferential pulmonary vein ablation in patients with atrial fibrillation: is it really mandatory? *Europace* 2011; **13**: 486-491 [PMID: [21186230](#) DOI: [10.1093/europace/euq456](#)]
 - 12 **Scherr D**, Dalal D, Chilukuri K, Dong J, Spragg D, Henrikson CA, Nazarian S, Cheng A, Berger RD, Abraham TP, Calkins H, Marine JE. Incidence and predictors of left atrial thrombus prior to catheter ablation of atrial fibrillation. *J Cardiovasc Electrophysiol* 2009; **20**: 379-384 [PMID: [19017348](#) DOI: [10.1111/j.1540-8167.2008.01336.x](#)]
 - 13 **McCready JW**, Nunn L, Lambiase PD, Ahsan SY, Segal OR, Rowland E, Lowe MD, Chow AW. Incidence of left atrial thrombus prior to atrial fibrillation ablation: is pre-procedural transoesophageal echocardiography mandatory? *Europace* 2010; **12**: 927-932 [PMID: [20304842](#) DOI: [10.1093/europace/euq074](#)]
 - 14 **Savarese G**, Giugliano RP, Rosano GM, McMurray J, Magnani G, Filippatos G, DelleGrottaglie S, Lund LH, Trimarco B, Perrone-Filardi P. Efficacy and Safety of Novel Oral Anticoagulants in Patients With Atrial Fibrillation and Heart Failure: A Meta-Analysis. *JACC Heart Fail* 2016; **4**: 870-880 [PMID: [27614940](#) DOI: [10.1016/j.jchf.2016.07.012](#)]
 - 15 **Ruff CT**, Giugliano RP, Braunwald E, Hoffman EB, Deenadayalu N, Ezekowitz MD, Camm AJ, Weitz JJ, Lewis BS, Parkhomenko A, Yamashita T, Antman EM. Comparison of the efficacy and safety of new oral anticoagulants with warfarin in patients with atrial fibrillation: a meta-analysis of randomised trials. *Lancet* 2014; **383**: 955-962 [PMID: [24315724](#) DOI: [10.1016/S0140-6736\(13\)62343-0](#)]
 - 16 **Jaber WA**, Prior DL, Thamilarasan M, Grimm RA, Thomas JD, Klein AL, Asher CR. Efficacy of anticoagulation in resolving left atrial and left atrial appendage thrombi: A transesophageal echocardiographic study. *Am Heart J* 2000; **140**: 150-156 [PMID: [10874278](#) DOI: [10.1067/mhj.2000.106648](#)]
 - 17 **Seidl K**, Rameken M, Drögemüller A, Vater M, Brandt A, Schwacke H, Bergmeier C, Zahn R, Senges J. Embolic events in patients with atrial fibrillation and effective anticoagulation: value of transesophageal echocardiography to guide direct-current cardioversion. Final results of the Ludwigshafen Observational Cardioversion Study. *J Am Coll Cardiol* 2002; **39**: 1436-1442 [PMID: [11985904](#) DOI: [10.1016/S0735-1097\(02\)01785-0](#)]
 - 18 **Wyrembak J**, Campbell KB, Steinberg BA, Bahnson TD, Daubert JP, Velazquez EJ, Samad Z, Atwater BD. Incidence and Predictors of Left Atrial Appendage Thrombus in Patients Treated With Nonvitamin K Oral Anticoagulants Versus Warfarin Before Catheter Ablation for Atrial Fibrillation. *Am J Cardiol* 2017; **119**: 1017-1022 [PMID: [28153350](#) DOI: [10.1016/j.amjcard.2016.12.008](#)]
 - 19 **Frenkel D**, D'Amato SA, Al-Kazaz M, Markowitz SM, Liu CF, Thomas G, Ip JE, Sharma SK, Yang H, Singh P, Lerman BB, Cheung JW. Prevalence of Left Atrial Thrombus Detection by Transesophageal Echocardiography: A Comparison of Continuous Non-Vitamin K Antagonist Oral Anticoagulant Versus Warfarin Therapy in Patients Undergoing Catheter Ablation for Atrial Fibrillation. *JACC Clin Electrophysiol* 2016; **2**: 295-303 [PMID: [29766887](#) DOI: [10.1016/j.jacep.2016.01.004](#)]
 - 20 **Da Costa A**, Delolme C, Guichard JB, Gerbay A, Pierrard R, Romeyer-Bouchard C, Isaaz K. Comparison of prevalence and management of left atrial appendage thrombi under old and new anticoagulants prior to left atrial catheter ablation. *Am Heart J* 2017; **193**: 8-15 [PMID: [29129259](#) DOI: [10.1016/j.ahj.2017.07.016](#)]

- 21 **Flaker G**, Lopes RD, Al-Khatib SM, Hermosillo AG, Hohnloser SH, Tinga B, Zhu J, Mohan P, Garcia D, Bartunek J, Vinereanu D, Husted S, Harjola VP, Rosenqvist M, Alexander JH, Granger CB; ARISTOTLE Committees and Investigators. Efficacy and safety of apixaban in patients after cardioversion for atrial fibrillation: insights from the ARISTOTLE Trial (Apixaban for Reduction in Stroke and Other Thromboembolic Events in Atrial Fibrillation). *J Am Coll Cardiol* 2014; **63**: 1082-1087 [PMID: 24211508 DOI: 10.1016/j.jacc.2013.09.062]
- 22 **Balouch M**, Gucuk Ipek E, Chrispin J, Bajwa RJ, Zghaib T, Berger RD, Ashikaga H, Calkins H, Nazarian S, Marine JE, Spragg DD. Trends in Transesophageal Echocardiography Use, Findings, and Clinical Outcomes in the Era of Minimally Interrupted Anticoagulation for Atrial Fibrillation Ablation. *JACC Clin Electrophysiol* 2017; **3**: 329-336 [PMID: 29759444 DOI: 10.1016/j.jacep.2016.09.011]
- 23 **Marzec LN**, Wang J, Shah ND, Chan PS, Ting HH, Gosch KL, Hsu JC, Maddox TM. Influence of Direct Oral Anticoagulants on Rates of Oral Anticoagulation for Atrial Fibrillation. *J Am Coll Cardiol* 2017; **69**: 2475-2484 [PMID: 28521884 DOI: 10.1016/j.jacc.2017.03.540]
- 24 **Lip GY**, Nieuwlaet R, Pisters R, Lane DA, Crijns HJ. Refining clinical risk stratification for predicting stroke and thromboembolism in atrial fibrillation using a novel risk factor-based approach: the euro heart survey on atrial fibrillation. *Chest* 2010; **137**: 263-272 [PMID: 19762550 DOI: 10.1378/chest.09-1584]
- 25 **Fatkin D**, Kelly RP, Feneley MP. Relations between left atrial appendage blood flow velocity, spontaneous echocardiographic contrast and thromboembolic risk in vivo. *J Am Coll Cardiol* 1994; **23**: 961-969 [PMID: 8106703 DOI: 10.1016/0735-1097(94)90644-0]
- 26 **Dobashi S**, Fujino T, Ikeda T. Use of apixaban for an elderly patient with left atrial thrombus. *BMJ Case Rep* 2014; **2014**: pii: bcr2014203870 [PMID: 24962484 DOI: 10.1136/bcr-2014-203870]
- 27 **Miwa Y**, Minamishima T, Sato T, Sakata K, Yoshino H, Soejima K. Resolution of a warfarin and dabigatran-resistant left atrial appendage thrombus with apixaban. *J Arrhythm* 2016; **32**: 233-235 [PMID: 27354873 DOI: 10.1016/j.joa.2016.01.009]
- 28 **Corrado G**, Tadeo G, Beretta S, Tagliagambe LM, Manzillo GF, Spata M, Santarone M. Atrial thrombi resolution after prolonged anticoagulation in patients with atrial fibrillation. *Chest* 1999; **115**: 140-143 [PMID: 9925075 DOI: 10.1378/chest.115.1.140]
- 29 **Cappato R**, Calkins H, Chen SA, Davies W, Iesaka Y, Kalman J, Kim YH, Klein G, Natale A, Packer D, Skanes A, Ambrogi F, Biganzoli E. Updated worldwide survey on the methods, efficacy, and safety of catheter ablation for human atrial fibrillation. *Circ Arrhythm Electrophysiol* 2010; **3**: 32-38 [PMID: 19995881 DOI: 10.1161/CIRCEP.109.859116]
- 30 **Awad S**, Mohajer K, Baranchuk A, Pal RS. Current clinical practice of transesophageal echocardiography and cardiac computed tomography prior to atrial fibrillation ablation in Canada. *Int J Cardiol* 2015; **189**: 300-301 [PMID: 25967570 DOI: 10.1016/j.ijcard.2015.03.318]
- 31 **Gula LJ**, Massel D, Redfearn DP, Krahn AD, Yee R, Klein GJ, Skanes AC. Impact of routine transoesophageal echocardiography on safety, outcomes, and cost of pulmonary vein ablation: inferences drawn from a decision analysis model. *Europace* 2010; **12**: 1550-1557 [PMID: 20716548 DOI: 10.1093/europace/euq306]
- 32 **Di Biase L**, Briceno DF, Trivedi C, Mohanty S, Gianni C, Burkhardt JD, Mohanty P, Bai R, Gunda S, Horton R, Bailey S, Sanchez JE, Al-Ahmad A, Hranitzky P, Gallingshouse GJ, Reddy YM, Zagrodzky J, Hongo R, Beheiry S, Lakkireddy D, Natale A. Is transesophageal echocardiogram mandatory in patients undergoing ablation of atrial fibrillation with uninterrupted novel oral anticoagulants? Results from a prospective multicenter registry. *Heart Rhythm* 2016; **13**: 1197-1202 [PMID: 26994940 DOI: 10.1016/j.hrthm.2016.03.024]

P- Reviewer: Hochholzer W, Kharlamov AN

S- Editor: Ji FF L- Editor: A E- Editor: Wu YXJ





Published By Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

