



停止显示此广告

为什么显示该广告？



时间不限

2018以来

2017以来

2014以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

☒ 创建快讯

Google 已关闭此广告

停止显示此广告

为什么显示该广告？

Google 已关闭此广告

停止显示此广告

为什么显示该广告？

Google 已关闭此广告

停止显示此广告

为什么显示该广告？

小提示：只搜索中文(简体)结果，可在 学术搜索设置 指定搜索语言

Analgesia for total hip and knee arthroplasty: a multimodal pathway featuring peripheral nerve block

TT Horlocker, SL Kopp, MW Pagnano... - ... Orthopaedic Surgeons, 2006 - journals.lww.com

You may be trying to access this site from a secured browser on the server. Please enable scripts and reload this page ...

☆ 99 被引用次数：196 相关文章 所有 4 个版本

sci-hub下载

Does regional anaesthesia improve outcome after total hip arthroplasty? A systematic review

AJR Macfarlane, GA Prasad... - British Journal of ..., 2009 - academic.oup.com

... You can change your cookie settings at any time ... Oxford University Press. Search; Account Menu; Menu; Sign In. Register. Navbar Search Filter All BJA: British Journal of Anaesthesia All Journals Mobile Microsite Search Term ... Canada. M5T 2S8. Search for other works by this ...

☆ 99 被引用次数：190 相关文章 所有 8 个版本

[HTML] oup.com

Full View

Pain management in total joint arthroplasty: a historical review

TT Horlocker - Orthopedics, 2010 - healio.com

... postoperative pain: seven years' experience with 5969 surgical patients at Indiana University ... Practice guidelines for the prevention, detection, and management of respiratory depression associated ... of the quality of postoperative analgesia and complications in 1,416 patients ...

☆ 99 被引用次数：69 相关文章 所有 10 个版本

[PDF] wyanokecdn.com

Single fascia iliaca compartment block for post-hip fracture pain relief

DG Monzon, KV Iserson, JA Vazquez - The Journal of emergency medicine, 2007 - Elsevier

... the FICB has been shown to be much simpler to perform, have a more rapid onset, and be more effective, at least in adults, for simultaneously blocking the ... The 3-in-1 block often fails because the catheter is not placed under the fascia iliaca, and the success rate does not ...

[PDF] academia.edu

Match Overview

1	Crossref 15 words Xu-feng Jia, Yong Ji, Guang-ping Huang, Yu Zhou, Miao Long. "Comparison of intrathecal and local infiltration analgesia	1%
2	Crossref 12 words Singelyn, F.J.E., "Continuous femoral and popliteal sciatic ... erve blockades", Techniques in Regional Anesthesia and Pa	1%
3	Internet 10 words crawled on 15-Mar-2019 f6publishing.blob.core.windows.net	1%

3
Name of Journal: *World Journal of Anesthesiology*

Manuscript NO: 43284

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Enhanced recovery after surgery pathway: The use of fascia iliaca blocks causes delayed ambulation after total hip arthroplasty

Jamie-Lee Metesky, Junping Chen, Meg Rosenblatt

Abstract

BACKGROUND

Fascia iliaca compartment blocks (FIBs) have been used to provide postoperative analgesia after total hip arthroplasty (THA). However, evidence of their efficacy remains limited. While pain control appears to be satisfactory, quadriceps weakness may be an untoward consequence of the block. Prior studies have shown femoral nerve blocks and fascia iliaca blocks as being superior for pain control and ambulation following THA when compared to standard therapy of parenteral pain control.

[国内版](#)[国际版](#)[Chat with Bing](#)

Enhanced recovery after surgery pathway: The use of fascia iliaca blocks causes de

[Sign in](#)[All](#)[Images](#)[Videos](#)[翻译成中文](#)[关闭取词](#)[Make Bing your search engine >](#)

4 Results

Any time ▾

A Modified Fascia Iliaca Compartment Block Has Significant ...

https://www.researchgate.net/publication/5763713_A_Modified_Fascia...

Despite being commonly used, there is a lack of evidence relating to the use of a fascia iliaca block for analgesia after total hip arthroplasty.

Journal of Pain Research - Dove Press Open Access Publisher

<https://www.dovepress.com/journal-of-pain-research-archive41> ▾

Effects of fascia iliaca compartment block combined with general laryngeal mask airway anesthesia in children undergoing femoral fracture surgery: a randomized trial ... Paravertebral blocks and enhanced recovery after surgery protocols in breast reconstructive surgery: patient selection and perspectives ... In-hospital outcomes with anterior ...

Practical Plastic Surgery - PDF Free Download

<https://epdf.tips/practical-plastic-surgery.html> ▾

Postoperative nausea and vomiting is the major factor contributing to unintentional hospital admission after outpatient surgery. It begins shortly after arrival in the recovery room and usually lasts no longer than 12-24 hours postoperatively.

(PDF) A randomised, controlled, double blind, non ...

https://www.researchgate.net/publication/308946287_A_randomised...

the provision of analgesia in the first 24 h after total hip arthroplasty. No increase in side-effects was noted in the ... Fascia Iliaca Blocks and Non-Physician ... intraoperative opioid use ...