

World Journal of *Clinical Cases*

World J Clin Cases 2019 April 26; 7(8): 908-1005





REVIEW

- 908** Exosomes in esophageal cancer: A review on tumorigenesis, diagnosis and therapeutic potential
Su LL, Chang XJ, Zhou HD, Hou LB, Xue XY

MINIREVIEWS

- 917** Food additives can act as triggering factors in celiac disease: Current knowledge based on a critical review of the literature
Mancuso C, Barisani D

ORIGINAL ARTICLE

Retrospective Study

- 928** Optimal use of fielder XT guidewire enhances the success rate of chronic total occlusion percutaneous coronary intervention
Wang QC, Lin HR, Han Y, Dong H, Xu K, Guan SY, Chen ZH, Hao HX, Bin JP, Liao YL, Jing QM

Observational Study

- 940** Association between ventricular repolarization variables and cardiac diastolic function: A cross-sectional study of a healthy Chinese population
Li ZD, Bai XJ, Han LL, Han W, Sun XF, Chen XM

CASE REPORT

- 951** Non-invasive home lung impedance monitoring in early post-acute heart failure discharge: Three case reports
Lycholip E, Palevičiūtė E, Aamodt IT, Hellesø R, Lie I, Strömberg A, Jaarsma T, Čelutkienė J
- 961** Bilateral adrenocortical adenomas causing adrenocorticotrophic hormone-independent Cushing's syndrome: A case report and review of the literature
Gu YL, Gu WJ, Dou JT, Lv ZH, Li J, Zhang SC, Yang GQ, Guo QH, Ba JM, Zang L, Jin N, Du J, Pei Y, Mu YM
- 972** Two case reports and literature review for hepatic epithelioid angiomyolipoma: Pitfall of misdiagnosis
Mao JX, Teng F, Liu C, Yuan H, Sun KY, Zou Y, Dong JY, Ji JS, Dong JF, Fu H, Ding GS, Guo WY
- 984** Diagnosis of follicular lymphoma by laparoscopy: A case report
Wei C, Xiong F, Yu ZC, Li DF, Luo MH, Liu TT, Li YX, Zhang DG, Xu ZL, Jin HT, Tang Q, Wang LS, Wang JY, Yao J
- 992** Extranodal natural killer/T-cell lymphoma (nasal type) presenting as a perianal abscess: A case report
Liu YN, Zhu Y, Tan JJ, Shen GS, Huang SL, Zhou CG, Huangfu SH, Zhang R, Huang XB, Wang L, Zhang Q, Jiang B

- 1001** Radiologic features of Castleman's disease involving the renal sinus: A case report and review of the literature

Guo XW, Jia XD, Shen SS, Ji H, Chen YM, Du Q, Zhang SQ

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Preet M Chaudhary, MD, PhD, Professor, Department of Medicine, Jane Anne Nohl Division of Hematology and Center for the Study of Blood Diseases, University of Southern California, Los Angeles, CA 90033, United States

AIMS AND SCOPE

World Journal of Clinical Cases (*World J Clin Cases*, *WJCC*, online ISSN 2307-8960, DOI: 10.12998) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

The primary task of *WJCC* is to rapidly publish high-quality Case Report, Clinical Management, Editorial, Field of Vision, Frontier, Medical Ethics, Original Articles, Meta-Analysis, Minireviews, and Review, in the fields of allergy, anesthesiology, cardiac medicine, clinical genetics, clinical neurology, critical care, dentistry, dermatology, emergency medicine, endocrinology, family medicine, gastroenterology and hepatology, *etc.*

INDEXING/ABSTRACTING

The *WJCC* is now indexed in PubMed, PubMed Central, Science Citation Index Expanded (also known as SciSearch®), and Journal Citation Reports/Science Edition. The 2018 Edition of Journal Citation Reports cites the 2017 impact factor for *WJCC* as 1.931 (5-year impact factor: N/A), ranking *WJCC* as 60 among 154 journals in Medicine, General and Internal (quartile in category Q2).

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: Yun-Xiaojuan Wu Proofing Editorial Office Director: Jin-Lei Wang

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Semimonthly

EDITORS-IN-CHIEF

Dennis A Bloomfield, Sandro Vento

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

EDITORIAL OFFICE

Jin-Lei Wang, Director

PUBLICATION DATE

April 26, 2019

COPYRIGHT

© 2019 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Observational Study

Association between ventricular repolarization variables and cardiac diastolic function: A cross-sectional study of a healthy Chinese population

Zhi-Dan Li, Xiao-Juan Bai, Lu-Lu Han, Wen Han, Xue-Feng Sun, Xiang-Mei Chen

ORCID number: Zhi-Dan Li (0000-0001-5558-2800); Xiao-Juan Bai (0000-0002-5919-7173); Lu-Lu Han (0000-0001-8021-8998); Wen Han (0000-0003-4410-3362); Xue-Feng Sun (0000-0003-1160-4371); Xiang-Mei Chen (0000-0002-8889-6360).

Author contributions: Li ZD analyzed the data and drafted the manuscript; Bai XJ designed the manuscript and approved the manuscript for final submission; Han LL and Han W measured some important parameters; Sun XF and Chen XM participated in the coordination of the study; all authors read and approved the final manuscript.

Supported by the National Basic Research Program of China, No. 973-Program #2007CB507405.

Institutional review board statement: The study was approved by the ethics committee of General Hospital of Chinese People's Liberation Army (Beijing, China).

Informed consent statement: All patients gave informed consent.

Conflict-of-interest statement: The authors declare no conflict of interest.

STROBE statement: The authors have read the STROBE Statement-checklist of items, and the manuscript was prepared and revised according to the STROBE Statement-checklist of items.

Zhi-Dan Li, Xiao-Juan Bai, Lu-Lu Han, Wen Han, Department of Gerontology and Geriatrics, Shengjing Hospital of China Medical University, Shenyang 110004, Liaoning Province, China

Xue-Feng Sun, Xiang-Mei Chen, Department of Kidney, General Hospital of Chinese People's Liberation Army, Beijing 100853, China

Corresponding author: Xiao-Juan Bai, PhD, Professor, Department of Gerontology and Geriatrics, Shengjing Hospital of China Medical University, 36 Sanhao Street, Shenyang 110004, Liaoning Province, China. xiaojuan_bai@126.com

Telephone: +86-24-62043437

Fax: +86-24-62043437

Abstract

BACKGROUND

Diastolic electromechanical couple, a well-described phenomenon in symptomatic heart failure, has not been well studied in healthy people. We hypothesized that ventricular repolarization variables, such as the QT interval, Tpeak-to-Tend (Tpe) interval and Tpe/QT ratio, are associated with cardiac diastolic function in the healthy Chinese population.

AIM

To assess the relationship between ventricular repolarization variables and cardiac diastolic function in apparently healthy Chinese individuals.

METHODS

This was a community-based cross-sectional study conducted in Shenyang, China. A total of 414 healthy subjects aged 35-91 years were enrolled. All subjects underwent standard 12-lead electrocardiography (ECG) and comprehensive echocardiography. ECG enabled the measurement of QT and Tpe intervals and Tpe/QT ratio. echocardiographic parameters, such as the ratio of mitral early diastolic inflow velocity (E) and late diastolic inflow velocity (A), E-wave deceleration time, left atrial volume (LAV) and LAV index, were measured to assess diastolic function. $E/A < 0.75$ was considered to indicate reduced diastolic function. ECG and echocardiography results were analyzed separately and in a blinded fashion. Correlation and regression analyses were applied to determine associations.

RESULTS

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Received: December 26, 2018

Peer-review started: December 27, 2018

First decision: March 10, 2019

Revised: March 28, 2019

Accepted: April 9, 2019

Article in press: April 9, 2019

Published online: April 26, 2019

P-Reviewer: Barik R, Karatza AA, Kharlamov AN

S-Editor: Ji FF

L-Editor: A

E-Editor: Wu YXJ



Ventricular repolarization variables, such as the QTc interval (393.59 ± 26.74 vs 403.86 ± 33.56 ; $P < 0.001$), Tpe interval (72.68 ± 12.41 vs 77.26 ± 17.86 ; $P < 0.01$), Tpe_c interval (76.36 ± 13.53 vs 83.32 ± 21.25 ; $P < 0.001$) and Tpe/QT ratio (0.19 ± 0.03 vs 0.20 ± 0.04 ; $P < 0.01$), were significantly different between the normal diastolic function group and the reduced diastolic function group. Significant associations were found between repolarization variables and diastolic function. After adjusting for all other possible confounders, the QTc and Tpe_c intervals were significantly associated with the E/A ratio ($P = 0.008$; $P = 0.010$). In men, the QTc interval was associated with abnormal diastolic function, and compared to the third QTc tertile, in the second QTc tertile, the odds ratio was 0.257 (95%CI: 0.102–0.649; $P = 0.004$).

CONCLUSION

Repolarization variables are associated with cardiac diastolic function even in healthy people. Moderate levels of the QTc interval exert a protective effect on diastolic dysfunction in men.

Key words: QT interval; Tpeak-to-Tend interval; Diastolic dysfunction; Ventricular repolarization; Electrocardiography

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: A community-based cross-sectional study conducted to assess the relationship between ventricular repolarization variables and cardiac diastolic function in apparently healthy Chinese individuals. We observed an independent association between ventricular repolarization variables and cardiac diastolic function parameters, and moderate levels of the QTc interval tend to have a protective effect on diastolic dysfunction in men. Electromechanical coupling may represent a relationship between heterogeneity of repolarization and abnormal myocardial mechanics.

Citation: Li ZD, Bai XJ, Han LL, Han W, Sun XF, Chen XM. Association between ventricular repolarization variables and cardiac diastolic function: A cross-sectional study of a healthy Chinese population. *World J Clin Cases* 2019; 7(8): 940-950

URL: <https://www.wjnet.com/2307-8960/full/v7/i8/940.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v7.i8.940>

INTRODUCTION

Heart failure (HF) is a complex and increasingly common syndrome that affects over 23 million patients worldwide^[1]. Approximately half of patients with HF present with diastolic dysfunction, and asymptomatic diastolic dysfunction is present in 21% of the population^[2]. Asymptomatic diastolic dysfunction may be present for a significant period of time before developing into symptomatic HF. The latency between dysfunction and symptoms represents the best time for using effective diagnostics and therapies^[3]. Therefore, the mechanisms underlying diastolic dysfunction need to be more extensively studied.

Growing evidence demonstrates that the pathologies of diastolic dysfunction and systolic dysfunction are different. Some studies have reported that electrical repolarization abnormalities are associated with declining diastolic function, suggesting that diastolic electromechanical coupling represents a unifier linking diastolic dysfunction, calcium handling, and repolarization abnormalities with the development of symptomatic HF^[4]. The QT interval is the most commonly used parameter in the electrocardiographic (ECG) assessment of repolarization by physicians in clinical practice. The Tpeak-to-Tend (Tpe) interval, defined as the time interval between the peak and the end of the T wave, is proposed as a readily available ECG measurement of the dispersion of ventricular repolarization. The Tpe/QT ratio is used to predict cardiac arrhythmias. Previous studies have evaluated the QT and Tpe intervals as potential mechanistic contributors to diastolic dysfunction in specific populations^[5-8]. Wilcox *et al*^[5] found QTc prolongation was independently associated with diastolic dysfunction in patients with clinical suspicion of HF. Another study by Khan *et al*^[6] confirmed that a prolonged QTc interval was a

useful tool for predicting diastolic dysfunction. In addition, Sauer *et al*^[7] showed that increased Tpe interval was associated with both resting and exercise-induced diastolic dysfunction. However, the subjects of the studies above were all patients with risk factors of HF, and risk factors such as hypertension and diabetes mellitus may influence both heart structure and heart function and are powerful potential confounders of electromechanical coupling. Thus, the aim of this study was to assess the relationship between ventricular repolarization variables, such as the QT interval, Tpe interval and Tpe/QT ratio, and diastolic function in an apparently healthy Chinese population and determine whether measuring repolarization variables may provide information valuable for predicting cardiac diastolic function in healthy people.

MATERIALS AND METHODS

Study population

The subjects who participated in this community-based cross-sectional study were recruited from a healthy Han Chinese population between 2007 and 2008. There were the following register criteria: (1) Age older than 35 years; (2) Being healthy by self-evaluation; (3) Having the ability to care for themselves and perform activities associated with daily living independently and without difficulty; (4) Having the ability to provide informed consent and self-reported data. Subjects with cardiovascular disease, hypertension [defined as a systolic blood pressure (SBP) ≥ 140 mmHg or a diastolic blood pressure (DBP) ≥ 90 mmHg and/or an antihypertensive medication requirement], diabetes (defined as a fasting plasma glucose > 7.0 mmol/L or an insulin or oral hypoglycemic medication requirement), and other chronic diseases were excluded from the study. The subjects who provided the informed consent conducted physical examination or laboratory tests. After excluding persons with an abnormal physical examination or laboratory results, 414 healthy subjects (186 men and 228 women) were included in the study. This study was approved by the Ethics Committee of China Medical University.

Clinical measurements and laboratory tests

The participants underwent a clinical examination and completed a detailed questionnaire at the time of enrollment. Before they were examined, the subjects rested for 10-15 min in a temperature-controlled environment. Physical parameters, including height and weight, were measured with a digital scale while the participants were wearing light clothing and no shoes on a digital scale. Blood pressure was measured using a manual stethoscope and a sphygmomanometer with an adjustable cuff. Two measurements were performed 2 min apart, and the average of the two measurements was calculated. Body mass index (BMI) was calculated as the mass in kilograms divided by height in meters squared, and body surface area (BSA) was calculated according to the following formula: $BSA (m^2) = 0.0061 \times \text{height (cm)} + 0.0128 \times \text{weight (kg)} - 0.1529$.

Blood samples were collected from the subjects between 8:00 and 9:00 am after the subjects had fasted for at least 10 h overnight. Blood biochemical parameters, including triglycerides (TG), total cholesterol (TC), high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein (LDL-C), fasting blood glucose (FBG) and serum creatinine (SCr) levels, were assayed on-site at the medical laboratory of the study center.

ECG

All subjects underwent a 12-lead ECG recorded at a paper speed of 25 mm/sec and a voltage of 10 mm/mV by a standard ECG system. For ECG analysis, we performed manual measurements of the values with a digital caliper using a computer program. The QT interval was measured between the QRS onset and the end of the T wave, and the Tpe interval was measured from the peak of the T wave to the end of the T wave. The QT interval and the Tpe interval were corrected separately for heart rate using the Bazett formula^[9]. We also calculated the Tpe/QT and Tpe/QTc ratios. All measurements were performed in lead V5^[10]. In cases in which lead V5 could not be used for the analysis, leads V4 and V6 (in that order) were utilized. ECG measurements were performed by a single trained reader who was blinded to the echocardiographic results.

Echocardiography

All subjects underwent a complete M-mode, two-dimensional and pulsed-wave (PW) Doppler echocardiographic examination using a Philips iE33 Ultrasound System, and all echocardiographic measurements were obtained according to published

guidelines^[11,12]. PW Doppler was performed in the apical 4-chamber view to obtain data pertaining to mitral inflow velocities and time intervals. The primary measurements performed during this procedure included measurements of mitral early diastolic inflow velocity (E), late diastolic inflow velocity (A), the E/A ratio, and E-wave deceleration time (DT). We used the apical four-chamber views to measure left atrial anterior-posterior diameter (LA-AP-D), left atrial medial-lateral diameter (LA-ML-D) and left atrial superior-inferior diameter (LA-SI-D). The left atrial volume (LAV) was computed by the equation $4\pi/3(LA-SI-D/2)(LA-AP-D/2)(LA-ML-D/2)$ according to the ellipsoid model, and the left atrial volume index (LAVI) was calculated as the LAV in milliliters divided by the BSA in meters squared. According to the American Society of Echocardiography guidelines^[12], in this study, $E/A < 0.75$ was considered to be reduced diastolic function.

Statistical analysis

The Kolmogorov-Smirnov test was used for all analyses of normally distributed data. All continuous data were normally distributed and were expressed as the means \pm SDs. The differences in these variables between different groups were examined using the independent-samples t-test. Categorical variables were expressed as the number {(percentile) $n(\%)$ }, and comparative analyses of different groups were performed using the chi-square (χ^2) test. Simple correlation analyses were performed by calculating Pearson's coefficients for the relationships between two variables. Multiple linear regression analyses were performed to adjust for possible confounding variables. The following three models were used: An un-adjusted model, a model adjusted only for age, and a model adjusted for age, BMI, SBP, DBP, TG, TC, HDL-C, LDL-C, FBG and SCr. Binary logistic regression was reformed to investigate the associations between abnormal diastolic function and the levels of QTc and Tpe. The subjects were categorized into tertiles (1-3) for QTc and Tpe, with tertile 3 showing the highest levels of QTc and Tpe. The following two models were applied: A model adjusting for age and a model adjusting for age, BMI, SBP, DBP, TG, TC, HDL-C, LDL-C, FBG and SCr. All statistical analyses were performed using SPSS 19.0 statistical software (SPSS, Chicago, IL, United States), and a P value < 0.05 was considered statistically significant.

RESULTS

Clinical, electrocardiographic and echocardiographic characteristics

A total of 414 subjects (186 men) were assessed in this study, and the mean age was 55 ± 14 years old. Subjects were divided into two groups based on their E/A values (one group with $E/A \geq 0.75$ and the other with $E/A < 0.75$). The differences in the clinical, ECG and echocardiographic characteristics between the groups with normal and reduced E/A ratios are shown in Table 1. Age, BMI, SBP, DBP and SCr were significantly higher in subjects with a reduced E/A ratio ($P < 0.01$). Furthermore, there were several ECG-related differences between the two groups: The RR interval, QTc interval, Tpe interval, Tpe_c interval and Tpe/QT ratio were higher in those with $E/A < 0.75$ than in those with $E/A \geq 0.75$ ($P < 0.01$). All echocardiographic parameters were significantly different between the two groups ($P < 0.05$).

Correlation analysis between electrocardiographic repolarization variables and other variables

Table 2 shows the Pearson correlation coefficients for the relationships between ECG repolarization variables and other variables. All the repolarization variables except the Tpe/QTc ratio were significantly associated with age ($P < 0.05$) and all except the QT interval were associated with BMI ($P < 0.05$). Moreover, the QTc interval was significantly associated with SBP, TC and LDL-C ($P < 0.05$). The Tpe interval was significantly associated with HDL-C ($P < 0.05$). The Tpe_c interval was significantly associated with HDL-C and LDL-C ($P < 0.05$). The Tpe/QT ratio was significantly associated with HDL-C, LDL-C and SCr ($P < 0.05$). The Tpe/QTc ratio was significantly associated with HDL-C and SCr ($P < 0.01$). With regard to the echocardiographic diastolic parameters, the E/A ratio was negatively and significantly correlated with QTc, Tpe and Tpe_c intervals and Tpe/QT ratio ($P < 0.01$). E was negatively correlated with the Tpe_c interval and the Tpe/QT ratio ($P < 0.05$). A was positively correlated with all the repolarization variables except the Tpe/QTc ratio ($P < 0.01$). DT was positively correlated with the QT interval and the Tpe/QTc ratio ($P < 0.05$). LAV was positively correlated with all the repolarization variables except the Tpe/QT ratio ($P < 0.05$). LAVI was positively correlated with QTc and Tpe intervals ($P < 0.01$).

Table 1 Clinical, electrocardiography and echocardiographic characteristics based on their E/A ratio

	All subjects (n = 414)	E/A ≥ 0.75 (n = 300)	E/A < 0.75 (n = 114)	P-value
Age (yr)	55.21 ± 14.18	55.25 ± 13.20	69.62 ± 11.05	< 0.001
Male n (%)	186 (44.9%)	124 (41.3%)	62 (55.4%)	0.017
BMI (kg/m ²)	23.68 ± 3.00	23.41 ± 2.93	24.37 ± 3.08	0.004
SBP (mmHg)	125.64 ± 12.69	123.05 ± 12.68	132.43 ± 9.93	< 0.001
DBP (mmHg)	75.97 ± 8.84	74.92 ± 8.97	78.73 ± 7.87	< 0.001
TG (mmol/L)	1.38 ± 1.10	1.39 ± 1.18	1.36 ± 0.84	0.840
TC (mmol/L)	5.23 ± 1.05	5.22 ± 1.08	5.26 ± 0.98	0.722
HDL-C (mmol/L)	1.64 ± 0.41	1.65 ± 0.40	1.62 ± 0.44	0.440
LDL-C (mmol/L)	2.90 ± 0.75	2.88 ± 0.76	2.95 ± 0.70	0.448
FBG (mmol/L)	5.55 ± 0.49	5.54 ± 0.48	5.60 ± 0.53	0.248
SCr (μmol/L)	64.37 ± 14.46	62.72 ± 13.23	68.71 ± 16.58	< 0.001
PR interval (ms)	157.08 ± 23.73	156.36 ± 24.18	158.97 ± 22.50	0.302
RR interval (ms)	905.53 ± 126.29	915.40 ± 118.32	879.56 ± 142.52	0.010
QT interval (ms)	372.81 ± 30.07	371.45 ± 29.49	376.39 ± 31.40	0.136
QTc interval (ms)	393.52 ± 29.44	393.59 ± 26.74	403.86 ± 33.56	< 0.001
Tpe interval (ms)	73.94 ± 14.25	72.68 ± 12.41	77.26 ± 17.86	0.003
Tpe _c interval (ms)	78.27 ± 16.30	76.36 ± 13.53	83.32 ± 21.25	< 0.001
Tpe/QT	0.20 ± 0.03	0.19 ± 0.03	0.20 ± 0.04	0.004
Tpe/QTc	0.19 ± 0.03	0.19 ± 0.03	0.19 ± 0.03	0.160
E (cm/s)	74.36 ± 17.83	80.27 ± 16.20	58.82 ± 11.49	< 0.001
A (cm/s)	75.07 ± 19.21	68.75 ± 16.70	91.69 ± 15.07	< 0.001
E/A	1.07 ± 0.41	1.23 ± 0.36	0.64 ± 0.08	< 0.001
MV-DT (ms)	162.78 ± 35.15	159.43 ± 31.14	171.60 ± 42.92	0.002
LAV (mL)	33.16 ± 9.73	32.30 ± 9.70	35.42 ± 9.48	0.003
LAVI (mL/m ²)	19.83 ± 5.31	19.38 ± 5.29	21.01 ± 5.22	0.005
LVEF (%)	62.54 ± 4.71	62.86 ± 4.47	61.68 ± 5.21	0.023

BMI: Body mass index; SBP: Systolic blood pressure; DBP: Diastolic blood pressure; TG: Triglyceride; TC: Total cholesterol; HDL-C: High-density lipoprotein cholesterol; LDL-C: Low-density lipoprotein cholesterol; FBG: Fasting blood glucose; SCr: Serum creatinine; E: Early diastolic inflow velocity; A: Late diastolic inflow velocity; DT: E-wave deceleration time; LAV: Left atrial volume; LAVI: Left atrial maximum volume index; LVEF: Left ventricular ejection fraction.

Associations between electrocardiographic repolarization variables and echocardiographic diastolic function parameters

Table 3 shows the results of the multiple linear regression analyses, in which echocardiographic diastolic function parameters served as the dependent variables, and ECG repolarization variables served as the independent variables. After adjusting for all other possible confounders, A and E/A ratio were significantly associated with QTc and Tpe intervals and the Tpe/QT ratio, and A was also significantly associated with the Tpe interval. LAV and LAVI showed significant and independent associations with the QTc interval, and LAV was also significantly associated with the QT interval. DT was significantly associated with the QT interval.

Table 4 shows the binary logistic regression analyses performed using reduced diastolic function as the dependent variable and cut-off values of the QTc and Tpe_c interval as the independent variables. In men, the odds ratio in subjects with reduced diastolic function was 2.715 (95%CI: 1.356–5.432; *P* < 0.001) for longer QTc interval after adjustment for age, compared to the subjects in normal QTc interval. This association was still significant after adjustment for all other variables, and the odds ratio was 2.567 (95%CI: 1.227–5.370; *P* = 0.012). In women, the QTc interval was not correlated with reduced diastolic function. With regard to the Tpe_c interval, in men, compared to the third Tpe_c tertile, in the Tpe_c second tertile, the odds ratio approached statistical significance (0.423; 95%CI: 0.175–1.022; *P* = 0.056) after adjustment for all other variables.

Table 2 Correlation analysis between electrocardiography repolarization variables and other variables

	QT (ms)	QTc (ms)	Tpe (ms)	Tpe _c (ms)	Tpe/QT	Tpe/QTc
Age (yr)	0.223 ^b	0.281 ^b	0.167 ^b	0.169 ^b	0.109 ^a	0.090
BMI (kg/m ²)	0.073	0.109 ^a	0.119 ^a	0.124 ^a	0.120 ^a	0.100 ^a
SBP (mmHg)	0.082	0.167 ^b	0.059	0.089	0.049	0.004
DBP (mmHg)	-0.031	0.029	-0.034	-0.008	-0.005	-0.036
TG (mmol/L)	-0.02	0.028	0.058	0.073	0.086	0.058
TC (mmol/L)	0.009	0.115 ^a	0.054	0.094	0.071	0.020
HDL-C (mmol/L)	-0.061	-0.067	-0.132 ^b	-0.122 ^a	-0.140 ^b	-0.130 ^b
LDL-C (mmol/L)	-0.046	0.155 ^b	0.088	0.127 ^a	0.098 ^a	0.042
FBG (mmol/L)	-0/089	0.080	0.015	0.074	0.081	0.001
SCr (μmol/L)	-0/013	-0.055	0.089	0.068	0.123 ^a	0.146 ^b
E (cm/s)	0.051	-0.093	-0.072	-0.121 ^a	-0.138 ^b	-0.060
A (cm/s)	0.152 ^b	0.324 ^b	0.189 ^b	0.237 ^b	0.157 ^b	0.079
E/A	-0.070	-0.288 ^b	-0.155 ^b	-0.223 ^b	-0.175 ^b	-0.067
DT (ms)	0.244 ^b	-0.028	0.073	-0.047	-0.039	0.102 ^a
LAV (mL)	0.312 ^b	0.209 ^b	0.157 ^b	0.099 ^a	0.038	0.100 ^a
LAVI (mL/m ²)	0.354	0.255 ^b	0.148 ^b	0.091	0.002	0.062

P-values are from analysis of variance:

^aSignificant difference ($P < 0.05$);

^bSignificant difference ($P < 0.01$). BMI: Body mass index; SBP: Systolic blood pressure; DBP: Diastolic blood pressure; TG: Triglyceride; TC: Total cholesterol; HDL-C: High-density lipoprotein cholesterol; LDL-C: Low-density lipoprotein cholesterol; FBG: Fasting blood glucose; SCr: Serum creatinine; E: Early diastolic inflow velocity; A: Late diastolic inflow velocity; DT: E-wave deceleration time; LAV: Left atrial volume; LAVI: Left atrial maximum volume index.

DISCUSSION

The most important findings of the present study are the associations between ventricular repolarization variables and cardiac diastolic function in a population of healthy adults. These relationships remained significant even after correction for other potential confounders.

The ECG QT interval is one of the ventricular repolarization variables most commonly used by physicians in clinical practice. Previous studies have demonstrated the existence of a relationship between a prolonged QT interval and abnormal myocardial mechanical function among patients with inherited long QT syndrome (LQTS), which has historically been considered a purely electrical disease^[13-15]. In recent years, clinicians have become increasingly interested in the association between the QT interval and echocardiographic parameters representing diastolic function in patients with suspected HF, hypertension, and diabetes mellitus and have obtained data supporting the idea that there is a correlation between diastolic dysfunction and the QT interval^[5,8,16]. However, diseases such as hypertension and diabetes mellitus influence both heart structure and heart function and are powerful potential confounders of electromechanical coupling measurements. The present community-based study included apparently healthy subjects selected from a Chinese population without hypertension, diabetes, cardiovascular disease, or other chronic diseases and thus excluded potential confounders. We found an independent linear association between the QTc interval and echocardiographic diastolic parameters. Moreover, this study also demonstrates that moderate levels of QTc exert a protective effect on diastolic dysfunction in healthy men. To the best of our knowledge, this is the first study to demonstrate an association between QTc interval levels and diastolic function in a healthy population. Possible explanations for this finding include the possibility that the link between electrical repolarization and diastolic mechanics may be commonly mediated by the effects of calcium handling^[4]. Previous studies have also described a U-shaped association between the QTc interval and risk of death^[17,18]. Therefore, it is biologically plausible that there is an optimum for the dispersion of ventricular repolarization and that the risk of diastolic dysfunction is increased at both ends of the spectrum. However, sex-related differences in the association between QTc interval levels and diastolic function remain to be explored.

Table 3 Relationship between electrocardiography repolarization variables and echocardiographic diastolic parameters using a stepwise multiple regression model

	Model 1		Model 2		Model 3	
	Beta	P-value	Beta	P-value	Beta	P-value
QT interval (ms)						
A (cm/s)	0.152	0.002	0.033	0.435	0.028	0.504
DT (ms)	0.244	< 0.001	0.194	< 0.001	0.182	< 0.001
LAV (mL)	0.312	< 0.001	0.271	< 0.001	0.248	< 0.001
QTc interval (ms)						
A (cm/s)	0.324	< 0.001	0.186	< 0.001	0.178	< 0.001
E/A	-0.288	< 0.001	-0.131	0.001	-0.105	0.008
LAV (mL)	0.209	< 0.001	0.152	0.002	0.105	0.019
LAVI (mL/m ²)	0.255	< 0.001	0.181	< 0.001	0.142	0.003
Tpe (ms)						
A (cm/s)	0.189	< 0.001	0.102	0.015	0.099	0.016
E/A	-0.155	0.002	-0.057	0.153	-0.035	0.355
LAV (mL)	0.157	0.001	0.119	0.014	0.167	0.120
LAVI (mL/m ²)	0.148	0.003	0.098	0.039	0.080	0.086
Tpe _c (ms)						
E (cm/s)	-0.121	0.014	-0.062	0.185	-0.032	0.489
A (cm/s)	0.237	< 0.001	0.150	< 0.001	0.143	< 0.001
E/A	-2.223	< 0.001	-0.126	0.002	-0.098	0.010
LAV (mL)	0.099	0.045	0.059	0.233	0.015	0.749
Tpe/QT						
E (cm/s)	-0.138	0.005	-0.100	0.030	-0.066	0.152
A (cm/s)	0.157	0.001	0.099	0.017	0.097	0.018
E/A	-0.175	< 0.001	-0.111	0.005	-0.083	0.028
Tpe/QTc						
DT (ms)	0.102	0.038	0.078	0.101	0.072	0.139
LAV (mL)	0.100	0.041	0.079	0.100	0.037	0.388

E: Early diastolic inflow velocity; A: Late diastolic inflow velocity; DT: E-wave deceleration time; LAV: Left atrial volume; LAVI: Left atrial maximum volume index. Model 1: Unadjusted model; Model 2: Adjusted for age; Model 3: Fully adjusted for age, gender, body mass index, systolic blood pressure, diastolic blood pressure, triglyceride, total cholesterol, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, fasting blood glucose and serum creatinine. Standardized coefficients and *P*-values were the outcomes of the regression analyses.

Tpe is the interval between the peak of the T wave and the end of the T wave. The Tpe interval is usually viewed as representative of the difference in repolarization times between subendocardial and subepicardial myocardial cells and has been proposed to reflect the transmural cardiac repolarization expressed through surface 12 ECG^[4]. Multiple recent studies have demonstrated that Tpe interval plays an important role as a potential ECG biomarker for predicting arrhythmia risk and cardiovascular death^[19-21]. Furthermore, some studies have evaluated the Tpe interval as a potential mechanism that contributes to mechanical dysfunction in patients with overt or suspected HF^[7,22]. In the present population-based study, we found a significant linear association between ECG Tpe and Tpe_c intervals and the echocardiographic E/A ratio and A wave. These findings add to the growing literature supporting the notion that electromechanical coupling of dispersion of repolarization is a potential mechanism of diastolic dysfunction. It is interesting to note that the Tpe_c interval was more strongly correlated with diastolic function parameters in this study, compared to the Tpe interval. The basis for this result is unclear and is beyond the scope of the current study. In the present study, we found that in men, compared to the third Tpe_c tertile, in the Tpe_c second tertile, the odds ratio approached statistical significance (0.423; 95%CI: 0.175–1.022; *P* = 0.056) after adjustment for all other variables. Moderate Tpe_c levels tend to have a protective effect on diastolic dysfunction in men. A large population study found U-shaped associations between the Tpe interval and the risks of all-cause and cardiovascular

Table 4 Association of E/A ratio within tertiles of QTc and Tpe_c using binary logistic regression model

	Tertile 1				Tertile 2				Tertile 3
	Model 1		Model 2		Model 1		Model 2		OR
	OR (95%CI)	<i>P</i> -value	OR (95%CI)	<i>P</i> -value	OR (95%CI)	<i>P</i> -value	OR (95%CI)	<i>P</i> -value	
Tertiles of QTc									
Men (<i>n</i> = 186)	0.604 (0.272-1.342)	0.216	0.763 (0.317-1.836)	0.545	0.210 (0.087-0.506)	0.001	0.257 (0.102-0.649)	0.004	1 (ref)
Women (<i>n</i> = 228)	0.527 (0.176-1.579)	0.253	0.982 (0.284-3.396)	0.977	1.230 (0.539-2.806)	0.624	1.480 (0.574-3.815)	0.417	1 (ref)
Tertiles of Tpe _c									
Men (<i>n</i> = 186)	0.890 (0.397-1.996)	0.777	0.980 (0.404-2.372)	0.963	0.465 (0.204-1.061)	0.069	0.423 (0.175-1.022)	0.056	1 (ref)
Women (<i>n</i> = 228)	0.483 (0.188-1.242)	0.131	0.477 (0.162-1.404)	0.179	0.487 (0.204-1.160)	0.104	0.532 (0.200-1.416)	0.206	1 (ref)

Model 1: Adjusted for age; Model 2: Fully adjusted for age, gender, body mass index, systolic blood pressure, diastolic blood pressure, triglyceride, total cholesterol, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, fasting blood glucose and serum creatinine. OR values and *P*-value were the outcome of binary logistic regression using abnormal diastolic function as the dependent variable.

mortality, atrial fibrillation, and HF^[19]. Therefore, similar to the QTc interval, there may be an optimum for the dispersion of cardiac repolarization, and the risk of cardiovascular outcomes may be increased at both ends of the spectrum. Further investigations are necessary to confirm and improve the present findings.

The Tpe/QT ratio is a novel ventricular repolarization variable that is used to predict cardiac arrhythmias^[23]. Some studies have suggested the applicability of Tpe/QT ratio as a potentially important index of arrhythmogenesis in congenital and acquired channelopathies^[23,24]. Furthermore, some studies have evaluated the Tpe/QT ratio as a potential ECG biomarker for predicting arrhythmia risk and cardiovascular death^[25-27]. In the present study, we found an inverse linear association between the Tpe/QT ratio and the E/A ratio. However, the Tpe/QTc ratio was not associated with any echocardiographic diastolic function parameters after adjustment for several important potential confounders. The basis for this finding is unclear, and additional studies are required.

Our results should not be interpreted to suggest that repolarization variables measured on surface ECG can be used to screen for diastolic dysfunction on echocardiography. Furthermore, our current understanding of the role of ventricular repolarization in the development or progression of HF is limited. Rather, our findings provide evidence showing that ventricular repolarization parameters may be used as markers of asymptomatic mild diastolic dysfunction. Epidemiological studies suggest that there is a latent phase during which diastolic dysfunction is present and progresses in severity before the symptoms of HF arise^[28]. This asymptomatic phase represents a potential time to intervene and thereby prevent symptomatic HF. The latency between dysfunction and symptoms represents the best time for using effective diagnostics and therapies. Identifying a pharmacological intervention to restore repolarization to a more normal state may be a novel target for therapy.

The present study had several limitations that require discussion. First, this research is part of a cross-sectional study, and it is therefore difficult to demonstrate cause-effect relationships between ventricular repolarization variables and cardiac diastolic function. A longitudinally designed study is required to confirm the current findings. Second, the subjects included in this study were recruited from communities in northern China. Therefore, the findings in the study are not representative of the general population. Third, a relatively low number of subjects were included in the study, and the proportion of women (55.1%) was higher than that of men (44.9%). Fourth, we chose to report the QT and Tpe intervals corrected for heart rate (QTc and Tpe_c) using the Bazett formula. The Bazett formula undercorrects the QT interval at lower heart rates and overcorrects it at higher heart rates^[29]. However, the Bazett formula is the formula that is most often used in research or clinical practice. Fifth, several important parameters, including LV mass and QRS durations, were not measured in this study but may affect ventricular repolarization variables^[30]. Further studies are needed to confirm and improve upon the findings of the present study.

In conclusion, this study reveals that even in healthy people, ventricular repolarization variables are linearly associated with cardiac diastolic function parameters. Furthermore, moderate levels of the QTc interval tend to have a protective effect on diastolic dysfunction in men. Electromechanical coupling may represent a relationship between heterogeneity of repolarization and abnormal

myocardial mechanics. A longitudinal study should be performed in the future to confirm the findings of the current study.

ARTICLE HIGHLIGHTS

Research background

Asymptomatic diastolic dysfunction present for a significant period of time before developing into symptomatic heart failure (HF), and diastolic electromechanical coupling may represent a unifier linking diastolic dysfunction, calcium handling, and repolarization abnormalities with the development of symptomatic HF. We assessed the relationship between ventricular repolarization variables, such as the QT interval, Tpe interval and Tpe/QT ratio, and diastolic function in an apparently healthy Chinese population through a community-based cross-sectional study.

Research motivation

Some studies have reported that electrical repolarization abnormalities are associated with declining diastolic function in patients with suspected HF, hypertension, and diabetes mellitus. However, diseases such as hypertension and diabetes mellitus influence both heart structure and heart function and are powerful potential confounders of electromechanical coupling measurements. Therefore, this study hopes to determine whether measuring repolarization variables may provide information valuable for predicting cardiac diastolic function in healthy people.

Research objectives

The research objective of this study was to explore the relationship between ventricular repolarization variables and cardiac diastolic function in apparently healthy Chinese individuals.

Research methods

We retrospectively analyzed 414 healthy subjects aged 35-91 years who were enrolled between September 2007 and June 2008. All subjects underwent standard 12-lead electrocardiography (ECG) and comprehensive echocardiography. ECG and echocardiography results were analyzed separately and in a blinded fashion. Correlation and regression analyses were applied to determine associations. This study is a community-based cross-sectional study.

Research results

Our research found that even in healthy people, ventricular repolarization variables are associated with cardiac diastolic function parameters. Furthermore, moderate levels of the QTc interval tend to have a protective effect on diastolic dysfunction in men.

Research conclusions

Repolarization variables are associated with cardiac diastolic function even in healthy people. Electromechanical coupling may represent a relationship between heterogeneity of repolarization and abnormal myocardial mechanics.

Research perspectives

We observed an independent association between ventricular repolarization variables and cardiac diastolic function in a community-based cross-sectional study. However, a relatively low number of subjects were included in the study, and the cross-sectional study has its own shortcomings. A longitudinally designed study is required to confirm the current findings.

REFERENCES

- 1 **Lloyd-Jones D**, Adams RJ, Brown TM, Carnethon M, Dai S, De Simone G, Ferguson TB, Ford E, Furie K, Gillespie C, Go A, Greenlund K, Haase N, Hailpern S, Ho PM, Howard V, Kissela B, Kittner S, Lackland D, Lisabeth L, Marelli A, McDermott MM, Meigs J, Mozaffarian D, Mussolino M, Nichol G, Roger VL, Rosamond W, Sacco R, Sorlie P, Stafford R, Thom T, Wasserthiel-Smoller S, Wong ND, Wylie-Rosett J; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Executive summary: heart disease and stroke statistics--2010 update: a report from the American Heart Association. *Circulation* 2010; **121**: 948-954 [PMID: 20177011 DOI: 10.1161/CIRCULATIONAHA.109.192666]
- 2 **Jeong EM**, Dudley SC. Diastolic dysfunction. *Circ J* 2015; **79**: 470-477 [PMID: 25746522 DOI: 10.1253/circj.CJ-15-0064]
- 3 **Halley CM**, Houghtaling PL, Khalil MK, Thomas JD, Jaber WA. Mortality rate in patients with diastolic dysfunction and normal systolic function. *Arch Intern Med* 2011; **171**: 1082-1087 [PMID: 21709107 DOI: 10.1001/archinternmed.2011.244]
- 4 **Prenner SB**, Shah SJ, Goldberger JJ, Sauer AJ. Repolarization Heterogeneity: Beyond the QT Interval. *J Am Heart Assoc* 2016; **5**: pii: e003607 [PMID: 27130347 DOI: 10.1161/JAHA.116.003607]
- 5 **Wilcox JE**, Rosenberg J, Vallakati A, Gheorghiadu M, Shah SJ. Usefulness of electrocardiographic QT interval to predict left ventricular diastolic dysfunction. *Am J Cardiol* 2011; **108**: 1760-1766 [PMID: 21907948 DOI: 10.1016/j.amjcard.2011.07.050]
- 6 **Khan HS**, Iftikhar I, Khan Q. Validity of Electrocardiographic QT Interval in Predicting Left Ventricular Diastolic Dysfunction in Patients with Suspected Heart Failure. *J Coll Physicians Surg Pak* 2016; **26**: 353-

- 356 [PMID: 27225136]
- 7 **Sauer A**, Wilcox JE, Andrei AC, Passman R, Goldberger JJ, Shah SJ. Diastolic electromechanical coupling: association of the ECG T-peak to T-end interval with echocardiographic markers of diastolic dysfunction. *Circ Arrhythm Electrophysiol* 2012; **5**: 537-543 [PMID: 22467673 DOI: 10.1161/CIRCEP.111.969717]
- 8 **Jani Y**, Kamberi A, Xhunga S, Pocesta B, Ferati F, Lala D, Zeqiri A, Rexhepi A. The influence of type 2 diabetes and gender on ventricular repolarization dispersion in patients with sub-clinic left ventricular diastolic dysfunction. *Am J Cardiovasc Dis* 2015; **5**: 155-166 [PMID: 26550530]
- 9 **Luo S**, Michler K, Johnston P, Macfarlane PW. A comparison of commonly used QT correction formulae: the effect of heart rate on the QTc of normal ECGs. *J Electrocardiol* 2004; **37** Suppl: 81-90 [PMID: 15534815 DOI: 10.1016/j.jelectrocard.2004.08.030]
- 10 **Castro-Torres Y**, Carmona-Puerta R, Katholi RE. Ventricular repolarization markers for predicting malignant arrhythmias in clinical practice. *World J Clin Cases* 2015; **3**: 705-720 [PMID: 26301231 DOI: 10.12998/wjcc.v3.i8.705]
- 11 **Nagueh SF**, Smiseth OA, Appleton CP, Byrd BF, Dokainish H, Edvardsen T, Flachskampf FA, Gillebert TC, Klein AL, Lancellotti P, Marino P, Oh JK, Alexandru Popescu B, Waggoner AD; Houston, Texas; Oslo, Norway; Phoenix, Arizona; Nashville, Tennessee; Hamilton, Ontario, Canada; Uppsala, Sweden; Ghent and Liège, Belgium; Cleveland, Ohio; Novara, Italy; Rochester, Minnesota; Bucharest, Romania; and St. Louis, Missouri. Recommendations for the Evaluation of Left Ventricular Diastolic Function by Echocardiography: An Update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging* 2016; **17**: 1321-1360 [PMID: 27422899 DOI: 10.1093/ehjci/ewj082]
- 12 **Nagueh SF**, Appleton CP, Gillebert TC, Marino PN, Oh JK, Smiseth OA, Waggoner AD, Flachskampf FA, Pellikka PA, Evangelisa A. Recommendations for the evaluation of left ventricular diastolic function by echocardiography. *Eur J Echocardiogr* 2009; **10**: 165-193 [PMID: 19270053 DOI: 10.1093/ejehoccard/jep007]
- 13 **Leren IS**, Hasselberg NE, Saberniak J, Håland TF, Kongsgård E, Smiseth OA, Edvardsen T, Haugaa KH. Cardiac Mechanical Alterations and Genotype Specific Differences in Subjects With Long QT Syndrome. *JACC Cardiovasc Imaging* 2015; **8**: 501-510 [PMID: 25890583 DOI: 10.1016/j.jcmg.2014.12.023]
- 14 **Savoie C**, Klug D, Denjoy I, Ennezat PV, Le Tourneau T, Guicheney P, Kacet S. Tissue Doppler echocardiography in patients with long QT syndrome. *Eur J Echocardiogr* 2003; **4**: 209-213 [PMID: 12928025 DOI: 10.1016/s1525-2167(03)00011-8]
- 15 **Letsas KP**, Efremidis M, Kounas SP, Pappas LK, Gavrielatos G, Alexanian IP, Dimopoulos NP, Filippatos GS, Sideris A, Kardaras F. Clinical characteristics of patients with drug-induced QT interval prolongation and torsade de pointes: identification of risk factors. *Clin Res Cardiol* 2009; **98**: 208-212 [PMID: 19031039 DOI: 10.1007/s00392-008-0741-y]
- 16 **Mayet J**, Shahi M, McGrath K, Poulter NR, Sever PS, Foale RA, Thom SA. Left ventricular hypertrophy and QT dispersion in hypertension. *Hypertension* 1996; **28**: 791-796 [PMID: 8901825 DOI: 10.1161/01.HYP.28.5.791]
- 17 **Nielsen JB**, Graff C, Rasmussen PV, Pietersen A, Lind B, Olesen MS, Struijk JJ, Haunsø S, Svendsen JH, Køber L, Gerdts TA, Holst AG. Risk prediction of cardiovascular death based on the QTc interval: evaluating age and gender differences in a large primary care population. *Eur Heart J* 2014; **35**: 1335-1344 [PMID: 24603310 DOI: 10.1093/eurheartj/ehu081]
- 18 **Zhang Y**, Post WS, Dalal D, Blasco-Colmenares E, Tomaselli GF, Guallar E. QT-interval duration and mortality rate: results from the Third National Health and Nutrition Examination Survey. *Arch Intern Med* 2011; **171**: 1727-1733 [PMID: 22025428 DOI: 10.1001/archinternmed.2011.433]
- 19 **Bachmann TN**, Skov MW, Rasmussen PV, Graff C, Pietersen A, Lind B, Struijk JJ, Olesen MS, Haunsø S, Køber L, Svendsen JH, Holst AG, Nielsen JB. Electrocardiographic Tpeak-Tend interval and risk of cardiovascular morbidity and mortality: Results from the Copenhagen ECG study. *Heart Rhythm* 2016; **13**: 915-924 [PMID: 26707793 DOI: 10.1016/j.hrthm.2015.12.027]
- 20 **Porthan K**, Viitasalo M, Toivonen L, Havulinna AS, Jula A, Tikkanen JT, Väänänen H, Nieminen MS, Huikuri HV, Newton-Cheh C, Salomaa V, Oikarinen L. Predictive value of electrocardiographic T-wave morphology parameters and T-wave peak to T-wave end interval for sudden cardiac death in the general population. *Circ Arrhythm Electrophysiol* 2013; **6**: 690-696 [PMID: 23881778 DOI: 10.1161/CIRCEP.113.000356]
- 21 **Rosenthal TM**, Stahls PF, Abi Samra FM, Bernard ML, Khatib S, Polin GM, Xue JQ, Morin DP. T-peak to T-end interval for prediction of ventricular tachyarrhythmia and mortality in a primary prevention population with systolic cardiomyopathy. *Heart Rhythm* 2015; **12**: 1789-1797 [PMID: 25998895 DOI: 10.1016/j.hrthm.2015.04.035]
- 22 **Sauer AJ**, Selvaraj S, Aguilar FG, Martinez EE, Wilcox JE, Passman R, Goldberger JJ, Freed BH, Shah SJ. Relationship between repolarization heterogeneity and abnormal myocardial mechanics. *Int J Cardiol* 2014; **172**: 289-291 [PMID: 24447740 DOI: 10.1016/j.ijcard.2013.12.232]
- 23 **Gupta P**, Patel C, Patel H, Narayanaswamy S, Malhotra B, Green JT, Yan GX. T(p-e)/QT ratio as an index of arrhythmogenesis. *J Electrocardiol* 2008; **41**: 567-574 [PMID: 18790499 DOI: 10.1016/j.jelectrocard.2008.07.016]
- 24 **Yamaguchi M**, Shimizu M, Ino H, Terai H, Uchiyama K, Oe K, Mabuchi T, Konno T, Kaneda T, Mabuchi H. T wave peak-to-end interval and QT dispersion in acquired long QT syndrome: a new index for arrhythmogenicity. *Clin Sci (Lond.)* 2003; **105**: 671-676 [PMID: 12857349 DOI: 10.1042/CS20030010]
- 25 **Aslan MM**, Atici A. Assessment of increased risk of arrhythmia in advanced age pregnancies. *Pak J Med Sci* 2018; **34**: 687-690 [PMID: 30034440 DOI: 10.12669/pjms.343.14313]
- 26 **Kuzu F**. The effect of type 2 diabetes on electrocardiographic markers of significant cardiac events. *Pak J Med Sci* 2018; **34**: 626-632 [PMID: 30034428 DOI: 10.12669/pjms.343.14562]
- 27 **Karakulak UN**, Yilmaz OH, Tutkun E, Gunduzoz M, Ercan Onay E. Comprehensive Electrocardiographic Analysis of Lead Exposed Workers: An Arrhythmic Risk Assessment Study. *Ann Noninvasive Electrocardiol* 2017; **22** [PMID: 27282320 DOI: 10.1111/ane.12376]
- 28 **Redfield MM**, Jacobsen SJ, Burnett JC, Mahoney DW, Bailey KR, Rodeheffer RJ. Burden of systolic and diastolic ventricular dysfunction in the community: appreciating the scope of the heart failure epidemic. *JAMA* 2003; **289**: 194-202 [PMID: 12517230 DOI: 10.1016/S0022-0248(01)00919-8]
- 29 **Mozos I**. The link between ventricular repolarization variables and arterial function. *J Electrocardiol* 2015; **48**: 145-149 [PMID: 25444569 DOI: 10.1016/j.jelectrocard.2014.11.008]

- 30 **Oikarinen L**, Nieminen MS, Viitasalo M, Toivonen L, Jern S, Dahlöf B, Devereux RB, Okin PM; LIFE Study Investigators. QRS duration and QT interval predict mortality in hypertensive patients with left ventricular hypertrophy: the Losartan Intervention for Endpoint Reduction in Hypertension Study. *Hypertension* 2004; **43**: 1029-1034 [PMID: [15037560](#) DOI: [10.1161/01.HYP.0000125230.46080.c6](#)]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

