

找到约 2,580 条结果 (用时 0.59 秒)

您是不是要找： Bilateral gangrene of fingers in a patient on empagliflozin- first case report

Case Report: Symmetrical peripheral gangrene of bilateral feet and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5836706/> - 翻译此页

作者：N Ruffin - 2018 - 被引用次数：2 - 相关文章

2018年2月7日 - One full year after the patient was first admitted to the hospital, and 3 ... who developed gangrene of the fingers, toes and earlobes after shock.

缺少字词： empagliflozin

Symmetrical peripheral gangrene in sepsis after treatment with inotropes

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5509204/> - 翻译此页

作者：JL Jiang - 2017 - 被引用次数：3 - 相关文章

We report a case of four limb SPG caused by septic shock with disseminated intravascular ...

Keywords: Inotropes, Sepsis, Symmetrical peripheral gangrene.

缺少字词： empagliflozin

The first case of hand infection caused by Dermabacter jinjuensis in a ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6206325/> - 翻译此页

作者：SH Cho - 2018 - 被引用次数：1

跳到Case report - A 78 - year - old female visited the outpatient clinic with bilateral multiple finger dry gangrene and a pus discharge. On physical ...

缺少字词： empagliflozin

Circulation on the Run

<https://circulation.libsyn.com/page/3/size/25> ▼ 翻译此页

The current study focused on patients with a history of stroke prior to randomization. Possible breaking the additional exposure in already recruited the Roth case. In this study from first author Dr. Frankenreiter, corresponding author Dr. How we can make centers better without finger pointing on specific centers, and ...

Diabetes Mellitus & Hypoglycemia | Current Medical Diagnosis ...

<https://accessmedicine.mhmedical.com/content.aspx?bookID=2449...> - 翻译此页

The diabetes mellitus usually presents in the first decade together with the optic atrophy. ... beta cell function; in a case control study and a large population cohort study Impaired circulation to the fingers (for example, in patients with Raynaud surgery, inoperable coronary artery disease, and

Name of Journal: *World Journal of Diabetes*

Manuscript NO: 45456

Manuscript Type: CASE REPORT

Bilateral gangrene of fingers in a patient on empagliflozin: First case report

Ramachandra Pai RP *et al.* Empagliflozin causing finger gangrene

Rajasree Pai Ramachandra Pai, Raghesh Varot Kangath

Abstract

BACKGROUND

Sodium glucose cotransporter 2 (SGLT2) inhibitors use has been associated with toe amputations and non-healing ulcers and gangrene mostly of lower extremities. There

Match Overview

1 match

1	Crossref 19 words Gian Paolo Fadini, Angelo Avogaro. "SGLT2 inhibitors and ... amputations in the US FDA Adverse Event Reporting System	2%
2	Internet 15 words crawled on 18-Sep-2017 www.nejm.org	1%
3	Crossref 15 words S. Kumar, A. J. Costello, P. G. Colman. "Fournier's gangrene ... e in a man on empagliflozin for treatment of Type 2 diabetes	1%
4	Internet 13 words lrd.yahooapis.com	1%
5	Internet 11 words crawled on 27-Jan-2019 www.rxfiles.ca	1%
6	Internet 10 words crawled on 22-Mar-2016 www.omicsgroup.org	1%



All

Images

Videos

翻译成中文

开启取词

23,200 Results

Any time ▾

(PDF) Bilateral gas gangrene of the hand - A unique case

https://www.researchgate.net/publication/8997614_Bilateral_gas...

Gas **gangrene** is a rare and rapidly progressive process with the propensity for devastating consequences. We **report** the **first case** of **bilateral gas gangrene of the hand** following **bilateral ...**

Successful resolution of symmetrical peripheral gangrene ...

<https://jmedicalcasereports.biomedcentral.com/articles/10.1186/s...> ▾

Sep 17, 2015 · Conclusions. To the best of our knowledge, this is the **first case report** of symmetrical peripheral **gangrene** occurring after acute **pancreatitis** with successful resolution. We highlight the importance of prompt and aggressive fluid **resuscitation** and consideration of multiple **treatment** options to prevent a **hypovolemic** state caused by acute pancreatitis.

Cited by: 1

Author: Chen-Yi Liao, Shih-Chung Huang, Cheng-...

Publish Year: 2015

Bilateral optic neuropathy, acral gangrene and visceral ...

www.thefreelibrary.com > ... > October 31, 2016 ▾

1,3,4 Here, we **report** a **case** with calciphylaxis very rarely presenting with **bilateral optic neuropathy**, acral **gangrene** and visceral ischaemia. **Case Report Bilateral papilloedema** was found upon evaluating visual impairment of sudden onset in a 43-year old caucasian female **patient** with chronic renal failure (CRF) on haemodialysis in February 2014.

Peripheral gangrene in patients with severe falciparum ...

www.ncbi.nlm.nih.gov > ... > Korean J Parasitol > v.44(2); 2006 Jun

DISCUSSION. The resolution of the **gangrene** in 2 of the 3 **patients** in this **report** suggests an initially conservative approach to the management of symmetrical peripheral **gangrene**. This allows time for the **patient's** condition to stabilize and allows the **gangrenous** lesions to become completely recovered.

Cited by: 6

Author: Vipa Thanachartwet, Srivicha Krudsood, ...

Publish Year: 2006

Bilateral optic neuropathy, acral gangrene and visceral ...

jpma.org.pk/full_article_text.php?article_id=7941 ▾

We **report** a **case** with calciphylaxis very rarely presenting with **bilateral optic neuropathy**, acral **gangrene** and visceral ischaemia. **Bilateral papilloedema** was found in a 43 year-old female with chronic renal failure.

[PDF] Symmetrical peripheral gangrene of bilateral feet and ...

<https://casereports.bmj.com/content/casereports/2018/bcr-2017...>