



All

Images

Videos

翻译成中文

关闭取词

24,900 Results

Any time ▾

## Necrosectomy and postoperative local lavage in necrotizing ...

<https://onlinelibrary.wiley.com/doi/abs/10.1002/bjs.1800750306>

Necrosectomy with postoperative **continuous local lavage** was performed in a prospective study involving 95 patients with necrotizing pancreatitis. In the same period 567 patients w

Cited by: 402

Author: H. G. Beger, M. Büchler, R. Bittner, S. Bl...

Publish Year: 1988

## Complications of pancreatic surgery - PubMed Central (PMC)

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > HPB (Oxford) > v.7(2); 2005

Certainly, being a lesser procedure, when compared to PD, **postoperative morbidity** rates following such **local pancreatic head resection** are predicted to be lower. In a prospective randomized trial by Izbicki et al. 99 comparing PPPD with the Frey's procedure, the **postoperative morbidity** rates were 53.3% for the former and 19.4% for the latter. DGE was observed to occur only in the PPPD group.

Cited by: 130

Author: Choon-Kiat. Ho, Jörg. Kleeff, Helmut. Frie...

Publish Year: 2005

## Surgical Management and Complex Treatment of Infected ...

<https://www.sciencedirect.com/science/article/pii/S1091255X05005469>

Material and Methods. **Continuous** lavage was applied for an average of 44.5 days (range, 21–95 days), with a median of 9.5 L (range, 5–20 L) of **normal saline** per 24 hours. In the first few postoperative days, the amount of **lavage** fluid was generally 15–20 L, which was ...

Cited by: 73

Author: Gyula Farkas, János Márton, Yvette Mán...

Publish Year: 2006

## Continuous irrigation around pancreatic remnant decreases ...

<https://www.researchgate.net/publication/256837838> **Continuous...**

1

**Name of Journal:** *World Journal of Clinical Cases*

**Manuscript No:** 46795

**Manuscript type:** CASE REPORT

**Impact of continuous local lavage on pancreatic juice-related postoperative complications: Three case reports**

Hori T *et al.* Postoperative pancreatic fistula

Tomohide Hori, Kohei Ogawa, Hidekazu Yamamoto, Hideki Harada, Kazuyoshi Matsumura, Michihiro Yamamoto, Masahiro Yamada, Takefumi Yazawa, Katsutoshi Kuriyama, Masaki Tani, Daiki Yasukawa, Yasuyuki Kamada, Yuki Aisu, Ryotaro Tani, Ryuhei Aoyama, Shinnosuke Nakayama, Yudai Sasaki, Koki Nishimoto, Masazumi Zaima

## Match Overview

1 Internet 56 words  
[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)

2%

[国内版](#)[国际版](#)[All](#)[Images](#)[Videos](#)[翻译成中文](#)[关闭取词](#)

23,400 Results

Any time ▾

### Article abstract | American Journal of Case Reports

<https://www.amjcaserep.com/abstract/index/idArt/915010>

Nevertheless, the intractable **pancreatic** fistula triggered a bacteriogenic infection, resulting in intraperitoneal abscess. **Continuous local lavage** via transnasal **continuous** infusion and endoscopic transgastric drainage was performed, until the fistula closed. He was healthy at 9 months after surgery.

### Get your full text copy in PDF | American Journal of Case ...

<https://www.amjcaserep.com/download/index/idArt/915010> ▾

Nevertheless, the intractable **pancreatic** fistula triggered a bacteriogenic infection, resulting in intraperitoneal abscess. **Continuous local lavage** via transnasal **continuous** infusion and endoscopic transgastric drainage was performed, until the fistula closed. He was healthy at 9 months after surgery.

**Author:** Ryotaro Tani, Tomohide Hori, Hideka... **Publish Year:** 2019

### Decision Making in Necrotizing Pancreatitis | Request PDF

[https://www.researchgate.net/.../304330897\\_Decision\\_Making\\_in\\_Necrotizing\\_Pancreatitis](https://www.researchgate.net/.../304330897_Decision_Making_in_Necrotizing_Pancreatitis)

CONCLUSIONS A SAA that had the **rare form and solitary** origin was treated. **Continuous local lavage** has a **therapeutic potential** for a **pancreatic juice-related bacteriogenic complication**.

### Clinical effects of laparotomy with perioperative ...

<https://wjes.biomedcentral.com/articles/10.1186/1749-7922-4-45> ▾

Dec 16, 2009 · Methods. From a consecutive group of 23 patients hospitalized for acute pancreatitis, we studied 6 patients all with Apache II scores  $\geq 19$ , who underwent emergency surgery for acute **complications** (5 for an abdominal compartment syndrome and 1 for septic shock) followed by **continuous** perioperative peritoneal **lavage** and postoperative CVVDH.

**Cited by:** 9 **Author:** Roberto Caronna, Michele Benedetti, An...**Publish Year:** 2009

### Continuous irrigation around pancreatic remnant decreases ...

[https://www.researchgate.net/publication/256837838\\_Continuous\\_irrigation\\_around...](https://www.researchgate.net/publication/256837838_Continuous_irrigation_around...)

Careful **necrosectomy** and a **continuous local lavage** of the **lesser sac** and the **necrotic cavities** are most effective. Conservation of **vital pancreatic tissue** is a major goal of **surgical treatment**.

[国内版](#)[国际版](#)[All](#)[Images](#)[Videos](#)[翻译成中文](#)[开启取词](#)

23,700 Results

Any time ▾

### Article abstract | American Journal of Case Reports

<https://www.amjcaserep.com/abstract/index/idArt/915010>

Nevertheless, the intractable pancreatic fistula triggered a bacteriogenic infection, resulting in intraperitoneal abscess. Continuous local lavage via transnasal continuous infusion and endoscopic transgastric drainage was performed, until the fistula closed. He was healthy at 9 months after surgery.

### Get your full text copy in PDF | American Journal of Case ...

<https://www.amjcaserep.com/download/index/idArt/915010> ▾

Nevertheless, the intractable pancreatic fistula triggered a bacteriogenic infection, resulting in intraperitoneal abscess. Continuous local lavage via transnasal continuous infusion and endoscopic transgastric drainage was performed, until the fistula closed. He was healthy at 9 months after surgery.

Author: Ryotaro Tani, Tomohide Hori, Hidekaz... Publish Year: 2019

### Clinical effects of laparotomy with perioperative ...

<https://wjcs.biomedcentral.com/articles/10.1186/1749-7922-4-45> ▾

Dec 16, 2009 · Methods. From a consecutive group of 23 patients hospitalized for acute pancreatitis, we studied 6 patients all with Apache II scores  $\geq 19$ , who underwent emergency surgery for acute complications (5 for an abdominal compartment syndrome and 1 for septic shock) followed by continuous perioperative peritoneal lavage and postoperative CVVDH.

Cited by: 9

Author: Roberto Caronna, Michele Benedetti, And...

Publish Year: 2009

### Decision Making in Necrotizing Pancreatitis | Request PDF

[https://www.researchgate.net/.../304330897\\_Decision\\_Making\\_in\\_Necrotizing\\_Pancreatitis](https://www.researchgate.net/.../304330897_Decision_Making_in_Necrotizing_Pancreatitis)

CONCLUSIONS A SAA that had the rare form and solitary origin was treated. Continuous local lavage has a therapeutic potential for a pancreatic juice-related bacteriogenic complication.

### Continuous irrigation around pancreatic remnant decreases ...

[https://www.researchgate.net/publication/256837838\\_Continuous\\_irrigation\\_around...](https://www.researchgate.net/publication/256837838_Continuous_irrigation_around...)

Careful necrosectomy and a continuous local lavage of the lesser sac and the necrotic cavities are most effective. Conservation of vital pancreatic tissue is a major goal of surgical treatment.

### Pancreatic Necrosis, Surgical Management of - Practice ...

<https://www.east.org/education/practice-management-guidelines/pancreatic-necrosis...> ▾

The oldest and most established approach includes open laparotomy or retroperitoneal flank incision with manual debridement. After necrosectomy, the abdomen may be left open or drains are placed for postoperative continuous lavage. The postoperative lavage technique was associated with less operations and a decrease in postoperative morbidity. [15]

### Early and late complications after pancreatic necrosectomy ...