



国内版

国际版

Germline mutations in Thai patients with nonmucinous epithe



All

Images

Videos

关闭取词

26,700 Results

Any time ▾

[Uptake of testing for germline BRCA mutations in patients ...](#)

<https://ijgc.bmj.com/content/early/2019/05/15/ijgc-2019-000389> ▾

May 16, 2019 · Introduction **Patients with non-mucinous epithelial tubo-ovarian cancers** should be referred for genetic testing because approximately 15% will carry an inherited **mutation** in the BRCA1 or **BRCA2 cancer susceptibility genes**.

Author: Grace Stearnes, Cassandra B Nichols... **Publish Year:** 2019

[Germline BRCA1 and BRCA2 Mutations in Ovarian Cancer ...](#)

[https://insights.ovid.com/obstetrics-gynecology/...](https://insights.ovid.com/obstetrics-gynecology/) ▾

OBJECTIVE: To estimate the frequency of **BRCA1** and **BRCA2 germline mutations** in women with **nonmucinous epithelial ovarian carcinoma** unselected for a family **history of breast or ovarian cancer**.

Cited by: 97 **Author:** Kasmintan A. Schrader, Jane Hurlburt, St...

Publish Year: 2012

[Germline BRCA1 and BRCA2 Mutations in Ovarian Cancer ...](#)

<journals.lww.com/greenjournal/Pages/articleviewer.aspx?year=2012&issue=08000&article=...>

OBJECTIVE: To estimate the frequency of **BRCA1** and **BRCA2 germline mutations** in women with **nonmucinous epithelial ovarian carcinoma** unselected for a family **history of breast or ovarian cancer**.

[High-risk epithelial ovarian cancer patients for ...](#)

<onlinelibrary.wiley.com/doi/10.1111/jog.13287/abstract>

Three **patients** (8.6%) had **MMR gene mutations** (one MLH1 and two MSH2) and one **patient** (2.9%) had other **gene mutation** (MUTYH). Of the 10 **patients with germline mutation**, 40% of **patients** had no significant family history of **cancer**. Conclusion. Up to 30% of **EOC patients** had risk factors for hereditary **ovarian cancer**.

Cited by: 1 **Author:** Seksit Chirasophon, Tarinee Manchana, ...

Publish Year: 2017



国内版

国际版

Germline mutations in Thai non-mucinous epithelial ovarian cancer patients



All

Images

Videos

翻译成中文

关闭取词

26,100 Results

Any time ▾

[High-risk epithelial ovarian cancer patients for ...](#)

onlinelibrary.wiley.com/doi/10.1111/jog.13287/abstract

The objectives of this study were to evaluate the frequency of high-risk **epithelial ovarian cancer (EOC) patients** and the frequency of **germline mutation** in these **patients**. Methods A total of 335 **patients** with histologically confirmed **non-mucinous EOC** were included.

Author: Seksit Chirasophon, Tarinee Manchan... **Publish Year:** 2017

[\(PDF\) High-risk epithelial ovarian cancer patients for ...](#)

https://www.researchgate.net/publication/313624753_High-risk...

High-risk **epithelial ovarian cancer patients** for hereditary **ovarian cancer**: Hereditary **epithelial ovarian cancer**. **Germline mutation** was identified in 28.6% of **patients** (11.4% **BRCA mutation**, 5.7% other HR genes **mutation**, 8.6% **MMR mutation**, and 2.9% **mutation** in other gene). The high cost of genetic testing is an important barrier. Selected **patients** with high-risk factors might be initially ...

[Evaluation of germline BRCA1 and BRCA2 mutations in a ...](#)

<https://www.sciencedirect.com/science/article/pii/S0090825815301736>

Germline mutations in BRCA1 and BRCA2 confer an increased risk of **breast and ovarian cancer**, with average cumulative risks of **ovarian cancer** by age 70 years of 39–40% for BRCA1 **mutation** carriers, and 11–18% for BRCA2 **mutation** carriers, compared to ...

Cited by: 1 **Author:** Hanis Nazihah Hasmad, Kah Nyin Lai, W...

Publish Year: 2016

[A universal genetic testing initiative for patients with ...](#)

[https://www.gynecologiconcology-online.net/article/S0090-8258\(17...](https://www.gynecologiconcology-online.net/article/S0090-8258(17...)

Approximately 10–20% of high-grade, **non-mucinous epithelial ovarian**, fallopian tube, and primary peritoneal cancers (HGOC) are hereditary, primarily due to **germline mutations** in the BRCA1 or BRCA2 genes [1–3]. A **mutation** in BRCA1 or BRCA2 confers a 40–66% lifetime risk of **breast cancer** and a 13–46% lifetime risk of **ovarian cancer** in women [4].

Cited by: 10 **Author:** Erica M. Bednar, Holly D. Oakley, Charlot...

Publish Year: 2017



时间不限

2019以来

2018以来

2015以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

包括专利

包含引用

创建快讯

您是不是要找: Germline mutations in Thai patients with *non mucinous* epithelial ovarian cancer

High-risk epithelial ovarian cancer patients for hereditary ovarian cancer

[PDF] researchgate.net

S Chirasophon, T Manchana... - Journal of Obstetrics ..., 2017 - Wiley Online Library

... Of 335 Thai EOC patients, 94 patients (28.1%) were identified as high-risk patients ... Three mutations (8.6%) were found in BRCA1 gene, one (2.8%) in BRCA2 gene, three (8.6%) in ... one RAD1C, and one MUTYH) (Table 2). Of the 10 patients with germline mutation, four patients ...

☆ 被引用次数: 3 相关文章 所有 5 个版本

Role of Biomarkers in Screening and Diagnosis of Ovarian Cancer

[PDF] tci-thaijo.org

Y Poomtavorn - Thai Journal of Obstetrics and Gynaecology, 2016 - tci-thaijo.org

... Oncology Unit, Department of Obstetrics and Gynecology, Faculty of Medicine, Thammasat University Hospital, Pathumthani 12120, Thailand, Phone: +66 ... 228 Thai J Obstet Gynaecol ... BRCA genetic mutations and other germline mutations are associated with type 2 carcinomas ...

☆ 相关文章

Surface epithelial tumors of the ovary

JD Seidman, KR Cho, BM Ronnett... - Blaustein's pathology of the ..., 2011 - Springer

... These findings highlight the importance of p53 mutation in the early development of ... The genes mutated in cancer are not selected randomly, but frequently encode proteins ... ovarian epithelial tumors for molecular genetic alterations such as point mutations, gene amplifications ...

☆ 被引用次数: 335 相关文章 所有 4 个版本

The Challenge of Evaluating Adnexal Masses in Patients With Breast Cancer

T Reinert, A Nogueira-Rodrigues, FP Kestelman... - ... breast cancer, 2018 - Elsevier

... BC and OC are summarized in Table 1. The identification of a germline mutation in one ... OC.42, 43 The use of PARP inhibitors is well-established for BRCA-mutated relapsed patients ... 44 In addition to these classic BC and OC hereditary syndromes, germline mutations in several ...

☆ 被引用次数: 1 相关文章 所有 5 个版本

Name of Journal: *World Journal of Clinical Oncology*

Manuscript NO: 46939

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Germline mutations in Thai patients with nonmucinous epithelial ovarian cancer

Manchana T *et al.* Germline mutations in epithelial ovarian cancer

Tarinee Manchana, Prasit Phowthongkum, Chinachote Teerapakpinyo

Abstract

BACKGROUND

Match Overview

1	Internet 47 words www.ncbi.nlm.nih.gov	1%
2	Internet 14 words crawled on 20-Jul-2019 scholars.uab.edu	<1%
3	Crossref 12 words Tarinee Manchana, Natacha Phoolcharoen, Patou Tantbiro ... "BRCA mutation in high grade epithelial ovarian cancers", G	<1%