

World Journal of *Gastroenterology*

World J Gastroenterol 2019 September 14; 25(34): 5017-5219



**OPINION REVIEW**

- 5017** Reducing anastomotic leak in colorectal surgery: The old dogmas and the new challenges
Meyer J, Naiken S, Christou N, Liot E, Toso C, Buchs NC, Ris F

REVIEW

- 5026** Diagnostic and prognostic potential of tissue and circulating long non-coding RNAs in colorectal tumors
Galamb O, Barták BK, Kalmár A, Nagy ZB, Szigeti KA, Tulassay Z, Igaz P, Molnár B
- 5049** Autoantibodies: Potential clinical applications in early detection of esophageal squamous cell carcinoma and esophagogastric junction adenocarcinoma
Xu YW, Peng YH, Xu LY, Xie JJ, Li EM

MINIREVIEWS

- 5069** Hepatic senescence, the good and the bad
Huda N, Liu G, Hong H, Yan S, Khambu B, Yin XM
- 5082** Role of endoscopic ultrasound in the screening and follow-up of high-risk individuals for familial pancreatic cancer
Lorenzo D, Rebours V, Maire F, Palazzo M, Gonzalez JM, Vullierme MP, Aubert A, Hammel P, Lévy P, Mestier LD
- 5097** Optimizing proton pump inhibitors in *Helicobacter pylori* treatment: Old and new tricks to improve effectiveness
Ierardi E, Losurdo G, Fortezza RFL, Principi M, Barone M, Leo AD

ORIGINAL ARTICLE**Basic Study**

- 5105** Regulatory effect of a Chinese herbal medicine formula on non-alcoholic fatty liver disease
Yang JM, Sun Y, Wang M, Zhang XL, Zhang SJ, Gao YS, Chen L, Wu MY, Zhou L, Zhou YM, Wang Y, Zheng FJ, Li YH
- 5120** Allyl isothiocyanate ameliorates lipid accumulation and inflammation in nonalcoholic fatty liver disease via the Sirt1/AMPK and NF- κ B signaling pathways
Li CX, Gao JG, Wan XY, Chen Y, Xu CF, Feng ZM, Zeng H, Lin YM, Ma H, Xu P, Yu CH, Li YM
- 5134** Effect of Tong Xie Yao Fang on endogenous metabolites in urine of irritable bowel syndrome model rats
Zhao XY, Wang JW, Yin Y, Li K, Zhang M, Yan FP

Case Control Study

- 5152** Association of XPG rs2094258 polymorphism with gastric cancer prognosis
Wang XQ, Terry PD, Li Y, Zhang Y, Kou WJ, Wang MX

Retrospective Study

- 5162** Timing, distribution, and microbiology of infectious complications after necrotizing pancreatitis
Lu JD, Cao F, Ding YX, Wu YD, Guo YL, Li F
- 5174** Risk factors for Mallory-Weiss Tear during endoscopic submucosal dissection of superficial esophageal neoplasms
Chen W, Zhu XN, Wang J, Zhu LL, Gan T, Yang JL

Clinical Trials Study

- 5185** Overlay of a sponge soaked with ropivacaine and multisite infiltration analgesia result in faster recovery after laparoscopic hepatectomy
Zhang H, Du G, Liu YF, Yang JH, A-Niu MG, Zhai XY, Jin B

Observational Study

- 5197** Analysis of 72 patients with colorectal high-grade neuroendocrine neoplasms from three Chinese hospitals
Wang ZJ, An K, Li R, Shen W, Bao MD, Tao JH, Chen JN, Mei SW, Shen HY, Ma YB, Zhao FQ, Wei FZ, Liu Q

META-ANALYSIS

- 5210** Bilateral *vs* unilateral placement of metal stents for inoperable high-grade hilar biliary strictures: A systemic review and meta-analysis
Ashat M, Arora S, Klair JS, Childs CA, Murali AR, Johlin FC

ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Matthias Ocker, MD, Professor, Department of Translational Medicine Oncology, Bayer AG, Berlin 13353, Germany

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. The *WJG* Editorial Board consists of 701 experts in gastroenterology and hepatology from 58 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, etc. The *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

The *WJG* is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, and Scopus. The 2019 edition of Journal Citation Report® cites the 2018 impact factor for *WJG* as 3.411 (5-year impact factor: 3.579), ranking *WJG* as 35th among 84 journals in gastroenterology and hepatology (quartile in category Q2). CiteScore (2018): 3.43.

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: Yan-Liang Zhang

Proofing Production Department Director: Xiang Li

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Subrata Ghosh, Andrzej S Tarnawski

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE

Ze-Mao Gong, Director

PUBLICATION DATE

September 14, 2019

COPYRIGHT

© 2019 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Reducing anastomotic leak in colorectal surgery: The old dogmas and the new challenges

Jeremy Meyer, Surennaidoo Naiken, Niki Christou, Emilie Liot, Christian Toso, Nicolas Christian Buchs, Frédéric Ris

ORCID number: Jeremy Meyer (0000-0003-3381-9146); Surennaidoo Naiken (0000-0002-7195-4974); Niki Christou (0000-0003-2125-0503); Emilie Liot (0000-0002-2856-5260); Christian Toso (0000-0003-1652-4522); Nicolas Christian Buchs (0000-0001-9255-3929); Frédéric Ris (0000-0001-7421-6101).

Author contributions: Meyer J conceived the review and wrote the draft of the manuscript; Meyer J, Naiken S, Christou N, Liot E, Toso C, Buchs NC and Ris F reviewed and accepted the manuscript.

Conflict-of-interest statement: The authors have no conflict of interest to declare.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Received: March 19, 2019

Peer-review started: March 19, 2019

Jeremy Meyer, Surennaidoo Naiken, Niki Christou, Emilie Liot, Christian Toso, Nicolas Christian Buchs, Frédéric Ris, Division of Digestive Surgery, University Hospitals of Geneva, Genève 1211, Switzerland

Corresponding author: Jeremy Meyer, MD, PhD, Surgeon, Division of Digestive Surgery, University Hospitals of Geneva, Rue Gabrielle-Perret-Gentil 4, Genève 1211, Switzerland. jeremy.meyer@hcuge.ch

Telephone: +41-79-5533182

Fax: +41-22-3727707

Abstract

Anastomotic leak (AL) constitutes a significant issue in colorectal surgery, and its incidence has remained stable over the last years. The use of intra-abdominal drain or the use of mechanical bowel preparation alone have been proven to be useless in preventing AL and should be abandoned. The role of oral antibiotics preparation regimens should be clarified and compared to other routes of administration, such as the intravenous route or enema. In parallel, preoperative antibiotherapy should aim at targeting collagenase-inducing pathogens, as identified by the microbiome analysis. AL can be further reduced by fluorescence angiography, which leads to significant intraoperative changes in surgical strategies. Implementation of fluorescence angiography should be encouraged. Progress made in AL comprehension and prevention might probably allow reducing the rate of diverting stoma and conduct to a revision of its indications.

Key words: Anastomotic leakage; Rectal surgery; Colic surgery; Prevention; Surgical site infection; Anastomosis; Complication

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: The present manuscript reviews the current evidence regarding the prevention of anastomotic leak in colorectal surgery. Oral antibiotics and fluorescence angiography might help reduce the incidence of anastomotic leak. Study of the microbiome might offer interesting paths for research. Progress made in anastomotic leak comprehension and prevention might allow reducing the rate of diverting stoma and conduct to a revision of its indications.

First decision: August 3, 2019**Revised:** August 9, 2019**Accepted:** August 19, 2019**Article in press:** August 19, 2019**Published online:** September 14, 2019**P-Reviewer:** Zhao WT**S-Editor:** Yan JP**L-Editor:** A**E-Editor:** Zhang YL

Citation: Meyer J, Naiken S, Christou N, Liot E, Toso C, Buchs NC, Ris F. Reducing anastomotic leak in colorectal surgery: The old dogmas and the new challenges. *World J Gastroenterol* 2019; 25(34): 5017-5025

URL: <https://www.wjgnet.com/1007-9327/full/v25/i34/5017.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v25.i34.5017>

INTRODUCTION

Anastomotic leak (AL) refers to the communication between hollow viscera lumen and the peritoneal cavity at the anastomotic level^[1]. Despite a lack of widely accepted consensus regarding the definition of AL^[2,3], AL was documented to occur in 8.1% of patients after right hemicolectomy according to the 2015 ESCP snapshot audit^[4], 6.4% after colonic cancer surgery according to a nationwide Danish study including 9'333 patients^[5] and 11% after rectal cancer surgery as reported by a systematic review and meta-analysis pooling 84 studies^[6]. AL is graded according to the therapeutic management it requires: Grade A (no management), grade B (non operative management), grade C (operative management)^[3].

In addition to the septic complications and prolonged hospitalization induced, AL leads to delayed adjuvant chemotherapy or no chemotherapy at all^[7]. Further, a recent systematic review and meta-analysis including 78'434 colorectal cancer patients showed that AL was associated with increased local recurrence [risk ratio (RR) 1.90] and reduced overall survival (RR 1.36)^[8]. Another systematic review and meta-analysis including 11'353 rectal cancer patients demonstrated that AL led to increased local recurrence [hazard ratio (HR) 1.71] and decreased survival (HR 1.67) and cancer-specific survival (HR 1.03) after anterior resection^[9].

Despite the human and financial costs generated by AL^[10], and the efforts put in reducing its occurrence, the incidence of AL has not evolved among the last years. Further, pre-operative prediction of AL and identification of at-risk patients are not accurate enough^[11], and AL is often diagnosed too late in the postoperative period^[12]. In an effort to optimize the therapeutic care of patients with colorectal anastomosis with the hope to reduce the occurrence of AL, we will review old dogmas regarding prevention of AL and confront them to the most recent evidence, and will define the new challenges in the field.

PREOPERATIVE MEASURES TO PREVENT AL

Patient-related factors

A recent systematic review identified several adjustable and non-adjustable risk factors for AL, including male gender, smoking, obesity, alcohol, steroid and non-steroidal anti-inflammatory drugs, operative time, transfusion, contamination of the operative field and emergency surgery^[13]. Further, albumin < 3.5 g/dL, anemia, hypotension and use of inotropes were reported to increase the risk of AL^[14]. Preoperative radiotherapy was also documented to constitute a risk factor for AL^[15], especially if surgery occurred within an interval of 11-17 d after radiotherapy^[16]. Therefore, adjustable risk factors should be corrected before proceeding to a digestive anastomosis, in order to reduce the risk of AL. This can be partly done through enhanced recovery programs^[17], whose implementation in colorectal surgery units led to decreased postoperative morbidity and length of stay^[18]. However, the effect of enhanced recovery protocols on the rate of AL remains to be demonstrated.

Preoperative oral antibiotics and mechanical bowel preparation

Preoperative mechanical bowel preparation alone has lost in interest after multiple publications demonstrating its absence of benefit in reducing AL in elective colorectal surgery. Contant *et al*^[19] randomized 1'431 patients to receive or not mechanical bowel preparation before elective colorectal surgery, and showed that patients who received mechanical bowel preparation did not have a lower rate of AL. Further, a systematic review and meta-analysis of randomized clinical trial (RCT) including 4'859 patients confirmed this finding^[20].

Recently, authors, such as Scarborough^[21], postulated that mechanical bowel preparation allowed to improve the delivery of oral antibiotic preparation to the bowel mucosa and could therefore not be assessed independently. Using the American College of Surgeons National Surgical Quality Improvement Program

database and including 4'999 patients, they showed that combined preoperative oral antibiotics and mechanical bowel preparation lowered the rate of AL from 5.7% to 2.8% in colorectal surgery when compared to patients not receiving any kind of preparation. However, neither oral antibiotics alone or MBP alone allowed to lower the rate of AL^[21]. Further publications reached the same conclusions but used the same database^[22,23]. However, latest studies using the same database only showed an effect of oral antibiotics alone and demonstrated that combination with MBP offered no additional advantage^[24,25].

Therefore, a large RCT is needed to determine whether oral antibiotics alone and intravenous antibiotics are sufficient in reducing AL after colorectal surgery or whether association with mechanical bowel preparation is needed^[26]. Furthermore, the type of MBP is very different among surgeon practice^[27] and probably needs standardization before conclusion could be drawn from MBP studies. The use of rectal enema associated or not with antibiotics should be assessed; as evidence is growing that the local microbiome at the anastomotic site might be implicated in AL, as discussed below.

Microbiology

Recent evidence supports the hypothesis that AL might result from a local infective complication, resulting in impaired healing at the anastomotic level due to a local increase in collagenase activity. For instance, Shogan *et al*^[28] showed in a rat model that *Enterococcus faecalis* led to the degradation of collagen IV at the anastomotic level through activation of MMP9. Further, topical antibiotherapy administered by enema targeting *Enterococcus faecalis* allowed to reduce AL to 0%, whereas intramuscular cefotixim – commonly used for elective surgery prophylaxis – did not reduce collagenase activity nor AL^[28]. Moreover, a recent study using rat model of colo-colic anastomosis demonstrated that the selective MMP-8, MMP-9, and MMP-12 inhibitor AZD3342 allowed to maintain the anastomosis baseline breaking strength and to reduce AL^[29]. Butyrate, a short-chain fatty acid, was shown to reduce AL^[30-32], probably through its inhibitory effect on *Pseudomonas aeruginosa*^[33]. In a case-control study including 8 patients with AL and 8 patients without AL after stapled colorectal anastomosis, van Praagh *et al*^[34] showed that patients with AL had lower microbial diversity and higher abundance of *Lachnospiraceae*. They postulated that the higher rate of AL might be explained by the presence of mucin-degrading *Ruminococci* within that family^[34]. Stumpf *et al*^[35] found lower collagen type I/III ratio and higher expression of MMP-1, -2 and -9 in biopsies of patients with impaired anastomotic healing when compared to controls. These results suggest that unfavorable microbiome comprising collagenase-inducing pathogens might impair anastomotic healing and result in AL.

Further, anastomosis creation was shown to result in a 200- and 500-fold increase in the relative abundance of *Enterococcus* and *Escherichia/Shigella*, respectively, in a rodent model^[36]. In a prospective multicentric cohort of patients undergoing colorectal surgery, including our center, Dubinsky-Pertsov *et al*^[37] showed that carriers of beta lactamase-producing *Enterobacteriaceae* receiving cephalosporin-based antibioprophylaxis were at risk of surgical site infection [odds ratio (OR) 2.36]. These findings suggest that changes in the local microbiome caused by surgery or inappropriate prophylactic antibiotherapy might worsen the situation of patients with already unfavorable microbiome profiles. Also, radiotherapy was documented to change the composition of the microbiome^[38,39], which constitutes a finding of importance for rectal cancer patients receiving neoadjuvant treatment but with no clear demonstration in increased AL so far.

Microbiome is a new and very promising field of research, especially when studying the aetiologies of AL in colorectal surgery. Identifying at risk patients with unfavorable microbiome, comprising pathogens with high collagenase activity, and treating them with appropriate antibiotic regimen (per os, intravenous or by enema) and/or faecal transplantation if required could help reducing AL rate. Further, studying the microbiome might help explaining the protective effect of preoperative oral antibiotics on the AL rate.

OPERATIVE MEASURES TO PREVENT AL

Surgical approach

The United States Nationwide Inpatient Sample database (including 244'129 elective colectomies) was analyzed to compare outcomes between robot-assisted colectomy (1'584 colectomies), laparoscopic colectomy (116'261 colectomies) and open colectomy (126'284 colectomies). AL was not reported, but the authors described laparoscopic

colectomy to lower the risk of complications (19.8% *vs* 33.2%) and stoma (3.5% *vs* 13.0%) when compared to the open approach. No difference could be found between laparoscopy and the robotic technique regarding these outcomes^[40]. On the opposite, analysis of the The Danish Colorectal Cancer Group database described laparoscopic colectomy as a risk factor for AL (OR 1.34) in 9'333 patients^[5]. Regarding right colectomies, a systematic review and meta-analysis including 7'780 patients found no difference in terms of AL between the laparoscopic and robotic approaches^[41]. Regarding elective and emergency sigmoidectomy for diverticulitis, a Cochrane systematic review and meta-analysis of RCT did not describe any difference in terms of reoperation due to AL between patients with laparoscopic colectomy and those with open colectomy (349 pooled patients for that outcome)^[42]. Further, the intermediate analysis of the ROLARR trial described no difference in AL rate between the two approaches for rectal cancer^[43]. Regarding the latest, the transanal total mesorectum excision (taTME) technique, bypassing the anatomic limitations of the narrow pelvis, might allow to reduce AL, but remains to be evaluated for that outcome.

Anastomosis technique

Handsewn and stapled anastomoses are still widely performed according to surgeons preferences, reflecting the lack of consensus regarding the anastomotic method. A Cochrane systematic review and meta-analysis including 1'233 patients from 9 RCT found no difference in terms of AL, clinical AL and radiological AL between patients with stapled or handsewn colorectal anastomoses^[44]. However, the authors did not perform subgroup analysis according to the underlying disease or to the presence or not of associated procedures (drainage, diverting stoma). Further, all included studies were anterior to 1995. In emergency procedures, another systematic review and meta-analysis did not identify any statistical differences between stapled and handsewn anastomoses (1'120 patients)^[45].

Regarding right colectomy or ileo-caecal resection, the 2015 ESCP audit described an AL rate of 8.1% among 3'208 patients. After adjustment for confounding factors, the use of a stapler was significantly associated with AL (OR 1.43)^[4]. Further, stapled anastomoses were more frequently used in low risk patients, resulting in a likely underestimation of the risk of AL after right colectomy or ileo-caecal resection.

We should note that lower anastomoses are more at risk of AL, as known since decades^[15,46]. A snapshot audit specifically concerning left colon, sigmoid and rectal resections is currently undergoing^[47] and therefore conclusion cannot be reached regarding left colon and rectal surgery. No evidence is supporting either of the construction methods used for colorectal anastomosis (side to side, end to side, side to end, end to end). The evidence seems to be more straightforward regarding the number of cartridges used for rectal division. A retrospective study from Austria demonstrated in 382 patients who benefited from rectal division using a linear stapler and colorectal anastomosis using a circular stapler or compression device, that the use of 3 or more cartridges increased the incidence of AL (19.4% AL in this subgroup)^[48]. Further, the number of intersections of staple lines also correlated to the rate of AL in colorectal anastomosis using a double stapling technique^[49]. A single stapling technique for colorectal anastomosis (in TaTME for example), in opposition with the conventional double stapling technique, was demonstrated to be safe in low anterior resection but lacks evidences in term of reduction of AL^[50].

Compression anastomosis consists of a stapler equipped with disposable rings used for colorectal anastomosis: The rings are applied on each side at the anastomotic level and are evacuated into the stools once tissue necrosis and healing have occurred. A study performed in pigs with colorectal anastomoses showed that compression anastomosis was associated with less inflammation and scarring when compared with the stapling technique^[51]. A retrospective multicentric study including 1'180 patients described an AL rate of 3.22% using the ColonRing device^[52]. Further, a prospective postmarketing evaluation of the ColonRing described an AL rate of 5.3% among 266 patients, but a septic anastomotic complication rate of 8.3%, which could reflect the true AL rate^[53]. A recent systematic review and meta-analysis including 10 RCT (1'969 patients) found no difference in terms of AL between patients with handsewn or stapled anastomosis (977 patients) and those with compression anastomoses (992 patients)^[54]. Compression anastomosis has however not gained in popularity.

Intraoperative assessment of the anastomosis

As previously reported^[55], many methods have been develop to perioperatively assess the integrity of colorectal anastomoses. Briefly, the air leak test, which consists in insufflating the bowel at the anastomotic level to detect any AL, was demonstrated to help identifying AL perioperatively and led to their repair, resulting in lower rate of postoperative AL^[56,57]. Intraoperative endoscopy, in addition, to evaluate the

anastomosis integrity, allows identifying bleeding at the anastomotic level or disruption of the anastomosis^[58]. However, it requires endoscopy skills, extra material, is time-consuming and requires further scientific validation in terms of AL prevention^[59].

New methods rely on the assessment of the blood supply to the anastomosis. Adequate perfusion of the healing tissue is key to prevent AL, and a reduction in the blood flow at the rectal stump was shown to correlate with AL^[60]. Historical methods include relying on the color of the bowel, as proposed by Goligher^[61], or observing the pulsatile flow at the cut section, as stated by Novell and Lewis^[62]. Objective and reliable methods assessing anastomosis vascularization have been developed since, as reported in our recent review^[55], mentioning notably Doppler ultrasound^[63] and light spectroscopy^[64]. More recently, fluorescence perfusion angiography has showed a widespread clinical use. Briefly, a fluorophore is injected intravenously, excited by a specific wavelength to emit in another specific wavelength (usually infrared) just after vessel division and/or completion of the anastomosis, allowing the surgeon to identify any defect in vascularization at the anastomotic level. Jafari *et al.*^[65] reported that fluorescence perfusion angiography allowed to reduce AL from 18% to 6% after robotic-assisted anterior resection. Using a prospective cohort of 504 patients undergoing elective colorectal surgery with anastomosis, our team demonstrated that fluorescence perfusion angiography allowed for a change in the strategy of bowel division due to insufficient perfusion in 5.8% of patients, with no subsequent AL^[66]. Results of the PILLAR II study documented a change in the surgical plan in 8% of 139 included patients undergoing anterior resection with no subsequent AL in those patients^[67]. A recent systematic review and meta-analysis pooling 1'302 patients confirmed these results by reporting that fluorescence perfusion angiography reduced the rate of AL in patients operated for colorectal cancer^[68].

Therefore, old methods allowing assessing the integrity of the anastomosis and the absence of AL should be combined to new technologies, such as fluorescence perfusion angiography, which aim at determining the vascularization of the anastomosis, a prerequisite to an efficient healing process without subsequent AL. New studies should aim at determining whether stimulation of the neoangiogenesis process, for example by the local administration of recombinant VEGF^[69], could help in further reducing the occurrence of AL.

Diverting stoma

The creation of a lateral ileostomy or colostomy in patients at risk of AL aims at diverting the bowel content away from the anastomosis in order to decrease the rate of AL and the related morbidity. However, diverting stoma expose the patients to the risk of dehydration or to stoma closure-related complications. Further, they lead to an additional scare or won't be closed in a significant proportion of patients^[70].

A Cochrane systematic review and meta-analysis including RCT assessing the use of prophylactic stoma versus no stoma in patients with low anterior resection for rectal cancer until November 2009 described the use of covering stoma to lower the incidence of AL (RR 0.33)^[71]. Thereafter, a review of 525 patients with colo-anal anastomosis from the NSQIP database identified the absence of stoma as a risk factor for developing postoperative sepsis (OR 6.29), although the rate of AL was not reported. Also, allocation to the stoma group was not randomized and the effect was not observed in patients with low pelvic anastomosis (1'266 patients)^[72]. A systematic review and meta-analysis including all studies published between 2014 and 2017 regarding the role of a protective stoma in patients undergoing low anterior resection, identified the presence of a stoma as a protective factor against AL (RR 0.38, 5'612 patients, 11 studies)^[73]. A later systematic review and meta-analysis including only RCT (4 RCT, 358 patients) confirmed that diverting stoma lowers the risk of AL (OR 0.32)^[74]. The Cochrane collaboration produced a systematic review and meta-analysis pooling 648 patients from 6 RCT and identified diverting stoma as a protective factor against clinical AL (RR 0.33) after low anterior resection^[71].

Evidence regarding "ghost ileostomy" – a bowel loop brought through the abdominal wall but left unopened, leaving the possibility to be transformed in an ileostomy if needed – is low and remains to be clarified^[3]. Therefore, we can conclude that diverting stoma allows reducing the occurrence of AL in at-risk patients (those with low anastomosis). Ghost ileostomy could constitute a solution to avoid the occurrence of stoma-related complications, but it should be kept in mind that ghost ileostomy won't allow to avoid AL but rather to decrease its morbidity.

Prophylactic intra-abdominal drainage

Prophylactic intra-abdominal drainage during elective colorectal surgery was thought to help monitoring the occurrence of AL and to reduce its morbidity by avoiding a generalized peritonitis. The GRECCAR 5 trial compared 236 randomized rectal cancer

patients allocated to the intra-abdominal drain group to 233 patients allocated to the group without drainage. Intra-abdominal drainage did not allow to reduce the rate of pelvic sepsis, the postoperative morbidity, the reoperation rate, the length of hospital stay and the rate of stoma closure^[75]. Later, a systematic review and meta-analysis pooling 760 patients from 4 RCT demonstrated that intra-abdominal drainage did not reduce AL, pelvic complications, reintervention and mortality. Contrariwise, the incidence of postoperative bowel obstruction was significantly higher in the drained group (OR 1.61)^[76]. A Cochrane systematic review obtained the same conclusion that prophylactic intra-abdominal drainage did not reduce the rate of AL^[77]. Therefore, prophylactic intra-abdominal drainage should be discouraged in elective colorectal surgery.

Prophylactic transanal tube decompression

Prophylactic transanal tube decompression was thought to lower the risk of AL whilst presenting less risks of complication than diverting stoma. A systematic review and meta-analysis pooling 1772 patients undergoing anterior resection described transanal tube decompression to lower the risk of AL (RR 0.44)^[78]. However, patients receiving diverting stoma were excluded, leading to a potential underestimation of the AL rate. Another systematic review and meta-analysis followed, including patients with diverting stoma, and obtained the same conclusion (a reduction of the risk of AL (RR 0.42) in patients with transanal tube decompression)^[79]. Therefore, prophylactic transanal tube decompression could constitute an efficient method to prevent AL in high risk patients without exposing them to the complications of diverting stoma. A well-conducted large scale RCT comparing the 2 techniques remains, however, to be conducted.

CONCLUSION

AL still constitutes a significant issue in colorectal surgery, and its incidence has remained stable over the last years. The use of intra-abdominal drain or the use of mechanical bowel preparation alone have been proven to be useless in preventing AL and should be abandoned. The role of oral antibiotics preparation regimens should be clarified and compared to other routes of administration, such as the intravenous route or enema. In parallel, the composition of the microbiome of patients with AL should be precisely determined, in order to identify patients at risk of AL and offer targeted preoperative antibiotics. AL can be further reduced by fluorescence angiography, which leads to significant intraoperative changes in surgical strategies. Implementation of fluorescence angiography should be encouraged. Progress made in AL comprehension and prevention might probably allow reducing the rate of diverting stoma and conduct to a revision of its indications.

REFERENCES

- 1 **Rahbari NN**, Weitz J, Hohenberger W, Heald RJ, Moran B, Ulrich A, Holm T, Wong WD, Tiet E, Moriya Y, Laurberg S, den Dulk M, van de Velde C, Büchler MW. Definition and grading of anastomotic leakage following anterior resection of the rectum: A proposal by the International Study Group of Rectal Cancer. *Surgery* 2010; **147**: 339-351 [PMID: 20004450 DOI: 10.1016/j.surg.2009.10.012]
- 2 **Bruce J**, Krukowski ZH, Al-Khairy G, Russell EM, Park KG. Systematic review of the definition and measurement of anastomotic leak after gastrointestinal surgery. *Br J Surg* 2001; **88**: 1157-1168 [PMID: 11531861 DOI: 10.1046/j.0007-1323.2001.01829.x]
- 3 **Vallance A**, Wexner S, Berho M, Cahill R, Coleman M, Haboubi N, Heald RJ, Kennedy RH, Moran B, Mortensen N, Motson RW, Novell R, O'Connell PR, Ris F, Rockall T, Senapati A, Windsor A, Jayne DG. A collaborative review of the current concepts and challenges of anastomotic leaks in colorectal surgery. *Colorectal Dis* 2017; **19**: O1-O12 [PMID: 27671222 DOI: 10.1111/codi.13534]
- 4 **2015 European Society of Coloproctology collaborating group**. The relationship between method of anastomosis and anastomotic failure after right hemicolectomy and ileo-caecal resection: An international snapshot audit. *Colorectal Dis* 2017 [PMID: 28263043 DOI: 10.1111/codi.13646]
- 5 **Krørup PM**, Jørgensen LN, Andreassen AH, Harling H; Danish Colorectal Cancer Group. A nationwide study on anastomotic leakage after colonic cancer surgery. *Colorectal Dis* 2012; **14**: e661-e667 [PMID: 22564292 DOI: 10.1111/j.1463-1318.2012.03079.x]
- 6 **Paun BC**, Cassie S, MacLean AR, Dixon E, Buie WD. Postoperative complications following surgery for rectal cancer. *Ann Surg* 2010; **251**: 807-818 [PMID: 20395841 DOI: 10.1097/SLA.0b013e3181dae4ed]
- 7 **Smith JD**, Paty PB, Guillem JG, Temple LK, Weiser MR, Nash GM. Anastomotic leak is not associated with oncologic outcome in patients undergoing low anterior resection for rectal cancer. *Ann Surg* 2012; **256**: 1034-1038 [PMID: 22584695 DOI: 10.1097/SLA.0b013e318257d2c1]
- 8 **Ha GW**, Kim JH, Lee MR. Oncologic Impact of Anastomotic Leakage Following Colorectal Cancer Surgery: A Systematic Review and Meta-Analysis. *Ann Surg Oncol* 2017; **24**: 3289-3299 [PMID: 28608118 DOI: 10.1245/s10434-017-5881-8]
- 9 **Wang S**, Liu J, Wang S, Zhao H, Ge S, Wang W. Adverse Effects of Anastomotic Leakage on Local

- Recurrence and Survival After Curative Anterior Resection for Rectal Cancer: A Systematic Review and Meta-analysis. *World J Surg* 2017; **41**: 277-284 [PMID: 27743072 DOI: 10.1007/s00268-016-3761-1]
- 10 **Ashraf SQ**, Burns EM, Jani A, Altman S, Young JD, Cunningham C, Faiz O, Mortensen NJ. The economic impact of anastomotic leakage after anterior resections in English NHS hospitals: Are we adequately remunerating them? *Colorectal Dis* 2013; **15**: e190-e198 [PMID: 23331871 DOI: 10.1111/codi.12125]
 - 11 **Karliczek A**, Harlaar NJ, Zeebregts CJ, Wiggers T, Baas PC, van Dam GM. Surgeons lack predictive accuracy for anastomotic leakage in gastrointestinal surgery. *Int J Colorectal Dis* 2009; **24**: 569-576 [PMID: 19221768 DOI: 10.1007/s00384-009-0658-6]
 - 12 **Hyman N**, Manchester TL, Osler T, Burns B, Cataldo PA. Anastomotic leaks after intestinal anastomosis: It's later than you think. *Ann Surg* 2007; **245**: 254-258 [PMID: 17245179 DOI: 10.1097/01.sla.0000225083.27182.85]
 - 13 **McDermott FD**, Heeney A, Kelly ME, Steele RJ, Carlson GL, Winter DC. Systematic review of preoperative, intraoperative and postoperative risk factors for colorectal anastomotic leaks. *Br J Surg* 2015; **102**: 462-479 [PMID: 25703524 DOI: 10.1002/bjs.9697]
 - 14 **Choudhuri AH**, Uppal R, Kumar M. Influence of non-surgical risk factors on anastomotic leakage after major gastrointestinal surgery: Audit from a tertiary care teaching institute. *Int J Crit Illn Inj Sci* 2013; **3**: 246-249 [PMID: 24459621 DOI: 10.4103/2229-5151.124117]
 - 15 **Pommergaard HC**, Gessler B, Burcharth J, Angenete E, Haglind E, Rosenberg J. Preoperative risk factors for anastomotic leakage after resection for colorectal cancer: A systematic review and meta-analysis. *Colorectal Dis* 2014; **16**: 662-671 [PMID: 24655784 DOI: 10.1111/codi.12618]
 - 16 **Pettersson D**, Cedermarck B, Holm T, Radu C, Pahlman L, Glimelius B, Martling A. Interim analysis of the Stockholm III trial of preoperative radiotherapy regimens for rectal cancer. *Br J Surg* 2010; **97**: 580-587 [PMID: 20155787 DOI: 10.1002/bjs.6914]
 - 17 **Carmichael JC**, Keller DS, Baldini G, Bordeianou L, Weiss E, Lee L, Boutros M, McClane J, Feldman LS, Steele SR. Clinical Practice Guidelines for Enhanced Recovery After Colon and Rectal Surgery From the American Society of Colon and Rectal Surgeons and Society of American Gastrointestinal and Endoscopic Surgeons. *Dis Colon Rectum* 2017; **60**: 761-784 [PMID: 28682962 DOI: 10.1097/DCR.0000000000000883]
 - 18 **Greco M**, Capretti G, Beretta L, Gemma M, Pecorelli N, Braga M. Enhanced recovery program in colorectal surgery: A meta-analysis of randomized controlled trials. *World J Surg* 2014; **38**: 1531-1541 [PMID: 24368573 DOI: 10.1007/s00268-013-2416-8]
 - 19 **Contant CM**, Hop WC, van't Sant HP, Oostvogel HJ, Smeets HJ, Stassen LP, Neijenhuis PA, Idenburg FJ, Dijkhuis CM, Heres P, van Tets WF, Gerritsen JJ, Weidema WF. Mechanical bowel preparation for elective colorectal surgery: A multicentre randomised trial. *Lancet* 2007; **370**: 2112-2117 [PMID: 18156032 DOI: 10.1016/S0140-6736(07)61905-9]
 - 20 **Slim K**, Vicaut E, Launay-Savary MV, Contant C, Chipponi J. Updated systematic review and meta-analysis of randomized clinical trials on the role of mechanical bowel preparation before colorectal surgery. *Ann Surg* 2009; **249**: 203-209 [PMID: 19212171 DOI: 10.1097/SLA.0b013e318193425a]
 - 21 **Scarborough JE**, Mantyh CR, Sun Z, Migaly J. Combined Mechanical and Oral Antibiotic Bowel Preparation Reduces Incisional Surgical Site Infection and Anastomotic Leak Rates After Elective Colorectal Resection: An Analysis of Colectomy-Targeted ACS NSQIP. *Ann Surg* 2015; **262**: 331-337 [PMID: 26083870 DOI: 10.1097/SLA.0000000000001041]
 - 22 **Kiran RP**, Murray AC, Chiuzan C, Estrada D, Forde K. Combined preoperative mechanical bowel preparation with oral antibiotics significantly reduces surgical site infection, anastomotic leak, and ileus after colorectal surgery. *Ann Surg* 2015; **262**: 416-25; discussion 423-5 [PMID: 26258310 DOI: 10.1097/SLA.0000000000001416]
 - 23 **Althumairi AA**, Canner JK, Pawlik TM, Schneider E, Nagarajan N, Safar B, Efron JE. Benefits of Bowel Preparation Beyond Surgical Site Infection: A Retrospective Study. *Ann Surg* 2016; **264**: 1051-1057 [PMID: 26727098 DOI: 10.1097/SLA.0000000000001576]
 - 24 **Garfinkle R**, Abou-Khalil J, Morin N, Ghitulescu G, Vasilevsky CA, Gordon P, Demian M, Boutros M. Is There a Role for Oral Antibiotic Preparation Alone Before Colorectal Surgery? ACS-NSQIP Analysis by Coarsened Exact Matching. *Dis Colon Rectum* 2017; **60**: 729-737 [PMID: 28594723 DOI: 10.1097/DCR.0000000000000851]
 - 25 **Koller SE**, Bauer KW, Eggleston BL, Smith R, Philp MM, Ross HM, Esnaola NF. Comparative Effectiveness and Risks of Bowel Preparation Before Elective Colorectal Surgery. *Ann Surg* 2018; **267**: 734-742 [PMID: 28151800 DOI: 10.1097/SLA.0000000000002159]
 - 26 **Beyer-Berjot L**, Slim K. Colorectal surgery and preoperative bowel preparation: Aren't we drawing hasty conclusions? *Colorectal Dis* 2018; **20**: 955-958 [PMID: 30199608 DOI: 10.1111/codi.14415]
 - 27 **Devane LA**, Proud D, O'Connell PR, Panis Y. A European survey of bowel preparation in colorectal surgery. *Colorectal Dis* 2017; **19**: O402-O406 [PMID: 28975694 DOI: 10.1111/codi.13905]
 - 28 **Shogan BD**, Belogortseva N, Luong PM, Zaborin A, Lax S, Bethel C, Ward M, Muldoon JP, Singer M, An G, Umanskiy K, Konda V, Shakhsher B, Luo J, Klabbers R, Hancock LE, Gilbert J, Zaborina O, Alverdy JC. Collagen degradation and MMP9 activation by *Enterococcus faecalis* contribute to intestinal anastomotic leak. *Sci Transl Med* 2015; **7**: 286ra68 [PMID: 25947163 DOI: 10.1126/scitranslmed.3010658]
 - 29 **Krarup PM**, Eld M, Jorgensen LN, Hansen MB, Ågren MS. Selective matrix metalloproteinase inhibition increases breaking strength and reduces anastomotic leakage in experimentally obstructed colon. *Int J Colorectal Dis* 2017; **32**: 1277-1284 [PMID: 28717842 DOI: 10.1007/s00384-017-2857-x]
 - 30 **Rolandelli RH**, Buckmire MA, Bernstein KA. Intravenous butyrate and healing of colonic anastomoses in the rat. *Dis Colon Rectum* 1997; **40**: 67-70 [PMID: 9102264 DOI: 10.1007/BF02055684]
 - 31 **Mathew AJ**, Wann VC, Abraham DT, Jacob PM, Selvan BS, Ramakrishna BS, Nair AN. The effect of butyrate on the healing of colonic anastomoses in rats. *J Invest Surg* 2010; **23**: 101-104 [PMID: 20497012 DOI: 10.3109/08941930903469367]
 - 32 **Bloemen JG**, Schreinemacher MH, de Bruine AP, Buurman WA, Bouvy ND, Dejong CH. Butyrate enemas improve intestinal anastomotic strength in a rat model. *Dis Colon Rectum* 2010; **53**: 1069-1075 [PMID: 20551761 DOI: 10.1007/DCR.0b013e3181d881b7]
 - 33 **Levison ME**. Effect of colon flora and short-chain fatty acids on growth in vitro of *Pseudomonas aeruginosa* and *Enterobacteriaceae*. *Infect Immun* 1973; **8**: 30-35 [PMID: 4198102 DOI: 10.1007/BF01639659]
 - 34 **van Praagh JB**, de Goffau MC, Bakker IS, Harmsen HJ, Olinga P, Havenga K. Intestinal microbiota and

- anastomotic leakage of stapled colorectal anastomoses: A pilot study. *Surg Endosc* 2016; **30**: 2259-2265 [PMID: 26385781 DOI: 10.1007/s00464-015-4508-z]
- 35 **Stumpf M**, Klinge U, Wilms A, Zabrocki R, Rosch R, Junge K, Kronen C, Schumpelick V. Changes of the extracellular matrix as a risk factor for anastomotic leakage after large bowel surgery. *Surgery* 2005; **137**: 229-234 [PMID: 15674206 DOI: 10.1016/j.surg.2004.07.011]
 - 36 **Shogan BD**, Smith DP, Christley S, Gilbert JA, Zaborina O, Alverdy JC. Intestinal anastomotic injury alters spatially defined microbiome composition and function. *Microbiome* 2014; **2**: 35 [PMID: 25250176 DOI: 10.1186/2049-2618-2-35]
 - 37 **Dubinsky-Pertsov B**, Temkin E, Harbarth S, Fankhauser-Rodriguez C, Carevic B, Radovanovic I, Ris F, Kariv Y, Buchs NC, Schiffer E, Cohen Percia S, Nutman A, Fallach N, Klausner J, Carmeli Y; R-GNOSIS WP4 study group. Carriage of extended-spectrum beta-lactamase-producing Enterobacteriaceae and the risk of surgical site infection after colorectal surgery: A prospective cohort study. *Clin Infect Dis* 2018 [PMID: 30204851 DOI: 10.1093/cid/ciy768]
 - 38 **Brook I**, Walker RI, MacVittie TJ. Effect of antimicrobial therapy on bowel flora and bacterial infection in irradiated mice. *Int J Radiat Biol Relat Stud Phys Chem Med* 1988; **53**: 709-716 [PMID: 3283066 DOI: 10.1080/09553008814551081]
 - 39 **Olivas AD**, Shogan BD, Valuckaite V, Zaborin A, Belogortseva N, Musch M, Meyer F, Trimble WL, An G, Gilbert J, Zaborina O, Alverdy JC. Intestinal tissues induce an SNP mutation in *Pseudomonas aeruginosa* that enhances its virulence: Possible role in anastomotic leak. *PLoS One* 2012; **7**: e44326 [PMID: 22952955 DOI: 10.1371/journal.pone.0044326]
 - 40 **Juo YY**, Hyder O, Haider AH, Camp M, Lidor A, Ahuja N. Is minimally invasive colon resection better than traditional approaches? First comprehensive national examination with propensity score matching. *JAMA Surg* 2014; **149**: 177-184 [PMID: 24352653 DOI: 10.1001/jamasurg.2013.3660]
 - 41 **Solaini L**, Bazzocchi F, Cavaliere D, Avanzolini A, Cucchetti A, Ercolani G. Robotic versus laparoscopic right colectomy: An updated systematic review and meta-analysis. *Surg Endosc* 2018; **32**: 1104-1110 [PMID: 29218671 DOI: 10.1007/s00464-017-5980-4]
 - 42 **Abraha I**, Binda GA, Montedori A, Arezzo A, Cirocchi R. Laparoscopic versus open resection for sigmoid diverticulitis. *Cochrane Database Syst Rev* 2017; **11**: CD009277 [PMID: 29178125 DOI: 10.1002/14651858.CD009277.pub2]
 - 43 **Collinson FJ**, Jayne DG, Pigazzi A, Tsang C, Barrie JM, Edlin R, Garbett C, Guillou P, Holloway I, Howard H, Marshall H, McCabe C, Pavitt S, Quirke P, Rivers CS, Brown JM. An international, multicentre, prospective, randomised, controlled, unblinded, parallel-group trial of robotic-assisted versus standard laparoscopic surgery for the curative treatment of rectal cancer. *Int J Colorectal Dis* 2012; **27**: 233-241 [PMID: 21912876 DOI: 10.1007/s00384-011-1313-6]
 - 44 **Neutzling CB**, Lustosa SA, Proenca IM, da Silva EM, Matos D. Stapled versus handsewn methods for colorectal anastomosis surgery. *Cochrane Database Syst Rev* 2012; CD003144 [PMID: 22336786 DOI: 10.1002/14651858.CD003144.pub2]
 - 45 **Naumann DN**, Bhangu A, Kelly M, Bowley DM. Stapled versus handsewn intestinal anastomosis in emergency laparotomy: A systemic review and meta-analysis. *Surgery* 2015; **157**: 609-618 [PMID: 25731781 DOI: 10.1016/j.surg.2014.09.030]
 - 46 **Mann B**, Kleinschmidt S, Stremmel W. Prospective study of hand-sutured anastomosis after colorectal resection. *Br J Surg* 1996; **83**: 29-31 [PMID: 8653354 DOI: 10.1002/bjs.1800830108]
 - 47 **ESCP Cohort Studies and Audits Committee**. The 2017 European Society of Coloproctology (ESCP) international snapshot audit of left colon, sigmoid and rectal resections - study protocol. *Colorectal Dis* 2018; **20** Suppl 6: 5-12 [PMID: 30255639 DOI: 10.1111/codi.14377]
 - 48 **Braunschmid T**, Hartig N, Baumann L, Dauser B, Herbst F. Influence of multiple stapler firings used for rectal division on colorectal anastomotic leak rate. *Surg Endosc* 2017; **31**: 5318-5326 [PMID: 28634627 DOI: 10.1007/s00464-017-5611-0]
 - 49 **Lee S**, Ahn B, Lee S. The Relationship Between the Number of Intersections of Staple Lines and Anastomotic Leakage After the Use of a Double Stapling Technique in Laparoscopic Colorectal Surgery. *Surg Laparosc Endosc Percutan Tech* 2017; **27**: 273-281 [PMID: 28614172 DOI: 10.1097/SLE.0000000000000422]
 - 50 **Kim HJ**, Choi GS, Park JS, Park SY. Comparison of intracorporeal single-stapled and double-stapled anastomosis in laparoscopic low anterior resection for rectal cancer: A case-control study. *Int J Colorectal Dis* 2013; **28**: 149-156 [PMID: 23014975 DOI: 10.1007/s00384-012-1582-8]
 - 51 **Berho M**, Wexner SD, Botero-Anug AM, Pelled D, Fleshman JW. Histopathologic advantages of compression ring anastomosis healing as compared with stapled anastomosis in a porcine model: A blinded comparative study. *Dis Colon Rectum* 2014; **57**: 506-513 [PMID: 24608308 DOI: 10.1097/DCR.0000000000000009]
 - 52 **Masoomi H**, Luo R, Mills S, Carmichael JC, Senagore AJ, Stamos MJ. Compression anastomosis ring device in colorectal anastomosis: A review of 1,180 patients. *Am J Surg* 2013; **205**: 447-451 [PMID: 23290352 DOI: 10.1016/j.amjsurg.2012.03.013]
 - 53 **D'Hoore A**, Albert MR, Cohen SM, Herbst F, Matter I, Van Der Speeten K, Dominguez J, Rutten H, Muldoon JP, Bardakcioglu O, Senagore AJ, Ruppert R, Mills S, Stamos MJ, Pahlman L, Choman E, Wexner SD; COMPRES collaborative study group. COMPRES: A prospective postmarketing evaluation of the compression anastomosis ring CAR 27(TM)/ColonRing(TM). *Colorectal Dis* 2015; **17**: 522-529 [PMID: 25537083 DOI: 10.1111/codi.12884]
 - 54 **Slessor AA**, Pellino G, Shariq O, Cocker D, Kontovounisios C, Rasheed S, Tekkis PP. Compression versus hand-sewn and stapled anastomosis in colorectal surgery: A systematic review and meta-analysis of randomized controlled trials. *Tech Coloproctol* 2016; **20**: 667-676 [PMID: 27554096 DOI: 10.1007/s10151-016-1521-8]
 - 55 **Ris F**, Yeung T, Hompes R, Mortensen NJ. Enhanced Reality and Intraoperative Imaging in Colorectal Surgery. *Clin Colon Rectal Surg* 2015; **28**: 158-164 [PMID: 26491408 DOI: 10.1055/s-0035-1555007]
 - 56 **Allaix ME**, Lena A, Degiuli M, Arezzo A, Passera R, Mistrangelo M, Morino M. Intraoperative air leak test reduces the rate of postoperative anastomotic leak: Analysis of 777 laparoscopic left-sided colon resections. *Surg Endosc* 2019; **33**: 1592-1599 [PMID: 30203203 DOI: 10.1007/s00464-018-6421-8]
 - 57 **Wu Z**, van de Haar RC, Sparreboom CL, Boersema GS, Li Z, Ji J, Jeekel J, Lange JF. Is the intraoperative air leak test effective in the prevention of colorectal anastomotic leakage? A systematic review and meta-analysis. *Int J Colorectal Dis* 2016; **31**: 1409-1417 [PMID: 27294661 DOI: 10.1007/s00384-016-2616-4]
 - 58 **Li VK**, Wexner SD, Pulido N, Wang H, Jin HY, Weiss EG, Nogeauras JJ, Sands DR. Use of routine intraoperative endoscopy in elective laparoscopic colorectal surgery: Can it further avoid anastomotic

- failure? *Surg Endosc* 2009; **23**: 2459-2465 [PMID: [19301071](#) DOI: [10.1007/s00464-009-0416-4](#)]
- 59 **Hirst NA**, Tiernan JP, Millner PA, Jayne DG. Systematic review of methods to predict and detect anastomotic leakage in colorectal surgery. *Colorectal Dis* 2014; **16**: 95-109 [PMID: [23992097](#) DOI: [10.1111/codi.12411](#)]
- 60 **Vignali A**, Gianotti L, Braga M, Radaelli G, Malvezzi L, Di Carlo V. Altered microperfusion at the rectal stump is predictive for rectal anastomotic leak. *Dis Colon Rectum* 2000; **43**: 76-82 [PMID: [10813128](#) DOI: [10.1007/BF02237248](#)]
- 61 **GOLIGHER JC**. The blood-supply to the sigmoid colon and rectum with reference to the technique of rectal resection with restoration of continuity. *Br J Surg* 1949; **37**: 157-162 [PMID: [15395084](#) DOI: [10.1002/bjs.18003714604](#)]
- 62 **Novell JR**, Lewis AA. Peroperative observation of marginal artery bleeding: a predictor of anastomotic leakage. *Br J Surg* 1990; **77**: 137-138 [PMID: [2317669](#) DOI: [10.1002/bjs.1800770636](#)]
- 63 **Ambrosetti P**, Robert J, Mathey P, Rohner A. Left-sided colon and colorectal anastomoses: Doppler ultrasound as an aid to assess bowel vascularization. A prospective evaluation of 200 consecutive elective cases. *Int J Colorectal Dis* 1994; **9**: 211-214 [PMID: [7876727](#) DOI: [10.1007/BF00292253](#)]
- 64 **Karliczek A**, Benaron DA, Baas PC, Zeebregts CJ, Wiggers T, van Dam GM. Intraoperative assessment of microperfusion with visible light spectroscopy for prediction of anastomotic leakage in colorectal anastomoses. *Colorectal Dis* 2010; **12**: 1018-1025 [PMID: [19681979](#) DOI: [10.1111/j.1463-1318.2009.01944.x](#)]
- 65 **Jafari MD**, Lee KH, Halabi WJ, Mills SD, Carmichael JC, Stamos MJ, Pigazzi A. The use of indocyanine green fluorescence to assess anastomotic perfusion during robotic assisted laparoscopic rectal surgery. *Surg Endosc* 2013; **27**: 3003-3008 [PMID: [23404152](#) DOI: [10.1007/s00464-013-2832-8](#)]
- 66 **Ris F**, Liot E, Buchs NC, Kraus R, Ismael G, Belfontali V, Douissard J, Cunningham C, Lindsey I, Guy R, Jones O, George B, Morel P, Mortensen NJ, Hompes R, Cahill RA, Near-Infrared Anastomotic Perfusion Assessment Network VOIR. Multicentre phase II trial of near-infrared imaging in elective colorectal surgery. *Br J Surg* 2018; **105**: 1359-1367 [PMID: [29663330](#) DOI: [10.1002/bjs.10844](#)]
- 67 **Jafari MD**, Wexner SD, Martz JE, McLemore EC, Margolin DA, Sherwinter DA, Lee SW, Senagore AJ, Phelan MJ, Stamos MJ. Perfusion assessment in laparoscopic left-sided/anterior resection (PILLAR II): A multi-institutional study. *J Am Coll Surg* 2015; **220**: 82-92.e1 [PMID: [25451666](#) DOI: [10.1016/j.jamcollsurg.2014.09.015](#)]
- 68 **Blanco-Colino R**, Espin-Basany E. Intraoperative use of ICG fluorescence imaging to reduce the risk of anastomotic leakage in colorectal surgery: A systematic review and meta-analysis. *Tech Coloproctol* 2018; **22**: 15-23 [PMID: [29230591](#) DOI: [10.1007/s10151-017-1731-8](#)]
- 69 **Ishii M**, Tanaka E, Imaizumi T, Sugio Y, Sekka T, Tanaka M, Yasuda M, Fukuyama N, Shinozaki Y, Hyodo K, Tanioka K, Mochizuki R, Kawai T, Mori H, Makuuchi H. Local VEGF administration enhances healing of colonic anastomoses in a rabbit model. *Eur Surg Res* 2009; **42**: 249-257 [PMID: [19346745](#) DOI: [10.1159/000210671](#)]
- 70 **Lindgren R**, Hallböök O, Rutegård J, Sjö Dahl R, Matthiessen P. What is the risk for a permanent stoma after low anterior resection of the rectum for cancer? A six-year follow-up of a multicenter trial. *Dis Colon Rectum* 2011; **54**: 41-47 [PMID: [21160312](#) DOI: [10.1007/DCR.0b013e3181fd2948](#)]
- 71 **Montedori A**, Cirocchi R, Farinella E, Sciannameo F, Abraha I. Covering ileo- or colostomy in anterior resection for rectal carcinoma. *Cochrane Database Syst Rev* 2010; CD006878 [PMID: [20464746](#) DOI: [10.1002/14651858.CD006878.pub2](#)]
- 72 **Nurkin S**, Kakarla VR, Ruiz DE, Cance WG, Tiszenkel HI. The role of faecal diversion in low rectal cancer: A review of 1791 patients having rectal resection with anastomosis for cancer, with and without a proximal stoma. *Colorectal Dis* 2013; **15**: e309-e316 [PMID: [23586660](#) DOI: [10.1111/codi.12248](#)]
- 73 **Wu SW**, Ma CC, Yang Y. Role of protective stoma in low anterior resection for rectal cancer: A meta-analysis. *World J Gastroenterol* 2014; **20**: 18031-18037 [PMID: [25548503](#) DOI: [10.3748/wjg.v20.i47.18031](#)]
- 74 **Hüser N**, Michalski CW, Erkan M, Schuster T, Rosenberg R, Kleeff J, Friess H. Systematic review and meta-analysis of the role of defunctioning stoma in low rectal cancer surgery. *Ann Surg* 2008; **248**: 52-60 [PMID: [18580207](#) DOI: [10.1097/SLA.0b013e318176bf65](#)]
- 75 **Denost Q**, Rouanet P, Faucheron JL, Panis Y, Meunier B, Cotte E, Meurette G, Kirzin S, Sabbagh C, Loriau J, Benoist S, Mariette C, Sieleznoff I, Lelong B, Mauvais F, Romain B, Barussaud ML, Germain C, Picat MQ, Rullier E, Laurent C; French Research Group of Rectal Cancer Surgery (GRECCAR). To Drain or Not to Drain Infraperitoneal Anastomosis After Rectal Excision for Cancer: The GRECCAR 5 Randomized Trial. *Ann Surg* 2017; **265**: 474-480 [PMID: [27631776](#) DOI: [10.1097/SLA.0000000000001991](#)]
- 76 **Guerra F**, Giuliani G, Coletta D, Boni M, Rondelli F, Bianchi PP, Coratti A. A Meta-Analysis of Randomized Controlled Trials on the Use of Suction Drains Following Rectal Surgery. *Dig Surg* 2018; **35**: 482-490 [PMID: [29232658](#) DOI: [10.1159/000485139](#)]
- 77 **Jesus EC**, Karliczek A, Matos D, Castro AA, Atallah AN. Prophylactic anastomotic drainage for colorectal surgery. *Cochrane Database Syst Rev* 2004; CD002100 [PMID: [15495028](#) DOI: [10.1002/14651858.CD002100.pub2](#)]
- 78 **Yang Y**, Shu Y, Su F, Xia L, Duan B, Wu X. Prophylactic transanal decompression tube versus non-prophylactic transanal decompression tube for anastomotic leakage prevention in low anterior resection for rectal cancer: A meta-analysis. *Surg Endosc* 2017; **31**: 1513-1523 [PMID: [27620910](#) DOI: [10.1007/s00464-016-5193-2](#)]
- 79 **Chen H**, Cai HK, Tang YH. An updated meta-analysis of transanal drainage tube for prevention of anastomotic leak in anterior resection for rectal cancer. *Surg Oncol* 2018; **27**: 333-340 [PMID: [30217286](#) DOI: [10.1016/j.suronc.2018.05.018](#)]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

