



All

Images

Videos

翻译成中文

关闭取词

2,670 Results

Any time ▾

Partial Second Toe Pulp Free Flap for Fingertip Reconstruction

https://www.researchgate.net/publication/5535386_Partial_Second...

Fingertip defects are reconstructed using a variety of techniques 1 2345678 . For the past 15 years, our first-line treatment of such **defects** has been the partial **second toe** pulp free **flap** because ...

The Tibial Second Toe Vascularized Neurocutaneous Free ...

https://www.researchgate.net/publication/6528349_The_Tibial_Second...

The **Tibial Second Toe** Vascularized Neurocutaneous Free **Flap** for Major Digital Nerve **Defects** Article in The Journal Of Hand Surgery 32(2):209-17 · March 2007 with 15 Reads DOI: 10.1016/j.jhsa.2006 ...

Innervated dorsoradial perforator free flap: A reliable ...

<https://www.sciencedirect.com/science/article/pii/S1748681517302164>

Introduction. **Fingertip defects** reconstruction is challenging. 1 Cosmesis and sensitivity are essential for good reconstructive outcomes. Traditional methods include cross-**finger flap** 2 and digital artery perforator **flap**. 3 Current literature reports various free **flap** options such as **toe flap**, 4 thenar **flap**, 5 lateral arm **flap**, 6 posterior inter-osseous **flap** 7 venous **flap**, 8 dorsoulnar ...

Cited by: 2

Author: Dajiang Song, Georgios Pafitanis, Peng Y...

Publish Year: 2017

Innervated dorsoradial perforator free flap: A reliable ...

[https://www.jprasurg.com/article/S1748-6815\(17\)30216-4/fulltext](https://www.jprasurg.com/article/S1748-6815(17)30216-4/fulltext)

Innervated dorsoradial perforator free **flap**: A reliable **supermicrosurgery** **fingertip** reconstruction technique ... Kim, H.Y., Rha, E.Y. et al. Esthetically successful **fingertip** reconstruction using a **second toe** pulp free **flap** with a **split** thickness skin graft and ... G. and Suzuki, K. Arterialized venous **flap** for treating **multiple** skin **defects** of ...

Cited by: 2

Author: Dajiang Song, Georgios Pafitanis, Peng Y...

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 47518

Manuscript Type: CASE REPORT

Supermicrosurgery in fingertip defects-split tibial flap of the second toe to reconstruct multiple fingertip defects: A case report

Wang KL *et al.* Supermicrosurgery in fingertip defects

Ke-Lie Wang, Zi-Qing Zhang, Joseph A Buckwalter, Yi Yang

Match Overview

1	Internet 12 words crawled on 19-Apr-2010 www.nipne.ro	1%
2	Internet 11 words crawled on 16-Oct-2018 www.thieme-connect.de	1%
3	Internet 11 words crawled on 25-Nov-2017 emedicine.staging.medscape.com	1%
4	Internet 10 words crawled on 19-Jan-2015 pt.slideshare.net	1%

[国内版](#)[国际版](#)[All](#)[Images](#)[Videos](#)[翻译成中文](#)[关闭取词](#)

1,750 Results

Any time ▾

Innervated dorsoradial perforator free flap: A reliable ...

<https://www.sciencedirect.com/science/article/pii/S1748681517302164>

Introduction. **Fingertip defects** reconstruction is challenging. 1 Cosmesis and sensitivity are essential for good reconstructive outcomes. Traditional methods include cross-**finger flap** 2 and digital artery perforator **flap**. 3 Current literature reports various free **flap** options such as **toe flap**, 4 thenar **flap**, 5 lateral arm **flap**, 6 posterior inter-osseous **flap** 7 venous **flap**, 8 dorsolunar ...

Cited by: 2

Author: Dajiang Song, Georgios Pafitanis, Peng...

Publish Year: 2017

An Evolutionary Perspective on the History of Flap ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4160825>

In 1897, Carl Nicoladoni applied the distant **flap** concept with a pedicled **thoracoepigastric flap** to cover a degloved **thumb**. 28,29 Nicoladoni later further developed the distant **flap** concept with application to **finger reconstruction** in his description of a pedicled **second toe** to **thumb** transfer in 1900. 29 In 1917, another distant **flap** for **finger reconstruction**, the **cross-hand flap** (substitution of **ring finger** for reconstruction of a **thumb**...

Cited by: 2

Author: Frank Fang, Kevin C. Chung

Publish Year: 2014

Innervated dorsoradial perforator free flap: A reliable ...

https://www.researchgate.net/publication/317246902_Innervated_dorsoradial_perforator...

Innervated dorsoradial perforator free **flap**: A reliable **supermicrosurgery fingertip** reconstruction technique ... **defects**. We **report** the results of **second toe** transfer for 1-stage **finger** ...

Fingertip Reconstruction Using Free Toe Tissue Transfer ...

europepmc.org/articles/PMC3474414

Sep 01, 2012 · Index **fingertip** reconstruction using **second toe** tissue. Right index **fingertip** was reconstructed with **second toe** tissue in the **case** 1 patient. (A) Preoperative state. Initially a composite graft was performed but failed. (B) A 2.0×2.0 cm 2 sized **flap** was designed on the mediolateral aspect of the left **second toe**. (C) A **toe flap**, based only on the ...

Published

in: Archives of Plastic Surgery · 2012

Authors: Won Young Yoon · Byung Il Lee



国内版

国际版

Supermicrosurgery in fingertip defects-split tibial flap of the second toe to recons



All

Images

Videos

翻译成中文

开启取词

1,800 Results

Any time ▾

Innervated dorsoradial perforator free flap: A reliable ...

<https://www.sciencedirect.com/science/article/pii/S1748681517302164>

Introduction. **Fingertip defects** reconstruction is challenging. 1 Cosmesis and sensitivity are essential for good reconstructive outcomes. Traditional methods include cross-**finger flap** 2 and digital artery perforator **flap**. 3 Current literature reports various free **flap** options such as **toe flap**, 4 thenar **flap**, 5 lateral arm **flap**, 6 posterior inter-osseous **flap** 7 venous **flap**, 8 dorsoulnar ...

Cited by: 2

Author: Dajiang Song, Georgios Pafitanis, Peng Y...

Publish Year: 2017

Innervated dorsoradial perforator free flap: A reliable ...

https://www.researchgate.net/publication/317246902_Innervated_dorsoradial_perforator...

Innervated dorsoradial perforator free **flap**: A reliable **supermicrosurgery fingertip** reconstruction technique ... **defects**. We report the results of **second toe** transfer for 1-stage **finger** ...

An Evolutionary Perspective on the History of Flap ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4160825>

In 1897, Carl Nicoladoni applied the distant **flap** concept with a pedicled **thoracoepigastric flap** to cover a degloved **thumb**. 28,29 Nicoladoni later further developed the distant **flap** concept with application to **finger reconstruction** in his description of a pedicled **second toe** to **thumb** transfer in 1900. 29 In 1917, another distant **flap** for **finger reconstruction**, the **cross-hand flap** (substitution of **ring finger** for reconstruction of a **thumb**...

Cited by: 2

Author: Frank Fang, Kevin C. Chung

Publish Year: 2014

aps :: Archives of Plastic Surgery

<https://www.e-aps.org/m/journal/view.php?number=93> ▾

Right index **fingertip** was reconstructed with **second toe** tissue in the **case 1** patient. (A) Preoperative state. Initially a composite graft was performed but failed. (B) A 2.0×2.0 cm 2 sized **flap** was designed on the mediovolar aspect of the left **second toe**. (C) A **toe flap**...

Fingertip Reconstruction Using Free Toe Tissue Transfer ...

europepmc.org/articles/PMC3474414

Sep 01, 2012 · Index **fingertip** reconstruction using **second toe** tissue. Right index **fingertip** was reconstructed with **second toe** tissue in the **case 1** patient. (A) Preoperative state. Initially a composite graft was performed but failed. (B) A 2.0×2.0 cm 2 sized **flap** was designed on the mediovolar aspect of the left **second toe**. (C) A **toe flap**, based only on the ...

Published in: Archives of Plastic Surgery · 2012

Authors: Won Young Yoon · Byung Il Lee

Affiliation: Korea University

About: Bioinformatics · Finger injury · Text mining