

# World Journal of *Clinical Cases*

*World J Clin Cases* 2019 August 6; 7(15): 1908-2133



**REVIEW**

- 1908** Bone alterations in inflammatory bowel diseases  
*Sgambato D, Gimigliano F, De Musis C, Moretti A, Toro G, Ferrante E, Miranda A, De Mauro D, Romano L, Iolascon G, Romano M*

**MINIREVIEWS**

- 1926** Extrahepatic hepcidin production: The intriguing outcomes of recent years  
*Daher R, Lefebvre T, Puy H, Karim Z*
- 1937** Neoadjuvant endocrine therapy: A potential strategy for ER-positive breast cancer  
*Yao LT, Wang MZ, Wang MS, Yu XT, Guo JY, Sun T, Li XY, Xu YY*

**ORIGINAL ARTICLE****Basic Study**

- 1954** Vestigial like family member 3 is a novel prognostic biomarker for gastric cancer  
*Zhang LH, Wang Z, Li LH, Liu YK, Jin LF, Qi XW, Zhang C, Wang T, Hua D*

**Retrospective Study**

- 1964** HER2 heterogeneity is a poor prognosticator for HER2-positive gastric cancer  
*Kaito A, Kuwata T, Tokunaga M, Shitara K, Sato R, Akimoto T, Kinoshita T*

**Case Control Study**

- 1978** Changes in corneal endothelial cell density in patients with primary open-angle glaucoma  
*Yu ZY, Wu L, Qu B*

**Observational Study**

- 1986** Myocardial bridge-related coronary heart disease: Independent influencing factors and their predicting value  
*Zhao DH, Fan Q, Ning JX, Wang X, Tian JY*
- 1996** Clinical significance and role of up-regulation of SERPINA3 expression in endometrial cancer  
*Zhou ML, Chen FS, Mao H*
- 2003** Evaluation of right ventricular volume and systolic function in normal fetuses using intelligent spatiotemporal image correlation  
*Sun JX, Cai AL, Xie LM*

- 2013** Correlation between intracoronary thrombus components and coronary blood flow after percutaneous coronary intervention for acute myocardial infarction at different onset time  
*Zhang MJ, Liu X, Liu LH, Li N, Zhang N, Wang YQ, Sun XJ, Huang PH, Yin HM, Liu YH, Zheng H*

**META-ANALYSIS**

- 2022** Performance of common imaging techniques *vs* serum biomarkers in assessing fibrosis in patients with chronic hepatitis B: A systematic review and meta-analysis  
*Xu XY, Wang WS, Zhang QM, Li JL, Sun JB, Qin TT, Liu HB*

**CASE REPORT**

- 2038** Acute bleeding after argon plasma coagulation for weight regain after gastric bypass: A case report  
*Moura DTHD, Sachdev AH, Lu PW, Ribeiro IB, Thompson CC*
- 2044** Left colonic metastasis from primary hepatocellular carcinoma: A case report  
*Tagliabue F, Burati M, Chiarelli M, Marando A, Simone MD, Cioffi U*
- 2049** ALK-positive anaplastic large cell lymphoma presenting multiple lymphomatous polyposis: A case report and literature review  
*Saito M, Izumiyama K, Ogasawara R, Mori A, Kondo T, Tanaka M, Morioka M, Miyashita K, Tanino M*
- 2058** Modified Tong Xie Yao Fang relieves solitary rectal ulcer syndrome: A case report  
*Zhang LL, Hao WS, Xu M, Li C, Shi YY*
- 2065** Hydrogen gas therapy induced shrinkage of metastatic gallbladder cancer: A case report  
*Chen JB, Pan ZB, Du DM, Qian W, Ma YY, Mu F, Xu KC*
- 2075** Giant nonfunctional ectopic adrenocortical carcinoma on the anterior abdominal wall: A case report  
*Zhou DK, Liu ZH, Gao BQ, Wang WL*
- 2081** Oncogenic osteomalacia caused by a phosphaturic mesenchymal tumor of the femur: A case report  
*Tang D, Wang XM, Zhang YS, Mi XX*
- 2087** Gastric duplication cyst mimicking large cystic lymphangioma in an adult: A rare case report and review of the literature  
*Xv FY, Sun A, Gan Y, Hu HJ*
- 2094** Endometriosis of the duplex appendix: A case report and review of the literature  
*Zhu MY, Fei FM, Chen J, Zhou ZC, Wu B, Shen YY*
- 2103** Fever and neck pain after pacemaker lead extraction: A case report  
*Wang SX, Bai J, Ma R, Lan RF, Zheng J, Xu W*

- 2110** c.753\_754delAG, a novel *CFTR* mutation found in a Chinese patient with cystic fibrosis: A case report and review of the literature  
*Wang YQ, Hao CL, Jiang WJ, Lu YH, Sun HQ, Gao CY, Wu M*
- 2120** Common iliac artery occlusion with small intestinal transection caused by blunt abdominal trauma: A case report and review of the literature  
*Zhou YX, Ji Y, Chen J, Yang X, Zhou Q, Lv J*
- 2128** Percutaneous coronary intervention for ostial lesions of the left main stem in a patient with congenital single left coronary artery: A case report  
*Wu Q, Li ZZ, Yue F, Wei F, Zhang CY*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Shu-Pin Huang, MD, Associate Professor, Attending Doctor, Director, Department of Urology, Kaohsiung Medical University, Kaohsiung 807, Taiwan

**AIMS AND SCOPE**

*World Journal of Clinical Cases* (*World J Clin Cases*, *WJCC*, online ISSN 2307-8960, DOI: 10.12998) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

The primary task of *WJCC* is to rapidly publish high-quality Case Report, Clinical Management, Editorial, Field of Vision, Frontier, Medical Ethics, Original Articles, Meta-Analysis, Minireviews, and Review, in the fields of allergy, anesthesiology, cardiac medicine, clinical genetics, clinical neurology, critical care, dentistry, dermatology, emergency medicine, endocrinology, family medicine, gastroenterology and hepatology, etc.

**INDEXING/ABSTRACTING**

The *WJCC* is now indexed in PubMed, PubMed Central, Science Citation Index Expanded (also known as SciSearch®), and Journal Citation Reports/Science Edition. The 2019 Edition of Journal Citation Reports cites the 2018 impact factor for *WJCC* as 1.153 (5-year impact factor: N/A), ranking *WJCC* as 99 among 160 journals in Medicine, General and Internal (quartile in category Q3).

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Responsible Electronic Editor: *Yan-Xia Xing*  
 Proofing Production Department Director: *Yun-Xiaojuan Wu*

**NAME OF JOURNAL**  
*World Journal of Clinical Cases*

**ISSN**  
 ISSN 2307-8960 (online)

**LAUNCH DATE**  
 April 16, 2013

**FREQUENCY**  
 Semimonthly

**EDITORS-IN-CHIEF**  
 Dennis A Bloomfield, Sandro Vento

**EDITORIAL BOARD MEMBERS**  
<https://www.wjnet.com/2307-8960/editorialboard.htm>

**EDITORIAL OFFICE**  
 Jin-Lei Wang, Director

**PUBLICATION DATE**  
 August 6, 2019

**COPYRIGHT**  
 © 2019 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**  
<https://www.wjnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**  
<https://www.wjnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**  
<https://www.wjnet.com/bpg/gerinfo/240>

**PUBLICATION MISCONDUCT**  
<https://www.wjnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**  
<https://www.wjnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**  
<https://www.wjnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**  
<https://www.f6publishing.com>

## Basic Study

**Vestigial like family member 3 is a novel prognostic biomarker for gastric cancer**

Li-Hua Zhang, Zhuo Wang, Long-Hai Li, Yan-Kui Liu, Lin-Fang Jin, Xiao-Wei Qi, Chun Zhang, Teng Wang, Dong Hua

**ORCID number:** Li-Hua Zhang (0000-0002-0931-5039); Zhuo Wang (0000-0001-8206-6035); Long-Hai Li (0000-0003-4387-8932); Yan-Kui Liu (0000-0003-4387-8942); Lin-Fang Jin (0000-0002-2487-5886); Xiao-Wei Qi (0000-0001-7761-5095); Chun Zhang (0000-0001-6422-3367); Teng Wang (0000-0001-6092-9522); Dong Hua (0000-0001-5569-1708).

**Author contributions:** Zhang LH and Wang Z contributed equally to this work; Wang T and Hua D designed the research; Li LH, Liu YK, Jin LF, Qi XW, and Zhang C performed the research; Wang T and Hua D contributed new reagents and analytic tools; Li LH, Zhang LH, and Wang Z analyzed the data; Zhang LH and Wang Z wrote the paper.

**Supported by** the Natural Science Foundation of Jiangsu Province, No. BK20171150; the National Natural Science Foundation of China, No. 81502042; Research Project of Health and Family Planning Commission of Wuxi, No. Q201758; and Nanchang Hongda Jianghua Educational Foundation.

**Institutional review board**

**statement:** This study was reviewed and approved by Affiliated Hospital of Jiangnan University Institutional Review Board.

**Informed consent statement:** Any personal item or information will not be published without explicit consent from the involved persons.

**Li-Hua Zhang, Dong Hua,** School of Pharmaceutical Sciences, Jiangnan University, Wuxi 214122, Jiangsu Province, China

**Li-Hua Zhang, Teng Wang, Dong Hua,** Department of Oncology, Affiliated Hospital of Jiangnan University, Wuxi 214062, Jiangsu Province, China

**Zhuo Wang,** Department of Geriatrics, Wuxi People's Hospital Affiliated to Nanjing Medical University, Wuxi 214023, Jiangsu Province, China

**Long-Hai Li, Chun Zhang,** Wuxi Medical College, Jiangnan University, Wuxi 214122, Jiangsu Province, China

**Yan-Kui Liu, Lin-Fang Jin, Xiao-Wei Qi,** Department of Pathology, Affiliated Hospital of Jiangnan University, Wuxi 214062, Jiangsu Province, China

**Corresponding author:** Dong Hua, MD, PhD, Chief Doctor, Professor, Department of Oncology, Affiliated Hospital of Jiangnan University, No. 200, Huihe Road, Wuxi 214062, Jiangsu Province, China. [wx89211@163.com](mailto:wx89211@163.com)

**Telephone:** +86-510-88682109

**Fax:** +86-510-85808820

**Abstract****BACKGROUND**

Vestigial like family member 3 (VGLL3) is associated with the prognosis of epithelial ovarian cancer and soft tissue sarcoma, but its role in gastric cancer (GC) is unclear.

**AIM**

To explore the expression pattern and clinical significance of VGLL3 in GC.

**METHODS**

Integrative analysis was performed on the GC transcriptome profiles and survival information deposited in the ONCOMINE, GEPIA, and ONCOLNC databases. The expression levels of VGLL3 mRNA and protein were analyzed in the freshly resected tumor and normal gastric tissues from GC patients by quantitative RT-PCR and Western blot, respectively. In addition, the *in situ* expression of VGLL3 in the GC tissues was determined by immunohistochemistry (IHC), and the patients were accordingly classified into the high and low expression groups. The correlation of VGLL3 expression status with patient prognosis was then determined by univariate and multivariate Cox

**Conflict-of-interest statement:** The authors report no conflicts of interest in this work.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Invited Manuscript

**Received:** March 22, 2019

**Peer-review started:** March 22, 2019

**First decision:** May 13, 2019

**Revised:** June 22, 2019

**Accepted:** July 3, 2019

**Article in press:** July 3, 2019

**Published online:** August 6, 2019

**P-Reviewer:** Bang YJ, Kruszewski WJ

**S-Editor:** Cui LJ

**L-Editor:** Wang TQ

**E-Editor:** Liu JH



regression analyses.

## RESULTS

Analysis of the ONCOMINE and GEPIA databases showed that VGLL3 was significantly up-regulated in GC tissues ( $P = 0.003$ ), and associated with the tumor TNM stage ( $P = 0.0163$ ). The high VGLL3 expression group had a significantly worse prognosis compared to the low expression group, as per both GEPIA ( $P = 0.0057$ ) and ONCOLNC ( $P = 0.01$ ). The bioinformatics results were validated by the significantly higher VGLL3 mRNA and protein levels in the GC tissues compared to the adjacent normal tissues ( $P < 0.001$ ) in a cohort of 30 GC patients. Furthermore, high *in situ* expression of VGLL3 protein was associated with more advanced N and TNM stages and HER2 mutation ( $P < 0.05$ ) in a cohort of 172 patients. Kaplan-Meier analysis showed that the high VGLL3 expression group had a worse prognosis compared to the low expression group ( $P = 0.019$ ). Multivariate analysis showed that VGLL3 expression status was an independent risk factor for prognosis. In addition, the prognostic risk model nomogram showed that VGLL3 was the most important indicator, with an area under the receiver operating characteristic (ROC) curve (AUC) of 0.613 for 3-year survival and 0.706 for 5-year survival. Finally, the protein interaction network analysis revealed that VGLL3 is likely involved in the Hippo signaling pathway.

## CONCLUSION

VGLL3 is overexpressed in GC tissues and associated with a poor prognosis, indicating its potential as a novel prognosis biomarker and therapeutic target for GC.

**Key words:** Vestigial like family member 3; Stomach adenocarcinoma; HER2 mutation; Gastric cancer; Bioinformatics analysis

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** The present study for the first time revealed the expression of vestigial like family member 3 (VGLL3) in gastric cancer (GC) and its correlation with HER2 mutation. Overall, the findings of the present study suggest that VGLL3 is a novel prognostic biomarker for GC and highlight the significance of VGLL3 as a promising therapeutic target for GC.

**Citation:** Zhang LH, Wang Z, Li LH, Liu YK, Jin LF, Qi XW, Zhang C, Wang T, Hua D. Vestigial like family member 3 is a novel prognostic biomarker for gastric cancer. *World J Clin Cases* 2019; 7(15): 1954-1963

**URL:** <https://www.wjgnet.com/2307-8960/full/v7/i15/1954.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v7.i15.1954>

## INTRODUCTION

Gastric cancer (GC) is a frequently occurring malignancy of the digestive tract<sup>[1]</sup>, and it was the fifth most commonly diagnosed and the third most common cause of cancer-related mortality worldwide in 2018<sup>[2]</sup>. Lack of early diagnosis and ineffective treatment options for the advanced stages are the primary reasons for the dismal 5-year survival rate of GC<sup>[3]</sup>. Therefore, it is imperative to identify novel biomarkers for early diagnosis and accurate prediction of prognosis. Vestigial like family member 3 (VGLL3) is a member of the vestigial like family of proteins<sup>[4]</sup>, and is associated with epithelial ovarian cancer<sup>[5]</sup> and soft tissue sarcoma<sup>[6]</sup>. The aim of this study was to determine the expression status and prognostic utility of VGLL3 in GC. To this end, we mined the transcriptomic data of GC from the ONCOMINE and GEPIA databases, and determined its correlation with the survival data from GEPIA and ONCOLINC. VGLL3 levels were upregulated in GC samples and associated with a poor prognosis. The bioinformatics data were successfully validated on the tumor and normal gastric tissues resected from GC patients. Based on the VGLL3 levels, the patients were stratified into the high and low expression groups, and a prognosis prediction model was established on the basis of VGLL3 status and clinical data. Our findings show a

strong prognostic role of VGLL3 in GC, which can potentially translate to clinical applications.

## MATERIALS AND METHODS

### *Patients and tissue samples*

A total of 172 tissue samples were obtained from gastric adenocarcinoma patients who underwent major surgery at the Affiliated Hospital of Jiangnan University between 2009 and 2012. The tissues were immediately fixed in formalin and embedded in paraffin for further analysis. In addition, fresh samples were dissected from 30 patients at the Affiliated Hospital of Jiangnan University in November 2018, and immediately stored in liquid nitrogen for molecular analysis. The mean follow-up duration was 43.1 mo and ranged from 0.2 to 86 mo. None of the patients received chemotherapy or radiotherapy before surgery<sup>[7]</sup>. The tumor stage classification was determined by three pathologists who were blinded to the patient data according to the guidelines of the American Joint Committee on Cancer (AJCC). Freshly resected tissues were obtained from 202 patients, and were fixed in formalin and embedded in paraffin. In addition, tissue specimens from 30 patients were flash frozen in liquid nitrogen. All participating clinicians and patients provided written informed consent.

### *Bioinformatics analysis*

The VGLL3 mRNA levels in the GC and normal gastric tissues were determined by integrated analysis of the Oncomine database ([www.oncomine.org](http://www.oncomine.org))<sup>[8]</sup> using Coexpedia ([www.coexpedia.org](http://www.coexpedia.org))<sup>[9]</sup>. ONCOLINC (<http://www.oncolnc.org/cancer/>) and Coexpedia were used to analyze the prognostic value of WISP1 in STAD. The STRING database (<http://www.string-db.org>) was utilized to construct the protein-protein interaction (PPI) network<sup>[10]</sup>.

### *RNA isolation and RT-qPCR*

Total RNA was extracted from frozen tissue samples using Trizol reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions, and reverse transcribed using the PrimeScript RT-PCR kit (Takara, Japan)<sup>[11]</sup>. RT-qPCR was performed on the ABI 7500 RealTime PCR System (Applied Biosystems, Inc. USA) using SYBR Green Master Mix (Takara, Japan), and the VGLL3 levels were normalized to  $\beta$ -actin. The following primers were used: Forward primer, 5'-CCAACTACAGTCACC-TCTGCTAC-3' and reverse primer, 5'-ACCACGGTGATTCCTTACTCTTG-3' for VGLL3; forward primer, 5'-CCTGTGGCATCCACGAAACT-3' and reverse primer, 5'-GAAGCATTGCG GTGGACGAT-3' for  $\beta$ -actin. All reactions were performed in triplicates, and the  $2^{-\Delta\Delta Ct}$  method<sup>[12]</sup> was used to quantify the relative expression levels of VGLL3.

### *Western blot analysis*

Total protein was extracted from GC and para-cancerous tissues using the RIPA lysis buffer (Pierce, Thermo Scientific, Cramlington, United Kingdom)<sup>[13]</sup>, and quantified with the enhanced BCA protein assay kit (KeyGEN BioTECH, Jiangsu, China). Equal amount (40 mg) of proteins per sample were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SD-PAGE), and then transferred to polyvinylidene difluoride membranes (Bio-Rad, Hercules, CA, United States). After blocking with 5% non-fat milk for 1 h at room temperature (RT), the membranes were incubated overnight with anti-VGLL3 (ab83555, Abcam) and  $\beta$ -actin (ab8226, Abcam) primary antibodies at 4 °C. After washing thrice with TBST, the membranes were incubated with HRP-conjugated secondary antibody (1:1000) for 1 h at RT. The protein bands were visualized with an ECL chemiluminescence system after short exposure to X-ray films (Kodak, Japan). Densitometric analysis was performed with Image Pro-Plus software, and the relative expression levels of VGLL3 were normalized to tubulin.

### *Immunohistochemistry (IHC)*

The formalin-fixed tissues were dehydrated, embedded in paraffin, and cut into 5  $\mu$ m thick sections. IHC was performed according to the manufacturer's protocol<sup>[14]</sup>. Briefly, the sections were heated in citrate buffer in a microwave, cooled, and incubated overnight with anti-VGLL3 antibody (1:50, clone PA5-68441, Invitrogen) at 4 °C. After labeling with the secondary antibody for 1 hour at RT, color was developed using liquid DAB substrate. Three pathologists blinded to the patient identity observed and graded VGLL3 positivity as 0, 1 (1%-29% positively-stained cells), 2 (30%-69%), or 3 (70%-100%), and the staining intensity as 0 (negative), 1 (weak staining), 2 (moderate staining), or 3 (strong staining). The immunoreactive

score (IRS) was then calculated for each sample by multiplying the staining intensity and positivity scores, and graded as: 0-1, “-”; 2-3, “+”; 4-6, “++”; and 6-9, “+++”<sup>[15]</sup>. Based on the IRS, the samples were stratified into the high (4-9) and low (0-3) VGLL3 expression groups.

### Fluorescence in situ hybridization (FISH)

The tissue sections were treated with denaturing solution to denature the DNA into single strands, and hybridized with the PathVysion probes (No.36-161060, PathVysion HER-2DNA Probe Kit). After washing the unbound probe with DNA, the sections were counterstained with the nuclear dye DAPI (4', 6 diamidino-2-phenylindole). Positive LSI HER-2/neu and CEP 17 signals were counted under a fluorescence microscope, and the ratio of the copy number of HER-2/neu gene to that of chromosome 17 was calculated.

### Statistical analysis

Statistical analyses were performed with R\*64 version 3.5.2 software and Graphpad 6.01. HER2 status (positive or negative) was designated on the basis of the combined FISH and IHC results as per the American Society of Clinical Oncology/College of American Pathologists Clinical Practice Guidelines. VGLL3 expression levels in the tumor and normal samples were compared by the Student's *t*-test, and the association between VGLL3 and clinico-pathological features was assessed by the chi-square test. The overall survival (OS) curve was plotted by the Kaplan-Meier method and analyzed by the log-rank test. Univariate and multivariate analyses of the prognostic factors were performed using the Cox proportional hazard regression model. A nomogram was formulated based on the results of the multivariate analysis with the package of rms in R version 3.5.2 (<http://www.r-project.org/>)<sup>[16]</sup>. The receiver operating characteristic (ROC) curve was plotted to determine the sensitivity and specificity of the VGLL3-based prognostic score<sup>[17]</sup>, and the area under the curve (AUC) was calculated. All *P*-values were two-tailed and considered statistically significant when less than 0.05.


## RESULTS

### VGLL3 mRNA and protein levels are upregulated in GC samples


GEPIA database profiling showed that although VGLL3 was significantly overexpressed in GC samples of advanced TNM stages relative to those at the lower stages ( $P = 0.0163$ ), there was no significant difference between the GC and normal tissues ( $P > 0.05$ ). Integrated analysis of multiple VGLL3 transcriptome datasets from the ONCOMINE database, however, showed that VGLL3 was significantly overexpressed in GC ( $P = 0.003$ ), and notably associated with a poor prognosis according to the ONCOLINC database analysis ( $P = 0.01$ ,  $n = 378$ ) (Figure 1). In addition, GEPIA online survival analysis further confirmed that VGLL3 was a marker of poor prognosis in GC ( $P = 0.0062$ ,  $n = 384$ ). To validate the *in silico* data, we analyzed the expression levels of VGLL3 in the tumor and normal gastric tissues resected from 30 GC patients, and observed significantly higher levels of VGLL3 mRNA ( $P < 0.001$ ) and protein ( $P < 0.001$ ) in the GC tissues compared to normal tissues (Figure 2). Taken together, VGLL3 is upregulated in GC and possibly associated with a worse prognosis.

### VGLL3 is a potential prognostic factor for GC

The relationships between VGLL3 levels and the clinico-pathological characteristics of the patients are summarized in Table 1. High levels of VGLL3 were positively correlated with tumor lymph node metastasis ( $P = 0.028$ ) and TNM stage ( $P = 0.041$ ). In addition, the median survival time of the GC patients was significantly lower in the high VGLL3 expression group than in the low expression group (35 mo *vs* 49 mo,  $P = 0.019$ ; Figure 3C). Univariate analysis showed that the VGLL3 expression level ( $P = 0.019$ ) was the only significant prognostic factor of GC, whereas age, gender, differentiation, T stage, lymph node status, TNM stage, and HER2 status did not show any significant results ( $P > 0.05$ ). After including the significant factors in the Cox proportional hazards regression model for multivariate prognostic analysis, we found that VGLL3 expression ( $P = 0.019$ ) was an independent risk factor for GC. ROC curves were plotted to determine the power of the VGLL3 prognostic score in predicting the 3- and 5-year survival, and the AUCs were 0.613 and 0.706, respectively (Figure 3I). We constructed a PPI network of VGLL3 using the STRING database, and identified YAP1, TEAD1, WWTR1, and VGLL4 as upstream of VGLL3, and POU1F1, TEAD3, AKAP11, SYT17, CHMP28, TEAD4, and ERBBE2 (HER2) as downstream proteins (Table 2).



**Figure 1** Vestigial like family member 3 mRNA expression levels in gastric cancer tissues determined by integrated analysis of different databases. A and B: Vestigial like family member 3 (VGLL3) expression in gastric cancer and normal gastric tissues (A) and in different TNM stages (B) in the GEPIA database; C: VGLL3 expression reported in the five published studies in the ONCOMINE database; D and E: Kaplan-Meier survival analysis of patients in the GEPIA (D) and ONCOLNC (E) databases stratified by VGLL3 expression status.



**Figure 2** Vestigial like family member 3 mRNA and protein expression levels in gastric cancer. The expression of vestigial like family member 3 mRNA (A) and protein (B and C) in 30 pairs of gastric cancer and normal gastric tissues is shown.

## DISCUSSION

GC is the third most common cause of cancer-related deaths worldwide, mainly due to ineffective diagnostic and therapeutic options. It is often detected at the advanced stage, and is highly heterogeneous. Therefore, novel molecular biomarkers for the early diagnosis and treatment of GC are urgently needed. VGLL3 is a member of the vestigial-like family proteins that are related to sex and maturation<sup>[18-20]</sup>, and is reportedly associated with epithelial ovarian cancer and soft tissue sarcoma. Through bioinformatics data mining and integrated analysis, we found that VGLL3 mRNA was not only upregulated in the GC tissues relative to normal gastric tissues, but also significantly associated with advanced TNM stage GC and poor survival outcomes. To validate the *in silico* data, we analyzed the gastric tissues of GC patients, and found abnormally high levels of VGLL3 in the tumor relative to normal tissues. In addition, high *in situ* expression of VGLL3 was correlated with tumor lymph node metastasis, TNM stage, and HER2 mutation, but not with age, gender, differentiation, T stage, or M stage. The patients were stratified into the high and low VGLL3 expression groups, and the former showed a significantly poor survival. Univariate and multivariate analyses further indicated that VGLL3 expression and TNM stage were independent risk factors for the prognosis of GC. Finally, the respective AUCs of the 3- and 5-year survival of the VGLL3-based prognostic model were 0.613 and 0.706, indicating a somewhat imperfect predictive ability. This could be due to the insufficient number of samples and other confounding factors such as the patient's state of mind, economic status, and family environment. Taken together, we identified VGLL3 as a novel prognostic biomarker for GC.

To further elucidate the underlying molecular mechanisms, we analyzed the PPI network of VGLL3 through the STRING database, and found that the Hippo pathway was significantly enriched. In addition, the identified upstream molecules of VGLL3 are YAP1<sup>[6]</sup>, TEAD1, WWTR1, VGLL4, and the downstream molecules are POU1F1, TEAD3, AKAP11, SYT17, CHMP28, and TEAD4. Interestingly, the expression level of VGLL3 was related to ERBBE2 (HER2) mutation, although the relevant mechanistic connection is unclear.

To conclude, VGLL3 is highly expressed in GC and an independent risk factor, and further studies are needed to determine its underlying mechanism. Nevertheless, VGLL3 is a novel biomarker for GC prognosis and a potential therapeutic target for this malignancy.


**Table 1** Characteristics of gastric cancer patients according to vestigial like family member 3 expression status

Characteristic	Cases	High expression	Low expression	P-value
Age (yr)				
≤60	97	47	50	0.548
>60	75	32	43	
Gender				
Female	101	43	58	0.369
Male	71	36	35	
Differentiation				
Well	84	40	44	0.779
Poor and moderate	88	39	49	
T stage				
T1/2	91	36	55	0.105
T3/4	81	43	38	
N stage				
N0/1	95	36	59	0.028
N2/3	77	43	34	
TNM stage				
I-II	119	48	71	0.041
III-IV	53	31	22	
HER2				
Negative	89	32	57	0.010
Positive	83	47	36	

**Table 2** Univariate and multivariate analyses of factors associated with overall survival

Characteristic	Univariate analysis			Multivariate analysis		
	Hazard ratio	95%CI	P-value	Hazard ratio	95%CI	P-value
Age (yr)	0.83	0.55-1.26	0.389			
Gender (Male vs Female)	1.31	0.87-1.96	0.194			
Differentiation (Well vs Poor and moderate)	0.81	0.54-1.22	0.304			
T stage (T3/4 vs T1/2)	1.16	0.77-1.73	0.484			
N stage (N2/3 vs N0/1)	0.94	0.63-1.42	0.778			
TNM stage (III/IV vs I/II)	1.27	0.82-1.97	0.283			
HER2 (Positive vs Negative)	1.19	0.79-1.8	0.392			
VGLL3 (High vs Low)	1.66	1.09-2.55	0.019	1.66	1.09-2.55	0.019

VGLL3: Vestigial like family member 3.



**Figure 3 Prognostic value and putative molecular mechanism of vestigial like family member 3 in gastric cancer.** A-D: HER2 mutation status detected by FISH (A and C) and IHC (B and D) in gastric cancer (GC) tissues; E and F: *In situ* vestigial like family member 3 (VGLL3) status in GC tissues; G: Kaplan-Meier survival analysis on patients stratified by VGLL3 expression status; H: Prognostic nomogram for gastric cancer based on VGLL3 expression status and clinical features; I: Time-dependent receiver operating characteristic curves for the combination of VGLL3-based prognostic score and clinico-pathological variables; J: Protein-protein interaction network of VGLL3 based on STRING database.

## ARTICLE HIGHLIGHTS

### Research background

Gastric cancer (GC) is the most prevalent gastrointestinal tract malignancy. The prognosis of GC patients remains relatively poor. It is urgent to explore prognostic markers for GC.

### Research motivation

There are insufficient reports about the correlation between VGLL3 and GC.

### Research objectives

The aim of the present study was to explore the expression pattern and clinical significance of VGLL3 in GC.

### Research methods

It was found that VGLL3 would be a potential prognostic marker by bioinformatics analysis. To validate the *in silico* data, the authors identified the expression of VGLL3 in GC patient samples by immunohistochemistry and evaluated clinical outcomes.

### Research results

Analysis of the ONCOMINE and GEPIA databases showed that VGLL3 was significantly up-regulated in GC tissues, and associated with the tumor TNM stage. In addition, the high VGLL3 expression group had a significantly worse prognosis compared to the low expression group, as per both GEPIA and ONCOLNC. The bioinformatics results were validated by the significantly higher VGLL3 mRNA and protein levels in the GC tissues compared to the adjacent normal tissues in a cohort of 30 GC patients. Furthermore, high *in situ* expression of VGLL3 protein was associated with more advanced N and TNM stages and HER2 mutation in a cohort of 172 patients. Kaplan-Meier analysis showed that the high VGLL3 expression group had a worse prognosis compared to the low VGLL3 expression group. Multivariate analysis showed that VGLL3 expression status was an independent risk factor for prognosis. In addition, the prognostic risk model nomogram showed that VGLL3 was the most important indicator, with an AUC of 0.613 for 3-year survival and 0.706 for 5-year survival. Finally, the protein interaction network analysis revealed that VGLL3 is likely involved in the Hippo signaling pathway.

### Research conclusions

VGLL3 is overexpressed in GC tissues and associated with a poor prognosis, indicating its potential as a novel prognosis biomarker and therapeutic target for GC.

### Research perspectives

The present study suggested that VGLL3 is a novel prognostic biomarker for GC, and the significance of VGLL3 as a promising therapeutic target for GC is highlighted.

## REFERENCES

- 1 **The Lancet.** GLOBOCAN 2018: counting the toll of cancer. *Lancet* 2018; **392**: 985 [PMID: 30264708 DOI: 10.1016/S0140-6736(18)32252-9]
- 2 **Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A.** Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018; **68**: 394-424 [PMID: 30207593 DOI: 10.3322/caac.21492]
- 3 **Siegel RL, Miller KD, Jemal A.** Cancer statistics, 2019. *CA Cancer J Clin* 2019; **69**: 7-34 [PMID: 30620402 DOI: 10.3322/caac.21551]
- 4 **Gambaro K, Quinn MC, Wojnarowicz PM, Arcand SL, de Laurantaye M, Barrès V, Ripeau JS, Killary AM, Davis EC, Lavoie J, Provencher DM, Mes-Masson AM, Chevrette M, Tonin PN.** VGLL3 expression is associated with a tumor suppressor phenotype in epithelial ovarian cancer. *Mol Oncol* 2013; **7**: 513-530 [PMID: 23415753 DOI: 10.1016/j.molonc.2012.12.006]
- 5 **Cody NA, Shen Z, Ripeau JS, Provencher DM, Mes-Masson AM, Chevrette M, Tonin PN.** Characterization of the 3p12.3-pcen region associated with tumor suppression in a novel ovarian cancer cell line model genetically modified by chromosome 3 fragment transfer. *Mol Carcinog* 2009; **48**: 1077-1092 [PMID: 19347865 DOI: 10.1002/mc.20535]
- 6 **Hélias-Rodzewicz Z, Pérot G, Chibon F, Ferreira C, Lagarde P, Terrier P, Coindre JM, Aurias A.** YAP1 and VGLL3, encoding two cofactors of TEAD transcription factors, are amplified and overexpressed in a subset of soft tissue sarcomas. *Genes Chromosomes Cancer* 2010; **49**: 1161-1171 [PMID: 20842732 DOI: 10.1002/gcc.20825]
- 7 **Sun X, Wang T, Zhang C, Ning K, Guan ZR, Chen SX, Hong TT, Hua D.** S100A16 is a prognostic marker for colorectal cancer. *J Surg Oncol* 2018; **117**: 275-283 [PMID: 28876468 DOI: 10.1002/jso.24822]
- 8 **Wang LL, Chen ZS, Zhou WD, Shu J, Wang XH, Jin R, Zhuang LL, Hoda MA, Zhang H, Zhou GP.** Down-regulated GATA-1 up-regulates interferon regulatory factor 3 in lung adenocarcinoma. *Sci Rep* 2017; **7**: 2551 [PMID: 28566697 DOI: 10.1038/s41598-017-02700-5]
- 9 **Yang S, Kim CY, Hwang S, Kim E, Kim H, Shim H, Lee I.** COEXPEDIA: exploring biomedical hypotheses via co-expressions associated with medical subject headings (MeSH). *Nucleic Acids Res* 2017; **45**: D389-D396 [PMID: 27679477 DOI: 10.1093/nar/gkx868]
- 10 **Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, Simonovic M, Doncheva NT, Morris JH, Bork P, Jensen LJ, Mering CV.** STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res* 2019; **47**: D607-D613 [PMID: 30476243 DOI: 10.1093/nar/gky1131]
- 11 **Petrov A, Beer M, Blome S.** Development and validation of a harmonized TaqMan-based triplex real-time RT-PCR protocol for the quantitative detection of normalized gene expression profiles of seven porcine cytokines. *PLoS One* 2014; **9**: e108910 [PMID: 25268123 DOI: 10.1371/journal.pone.0108910]
- 12 **Sun X, Wang T, Guan ZR, Zhang C, Chen Y, Jin J, Hua D.** FBXO2, a novel marker for metastasis in human gastric cancer. *Biochem Biophys Res Commun* 2018; **495**: 2158-2164 [PMID: 29269301 DOI: 10.1016/j.bbrc.2017.12.097]
- 13 **Cai X, Zheng Y, Speck NA.** A Western Blotting Protocol for Small Numbers of Hematopoietic Stem Cells. *J Vis Exp* 2018 [PMID: 30199018 DOI: 10.3791/56855]
- 14 **Wester K, Wahlund E, Sundström C, Ranefall P, Bengtsson E, Russell PJ, Ow KT, Malmström PU, Busch C.** Paraffin section storage and immunohistochemistry. Effects of time, temperature, fixation, and retrieval protocol with emphasis on p53 protein and MIB1 antigen. *Appl Immunohistochem Mol Morphol* 2000; **8**: 61-70 [PMID: 10937051]
- 15 **Wu J, Wang F, Liu X, Zhang T, Liu F, Ge X, Mao Y, Hua D.** Correlation of IDH1 and B7H3 expression with prognosis of CRC patients. *Eur J Surg Oncol* 2018; **44**: 1254-1260 [PMID: 29871819 DOI: 10.1016/j.ejso.2018.05.005]
- 16 **Wang Y, Li J, Xia Y, Gong R, Wang K, Yan Z, Wan X, Liu G, Wu D, Shi L, Lau W, Wu M, Shen F.** Prognostic nomogram for intrahepatic cholangiocarcinoma after partial hepatectomy. *J Clin Oncol* 2013; **31**: 1188-1195 [PMID: 23358969 DOI: 10.1200/JCO.2012.41.5984]
- 17 **Kuang M, Zheng D, Tao X, Peng Y, Pan Y, Zheng S, Zhang Y, Li H, Yuan C, Zhang Y, Xiang J, Li Y, Chen H, Sun Y.** tRNA-based prognostic score in predicting survival outcomes of lung adenocarcinomas. *Int J Cancer* 2019 [PMID: 30838640 DOI: 10.1002/ijc.32250]

- 18 **Barson NJ**, Aykanat T, Hindar K, Baranski M, Bolstad GH, Fiske P, Jacq C, Jensen AJ, Johnston SE, Karlsson S, Kent M, Moen T, Niemelä E, Nome T, Næsje TF, Orell P, Romakkaniemi A, Sægrov H, Urdal K, Erkinaro J, Lien S, Primmer CR. Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature* 2015; **528**: 405-408 [PMID: 26536110 DOI: 10.1038/nature16062]
- 19 **Tu W**, Wagner EK, Eckert GJ, Yu Z, Hannon T, Pratt JH, He C. Associations between menarche-related genetic variants and pubertal growth in male and female adolescents. *J Adolesc Health* 2015; **56**: 66-72 [PMID: 25287989 DOI: 10.1016/j.jadohealth.2014.07.020]
- 20 **Liang Y**, Tsoi LC, Xing X, Beamer MA, Swindell WR, Sarkar MK, Berthier CC, Stuart PE, Harms PW, Nair RP, Elder JT, Voorhees JJ, Kahlenberg JM, Gudjonsson JE. A gene network regulated by the transcription factor VGLL3 as a promoter of sex-biased autoimmune diseases. *Nat Immunol* 2017; **18**: 152-160 [PMID: 27992404 DOI: 10.1038/ni.3643]



Published By Baishideng Publishing Group Inc  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

