

World Journal of *Gastroenterology*

World J Gastroenterol 2019 June 14; 25(22): 2699-2832



**OPINION REVIEW**

- 2699** Which factors determine exocrine pancreatic dysfunction in diabetes mellitus?
Altay M

REVIEW

- 2706** Proton pump inhibitors and dysbiosis: Current knowledge and aspects to be clarified
Bruno G, Zaccari P, Rocco G, Scalese G, Panetta C, Porowska B, Pontone S, Severi C
- 2720** Diagnosis and therapeutic strategies for small bowel vascular lesions
Sakai E, Ohata K, Nakajima A, Matsuhashi N

MINIREVIEWS

- 2734** Advanced diagnostics for pancreatic cysts: Confocal endomicroscopy and molecular analysis
Durkin C, Krishna SG
- 2743** Long-lasting discussion: Adverse effects of intraoperative blood loss and allogeneic transfusion on prognosis of patients with gastric cancer
Nakanishi K, Kanda M, Kodera Y

ORIGINAL ARTICLE**Basic Study**

- 2752** MiR-34a overexpression enhances the inhibitory effect of doxorubicin on HepG2 cells
Zheng SZ, Sun P, Wang JP, Liu Y, Gong W, Liu J
- 2763** Long noncoding RNA HOXA11-AS promotes gastric cancer cell proliferation and invasion *via* SRSF1 and functions as a biomarker in gastric cancer
Liu Y, Zhang YM, Ma FB, Pan SR, Liu BZ
- 2776** Sp1 contributes to overexpression of stanniocalcin 2 through regulation of promoter activity in colon adenocarcinoma
Li JB, Liu ZX, Zhang R, Ma SP, Lin T, Li YX, Yang SH, Zhang WC, Wang YP

Retrospective Study

- 2788** Increased risk of atrial fibrillation in patients with inflammatory bowel disease: A nationwide population-based study
Choi YJ, Choi EK, Han KD, Park J, Moon I, Lee E, Choe WS, Lee SR, Cha MJ, Lim WH, Oh S

Clinical Trials Study

- 2799** Effects of early enteral nutrition on Th17/Treg cells and IL-23/IL-17 in septic patients
Sun JK, Zhang WH, Chen WX, Wang X, Mu XW

SYSTEMATIC REVIEWS

- 2809** Autoimmune hepatitis treatment in the elderly: A systematic review
Durazzo M, Lupi G, Scandella M, Ferro A, Gruden G

META-ANALYSIS

- 2819** Mini-invasive *vs* open resection of colorectal cancer and liver metastases: A meta-analysis
Ye SP, Qiu H, Liao SJ, Ai JH, Shi J

ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Andrew Stewart Day, MD, Professor, Paediatrics Department, University of Otago, Christchurch 8041, New Zealand

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. The *WJG* Editorial Board consists of 642 experts in gastroenterology and hepatology from 59 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, etc. The *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

The *WJG* is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, Scopus and Directory of Open Access Journals. The 2018 edition of Journal Citation Report® cites the 2017 impact factor for *WJG* as 3.300 (5-year impact factor: 3.387), ranking *WJG* as 35th among 80 journals in gastroenterology and hepatology (quartile in category Q2).

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: Yan-Liang Zhang

Proofing Production Department Director: Yun-Xiaojian Wu

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Subrata Ghosh, Andrzej S Tarnawski

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE

Ze-Mao Gong, Director

PUBLICATION DATE

June 14, 2019

COPYRIGHT

© 2019 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Basic Study

Long noncoding RNA HOXA11-AS promotes gastric cancer cell proliferation and invasion *via* SRSF1 and functions as a biomarker in gastric cancer

Yun Liu, Yu-Mei Zhang, Feng-Bo Ma, Su-Rong Pan, Bao-Zhen Liu

ORCID number: Yun Liu (0000-0003-1622-0078); Yu-Mei Zhang (0000-0003-2087-4816); Feng-Bo Ma (0000-0002-2275-073X); Su-Rong Pan (0000-0002-4891-1941); Bao-Zhen Liu (0000-0002-2453-4643).

Author contributions: Liu BZ designed the research; Liu Y, Zhang YM, Ma FB, and Pan SR performed the research; Liu Y analyzed the data; Liu Y and Liu BZ wrote the paper.

Institutional review board statement: This study was reviewed and approved by the Institutional Review Board Committee of Binzhou People's Hospital.

Conflict-of-interest statement: We declare no conflict of interest.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Yun Liu, Department of Operating Room, Binzhou People's Hospital, Binzhou 256610, Shandong Province, China

Yu-Mei Zhang, Department of Return Visit, Binzhou People's Hospital, Binzhou 256610, Shandong Province, China

Feng-Bo Ma, Su-Rong Pan, Bao-Zhen Liu, Department of Gastroenterology, Binzhou People's Hospital, Binzhou 256610, Shandong Province, China

Corresponding author: Bao-Zhen Liu, PhD, Doctor, Department of Gastroenterology, Binzhou People's Hospital, Binzhou 256610, Shandong Province, China. zhebaonnt3411@163.com
Telephone: +86-543-3282396

Abstract

BACKGROUND

Gastric cancer (GC) is the fourth most frequent malignancy all over the world. The diagnosis of GC is challenging and the prognosis of GC is very unfavorable. Accumulating evidence reveals that serum long noncoding RNAs (lncRNAs) can function as biomarkers in various types of cancers, including GC.

AIM

To explore the level and molecular mechanism of the lncRNA HOXA11-AS in GC and the diagnostic and prognostic significance of serum HOXA11-AS in GC.

METHODS

HOXA11-AS levels in GC tissue, cell lines, and serum samples were measured. The correlation between HOXA11-AS expression and clinicopathological characteristics was analyzed. The role of HOXA11-AS in the diagnosis and prognosis of GC was evaluated. Cell function assays were performed for exploration of the roles of HOXA11-AS in GC cells. Moreover, Western blot was performed to explore the target regulated by HOXA11-AS in GC cells.

RESULTS

Up-regulation of HOXA11-AS was found in GC tissues, cell lines, and serum samples. In GC patients, decreased serum HOXA11-AS levels were negatively related with tumor size, TNM stage, and lymph node metastasis. The area under the receiver operating characteristic curve of serum HOXA11-AS in the diagnosis of GC was 0.924 (95%CI: 0.881-0.967; sensitivity, 0.787; specificity 0.978). Results of the Kaplan-Meier survival curves suggested the GC patients with a lower

Manuscript source: Unsolicited manuscript

Received: March 28, 2019

Peer-review started: March 28, 2019

First decision: April 11, 2019

Revised: April 15, 2019

Accepted: May 3, 2019

Article in press: May 3, 2019

Published online: June 14, 2019

P-Reviewer: Lazăr DC, Tanabe S

S-Editor: Gong ZM

L-Editor: Wang TQ

E-Editor: Zhang YL



HOXA11-AS level having a better overall survival rate. HOXA11-AS promoted GC cell proliferation and invasion. SRSF1 may be the target regulated by HOXA11-AS in GC cells.

CONCLUSION

HOXA11-AS promotes GC cell proliferation and invasion *via* SRSF1 and may function as a promising marker in GC.

Key words: Long noncoding RNA; HOXA11-AS; SRSF1; Gastric cancer; Biomarker

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: The long noncoding RNA (lncRNA) HOXA11-AS is up-regulated in gastric cancer (GC) tissues, cell lines, and serum samples. In GC patients, increased serum HOXA11-AS levels were positively related with tumor size, TNM stage, and lymph node metastasis. HOXA11-AS functions as a diagnostic and prognostic biomarker in GC, and promotes GC cell proliferation and invasion. SRSF1 may be the target regulated by HOXA11-AS in gastric cells.

Citation: Liu Y, Zhang YM, Ma FB, Pan SR, Liu BZ. Long noncoding RNA HOXA11-AS promotes gastric cancer cell proliferation and invasion *via* SRSF1 and functions as a biomarker in gastric cancer. *World J Gastroenterol* 2019; 25(22): 2763-2775

URL: <https://www.wjgnet.com/1007-9327/full/v25/i22/2763.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v25.i22.2763>

INTRODUCTION

Gastric cancer (GC) is a global health problem, holding the fourth most frequent malignancy and the second most common cause of death from cancer all over the world^[1]. Estimated 27510 cases diagnosed with GC and 11140 deaths from GC will occur in the United States in 2019^[2]. The diagnosis of GC is challenging owing to the fact that patients often present with vague and non-specific symptoms^[3], which causes many patients being at the progressive stage at first diagnosis. However, patients with metastatic GC carry a very poor prognosis and their median survival ranges from 4 mo to around 12 mo^[4,5]. Although GC is associated with *Helicobacter pylori* infection, the molecular mechanisms of its occurrence and progression remain unclear. Consequently, it is ineluctably necessary to probe into the molecular mechanism of the occurrence as well as progression of GC and finding a new marker applying to the detection, therapy, and prognosis of GC.

Long noncoding RNAs (lncRNAs), a class of transcripts > 200 nucleotides (nts) in length, exert their significant functions in the progression and metastasis of malignancy^[6]. lncRNAs typically exhibit tissue-specific expression patterns and are readily detectable in body fluids because of their high stability, in comparison with other protein biomarkers expressed in various types of tissues, making them ideal biomarkers^[7]. Accumulating evidence reveals that serum lncRNAs function as biomarkers in various types of cancers, such as HOTTIP in GC^[8], MALAT1 in epithelial ovarian cancer^[9], GIHCG in cervical cancer^[10], and LRB1 in hepatocellular carcinoma^[11].

Previous studies suggested that aberrantly expressed lncRNA HOXA11-AS plays significant roles in the development and progression of malignancies^[12]. HOXA11-AS functions as an oncogene and promotes cell proliferation, invasion, and metastasis in GC^[13,14]. However, the molecular mechanism of HOXA11-AS in GC is far from fully elucidated and the diagnostic and prognostic role of HOXA11-AS in GC is still unclear. In this study, we thoroughly investigated the molecular mechanism of HOXA11-AS and the diagnostic and prognostic roles of serum HOXA11-AS in GC.

MATERIALS AND METHODS

Tissue and serum specimens

GC tissue specimens and corresponding paracancerous gastric tissue specimens were

obtained from 25 GC patients from Binzhou People's Hospital. Each patient was diagnosed with GC pathologically and none of the patients received any treatment before operation. The tissue samples were processed within 1 hour and submerged in RNAlater reagent from Qiagen GmbH (Hilden, Germany) for half an hour. After that, the samples were stored at -80 °C till RNA extraction. All tissue samples were examined and classified under the management of experienced pathologists.

As for blood samples, the present study totaled 134 participants, consisting of 94 patients with GC and 40 healthy controls. Postoperative blood specimens were also obtained from 25 patients with GC. All the blood samples were disposed within 2 h. Serum was disposed at 4 °C and recruited by centrifugation ($1200 \times g$, 10 min). Another centrifugation ($10000 \times g$, 10 mins, at 4 °C) was performed for completely removing residual cellular debris. The serum specimens were kept in liquid nitrogen till RNA extraction.

Our research was managed by the ethics committee of Binzhou People's Hospital. Consent statement was obtained from each participant. Table 1 lists the clinical characteristics of the GC patients.

Cell culture and transfection

A culture medium consisting of 88% of Dulbecco's modified Eagle's medium (DMEM; Gibco; Thermo Fisher Scientific, Inc., Waltham, MA, United States), 10% of fetal bovine serum (FBS; Gibco; Thermo Fisher Scientific, Inc.), 1% of antibiotics (100 μ L/mL penicillin and 100 mg/mL streptomycin sulfate), and 1% of glutamine was used to cultivate normal gastric mucosal epithelial cells (GES-1) and GC cells (MKN-45). MKN-45 and GES-1 cells were maintained in an incubator (37 °C, 5% CO₂). MKN-45 cells were pre-seeded in the 6-well plates (3.0×10^5 cells/per well) and maintained until the confluence of 60%. HOXA11-AS primers, siRNA, and negative control (si-NC) were provided by Ribobio (Guangzhou, China); pcDNA-HOXA11-AS and pcDNA3.1 were constructed by Generay (Shanghai, China). siRNAs or pcDNAs were transfected into GES-1 cells with Lipofectamine RNAiMAX Reagent (Life Technologies, Carlsbad, CA, United States).

RNA isolation

The extraction of total RNA from tissue and cell samples was done using TRIzol reagent from Invitrogen (Thermo Fisher Scientific) following the manufacturer's procedures. The isolation of total RNA from serum samples was done with Qiagen miRNeasy Serum/Plasma Kit from Qiagen GmbH (Hilden, Germany) following the manufacturers' procedures. The quantity and purity of the extracted RNA samples were detected with NanoDrop 2000c (Thermo Fisher Scientific, Inc.). The RNA samples with an optical density ratio (260/280) of 1.8-2.0 were accepted into later study. RNA samples were either kept at -80 °C or used for direct synthesis of cDNA.

Reverse transcription and quantitative real-time polymerase chain reaction (qRT-PCR)

The PrimerScript RT Master from Takara Biotechnology (Dalian, China) was used for cDNA synthesis. The level of HOXA11-AS was measured by RT-qPCR with SYBR-Green PCR Master mix (Roche, Mannheim, Germany) on the Roche Lightcycler 480 Real-Time PCR system (Roche Diagnostics, Basel, Switzerland). The qRT-PCR reactions were performed in triplicate. Changes in HOXA11-AS level were calculated using the $2^{-\Delta\Delta C_t}$ method. Gene expression values were normalized to the expression of GAPDH. The sequences used in qRT-PCR are as follows: HOXA11-AS-F: 5'-GATTTCTCCAGCCTCCCTTC-3' and HOXA11-AS-R: 5'-AGAAATTG GACGAGACTGCG-3'; GAPDH-F: 5'-TGTTATTCGTGGAAGGACTCAT-3' and GAPDH-R: 5'-GTGGGTGTCGCTGTTGAAGTC-3'.

Cell counting kit 8 (CCK-8) assay

CCK-8 assay was conducted to explore cell proliferation ability. In this assay, a 96-well plate was used and transfected MKN-45 cells were seeded in each well at a density of 3.0×10^3 cells/well. The cells were maintained in a cell incubator. Ten microliters of CCK-8 reagent (US Everbright Inc.) were added to each well at 24, 48, and 72 h after the cells were incubated, respectively. The optical density of each well of 96-well plates was detected using an enzyme immunoassay instrument from Bio Rad Laboratories (Hercules, CA, USA).

Transwell assay

Transwell assay was conducted to explore cell invasive capability. Transfected MKN-45 cells (4×10^4) with 200 μ L of DMEM medium without serum were added into a transwell chamber (BD Biosciences, New York, NJ, United States) with Matrigel, which was inserted into a 24-well plate. Each well of the 24-well plate was added with

Table 1 Clinicopathological characteristics of the study population

	Tissue samples (n = 25)	Serum samples	Healthy controls
Gender			
Male	15	57	26
Female	10	37	14
Age			
Mean \pm SD	60 \pm 7	59 \pm 7	59 \pm 8
Median (range)	61 (45-72)	60 (40-74)	60 (41-74)
Tumor size (cm)			
<5	19	59	
\geq 5	6	35	
TNM stage			
I	13	18	
II	9	21	
III	3	42	
IV	0	13	
Differentiation			
Well and moderate	13	50	
Poor	12	44	
Invasion depth			
T1	3	9	
T2	7	32	
T3	12	35	
T4	3	18	
Lymph node metastasis			
Positive	18	50	
Negative	7	44	

450 μ L of DMEM medium and 50 μ L of FBS. MKN-45 cells were cultured for another 36 hours. The MKN-45 cells that did not invade through the membrane were washed and cleaned. Crystal violet (1%) was used and the invaded MKN-45 cells invading through the membrane were stained for 25 min. After washing cells with PBS three times, the number of MKN-45 cells invading through the membrane was calculated in three random visual places using a light microscope.

Protein extraction and Western blot analysis

After rinsing with pre-cooling PBS, MKN-45 cells were dissociated with RIPA buffer (Thermo Fisher Scientific) mixed with protease and phosphatase inhibitor cocktail (Roche). The total protein of MKN-45 cells was isolated on the basis of instructions. The Bio-Rad assay system (Bio-Rad Laboratories, Hercules, CA, United States) was used to detect the protein concentration. In Western blot analysis, 12% sodium dodecyl sulfate-polyacrylamide gels were prepared and the same amounts of protein extract were separated by electrophoresis and then transferred to polyvinylidene difluoride (PVDF) membranes (Millipore, Billerica, MA, USA). After blocking in 5% skim milk, the blotted membranes were immersed in primary antibodies (SRSF1, 1:1000; GAPDH, 1:2000, Proteintech, Wuhan, China) for 24 h. The membranes were rinsed with TBST three times (10 min/time). Subsequently, the membranes were immersed in a secondary antibody (HRP conjugated goat anti-rabbit antibody, 1:1000, Sigma) at room temperature for 1 h. GAPDH was used as the internal control. Kodak film (Kodak, Rochester, NY, United States) was used to detect the blots with an enhanced chemiluminescence kit (Merck Millipore).

Statistical analysis

SPSS 21.0 software and Graphpad prism 7 software were used for statistical analyses. The Student's *t*-test was conducted to evaluate the differences between subgroups. The Chi-square test and Fisher's exact test was conducted to explore the associations between HOXA11-AS expression and clinicopathological features. ROC curve

analysis was performed for the assessment of the possibility of serum HOXA11-AS for GC detection. The significance of serum HOXA11-AS for prognosis evaluation of GC was evaluated by the Kaplan-Meier method. Cox regression analysis was conducted for evaluating the possible factors that can predict the prognosis of GC. $P < 0.05$ was considered statistically significant.

RESULTS

HOXA11-AS levels in tissue and serum samples

We first explored the levels of HOXA11-AS in tissue and serum samples. As presented in **Table 1**, there was no significant difference in age or gender between GC patients and healthy controls. A significantly increased level of HOXA11-AS was found in GC tissues *vs* the paired normal gastric tissues (**Figure 1A**, $P < 0.001$). Furthermore, a similar result was obtained in serum samples. Significant up-regulation of HOXA11-AS was found in the serum of GC patients *vs* healthy individuals (**Figure 1B**, $P < 0.001$). As expected, the same result was obtained in GC cell lines, which revealed that HOXA11-AS was up-regulated in MKN-45 cells compared with GES-1 cells (**Figure 1C**, $P < 0.01$).

Association between HOXA11-AS level and clinicopathologic features of GC patients.

We next thoroughly investigated the association between HOXA11-AS level and GC patients' clinicopathologic features. The 94 GC patients were split into two parts using median serum HOXA11-AS l as the cutoff (47 with a high serum HOXA11-AS level and 47 with a low serum HOXA11-AS level). As shown in **Table 2**, there was no significant correlation between serum HOXA11-AS and gender, age, differentiation, or invasion depth ($P > 0.05$), while a remarkable association was obtained between serum HOXA11-AS and TNM stage ($P = 0.001$), tumor size ($P = 0.001$), as well as lymph node metastasis ($P = 0.011$). We found that the GC patients with decreased serum HOXA11-AS more probably had smaller tumor, earlier TNM stage, and negative lymph node metastasis.

HOXA11-AS functions as a potential diagnostic marker in GC

We further tested HOXA11-AS expression in serum samples from postoperative GC patients. The data demonstrated that compared with preoperative samples, serum HOXA11-A levels showed no significant difference in the samples obtained 1 d after surgery ($P > 0.05$), but they were greatly decreased in the samples obtained 1 mo after surgery ($P = 0.001$) (**Figure 2A**). Furthermore, the area under the ROC curve (AUC) of serum HOXA11-AS in the diagnosis of GC was 0.924 (95%CI: 0.881-0.967, sensitivity 0.787, specificity 0.978) (**Figure 2A**). These data demonstrated that HOXA11-AS possibly acts as a potential marker in the diagnosis of GC.

HOXA11-AS may function as a potential prognostic biomarker in GC

Subsequently, we analyzed the role of HOXA11-AS expression in the prognosis evaluation of GC patients. As shown in **Table 3**, the univariate Cox proportional hazard regression analysis revealed that tumor size (HR = 0.527, 95%CI: 0.301-0.924, $P = 0.025$), TNM stage (HR = 0.195, 95%CI: 0.092-0.417, $P = 0.001$), and serum HOXA11-AS (HR = 0.201, 95%CI: 0.104-0.835, $P = 0.001$) were potential factors affecting the overall survival of GC patients, and the multivariate Cox proportional hazard regression analysis demonstrated that serum HOXA11-AS (HR = 0.338, 95%CI: 0.115-0.989, $P = 0.048$) was a potential factor affecting the overall survival of GC patients. What's more, the Kaplan-Meier survival curve was drawn, which demonstrated that GC individuals with lower expression of HOXA11-AS had a better overall survival rate ($P = 0.001$, **Figure 3**). These data demonstrated that HOXA11-AS may act as a potential marker for the prognosis of GC.

Cell transfection efficiency

The down-regulation or up-regulation of HOXA11-AS in MKN-45 cells transfected with pcDNA HOXA11-AS or siRNA was authenticated by RT-qPCR. The amount of HOXA11-AS in MKN-45 cells transfected with pc-DNA-HOXA11-AS was noteworthy upregulated *vs* those transfected with pcDNA3.1 ($P < 0.01$, **Figure 4A**). Moreover, MKN-45 cells transfected with HOXA11-AS-siRNA1 ($P < 0.001$) or HOXA11-AS -siRNA2 ($P < 0.01$) showed a higher level of HOXA11-AS than those transfected with si-NC (**Figure 4B**).

HOXA11-AS promotes MKN-45 cell proliferation

In our study, CCK8 assay was conducted to evaluate cell proliferative capability. As




Figure 1 HOXA11-AS expression in GC tissues, cell lines, and serum samples. A: Relative expression of HOXA11-AS in gastric cancer and normal tissues; B: Relative levels of serum HOXA11-A in gastric cancer patients and healthy controls; C: Relative levels of HOXA11-A in MNK-45 and GES-1 cells. The assays were repeated three times. ^b $P < 0.01$, ^c $P < 0.001$. GC: Gastric cancer.

expected, cell proliferation was faster in MKN-45 cells transfected with pcDNA HOXA11-AS compared with those transfected with pcDNA 3.1 (Figure 4C). On the contrary, cell proliferation was slower in MKN-45 cell transfected with siRNA1 or siRNA2 in comparison with those transfected with si-NC (Figure 4D).

HOXA11-AS promotes MKN-45 cell invasion

Transwell assay was conducted to evaluate cell invasion capability, and the results are presented in Figure 4E and F, which revealed that the quantity of invaded cells was significantly higher in MKN-45 cells transfected with pcDNA HOXA11-AS than in those transfected with pcDNA 3.1 ($P < 0.01$, Figure 4E). However, the quantity of invaded cells was remarkably lower in MKN-45 cells transfected with siRNA1 ($P < 0.05$) or siRNA2 ($P < 0.05$) than in those transfected with si-NC (Figure 4F).

HOXA11-AS regulates expression of SRSF1 in MKN-45 cells

The result of Western blot indicated that the protein expression of SRSF1 in MKN-45 cells was upregulated when the expression of HOXA11-AS was increased ($P < 0.001$, Figure 5A). On the contrary, the protein expression of SRSF1 in MKN-45 cells was downregulated when the expression of HOXA11-AS was decreased ($P < 0.01$, Figure 5B).

DISCUSSION

GC is one of the most frequent and fatal malignancies, seriously threatening human health. Because of the position of the stomach and atypical clinical features of GC, it is quite difficult to identify GC patients from the healthy control at an early curable stage. Therefore, over half of GC patients are already in the advanced stage when they are initially diagnosed with GC and the 5-year overall survival (OS) rate of GC is less than 30%^[15]. However, there are no biomarkers that are accurate and easy to detect for the diagnosis, prognosis evaluation, and treatment of GC clinically, which may be attributed to the mystery of the molecular mechanism about the genesis and progression of GC.

HOXA11-AS, located at the 5' region of the HOXA cluster, is one of frequently studied lncRNAs^[12]. The level of HOXA11-AS was significantly increased in lung cancer^[16], breast cancer^[17], renal cancer^[18], and osteosarcoma^[19]. Previous studies have found that HOXA11-AS served as a biomarker in head and head-neck carcinoma^[20] and esophageal carcinoma^[21]. Moreover, a meta-analysis revealed that HOXA11-AS functioned as a prognostic marker in human solid tumors^[22]. As for non-small cell lung cancer (NSCLC), an increased level of HOXA11-AS had a promotive role in cell epithelial-mesenchymal transition, which was achieved through suppressing miR-200b^[23]. Moreover, the levels of HOXA11-AS in hepatocellular carcinoma tissues and cell lines were upregulated. HOXA11-AS promoted cell proliferation by regulating cell cycle and apoptosis *via* DUSP5 in hepatocellular carcinoma^[24]. These results demonstrated that HOXA11-AS may play a significant role in various types of cancers, including GC. In GC, HOXA11-AS was reported to be act as an oncogene and promote GC cell proliferation, invasion, and metastasis^[13,14]. However, the roles of HOXA11-AS in the diagnosis and prognosis of GC have yet to be fully elucidated.

In our study, the levels of HOXA11-AS in GC tissue samples, cell lines, and serum

Table 2 Association of serum HOXA11-AS with clinicopathological characteristics in patients with gastric cancer

Clinical feature	Cases	Serum HOXA11-AS	P-value
Gender			0.824
Male	57	0.345 ± 0.104	
Female	37	0.349 ± 0.091	
Age (yr)			0.304
<60	38	0.337 ± 0.111	
≥60	56	0.355 ± 0.089	
Tumor size (cm)			0.001
<5	59	0.320 ± 0.094	
≥5	35	0.391 ± 0.092	
TNM stage			0.001
I and II	39	0.250 ± 0.073	
III and IV	55	0.415 ± 0.040	
Differentiation			0.934
Well and moderate	50	0.346 ± 0.112	
Poor	44	0.347 ± 0.084	
Invasion depth			0.867
T1 and T2	41	0.348 ± 0.100	
T3 and T4	53	0.345 ± 0.099	
Lymph node metastasis			0.011
Positive	50	0.377 ± 0.088	
Negative	44	0.312 ± 0.100	

samples were detected, which revealed increased levels of HOXA11-AS in these samples and cell lines. Moreover, in GC patients, serum HOXA11-AS was negatively related to tumor size, TNM stage, and lymph node metastasis. The role of HOXA11-AS in the diagnosis of GC was explored, which revealed that the AUC of serum HOXA11-AS in the diagnosis of GC was 0.924 (95%CI: 0.881-0.967, sensitivity 0.787, specificity 0.978). The Kaplan-Meier survival curve was drawn to evaluate HOXA11-AS in the prognosis of GC and the data revealed that the GC patients with low HOXA11-AS expression has a better overall survival rate. Further functional assays were conducted to probe into how HOXA11-AS affects the biological behavior of GC cells, which revealed that HOXA11-AS promoted GC cell proliferation and invasion.

Serine/arginine splicing factor 1 (SRSF1), an important family member of SR proteins, was reported to participate in multiple biological functions, including splicing regulation, translation control, RNA transport, cell proliferation, invasion, and senescence^[25-29]. SRSF1 has emerged as a key oncogene in numerous solid tumors, such as GC. In our research, SRSF1 was found to be positively co-expressed with HOXA11-AS in GC by StarBase 3.0 (<http://starbase.sysu.edu.cn/index.php>). Therefore, we assume that SRSF1 may be a target of HOXA11-AS in GC cells. Western blot analysis was performed, and as expected, the results showed that SRSF1 may be the target of HOXA11-AS in GC cells.

LncRNAs play a significant role in the genesis and progression of malignant tumors. Previous research found that downregulation of HOXA11-AS caused positive changes in NSCLC cell proliferation and migration, reversed the epithelial mesenchymal transition process, and induced cell apoptosis^[30]. HOXA11-AS was increased in breast cancer and promoted cell invasion and metastasis *via* EMT, providing a theoretical basis and important molecular target for the therapy of breast cancer^[17]. Results of previous research verified that HOXA11-AS promoted liver metastasis in colorectal cancer by acting as a miR-125a-5p sponge^[31]. HOXA11-AS also exerted its functions in glioblastoma, uveal melanoma, renal cancer, and cervical cancer^[18,32-34].

Previous studies revealed that lncRNAs can function as diagnostic biomarkers in GC. It was found that serum lncRNA CTC-497E21.4 may function as a potentially diagnostic marker for GC^[35]. A lower level of circulating LINC00086 expression was identified in GC patients than in normal individuals. LINC00086 distinguished GC patients from healthy controls with a high sensitivity and specificity, suggested that LINC00086 is a potential biomarker for the diagnosis of GC^[36]. Moreover, circulating




Figure 2 Performance of HOXA11-AS in the diagnosis of gastric cancer. A: Serum HOXA11-AS levels in the samples obtained 1 d and 1 mo after surgery; B: The ROC curve demonstrates that serum HOXA11-AS can discriminate gastric cancer patients from healthy controls (AUC = 0.924, 95%CI: 0.881-0.967, $P = 0.001$) with a sensitivity of 78.7% and specificity of 97.8%. $^cP < 0.001$. The assays were repeated three times. PreOp: Pre-operation; 1 d: The first day after surgical removal of the tumor; 1 mo: The 30th day after surgical removal of the tumor; ROC: Receiver operating characteristic curve; AUC: Area under the curve.


H19 was upregulated in GC patients in comparison with healthy individuals and H19 expression declined significantly upon surgical removal of the tumors, suggesting that H19 can act as a diagnostic biomarker of GC^[37].

Previous studies also revealed that lncRNAs can function as prognostic biomarkers in GC. It was found that serum lncRNA HULC was upregulated in GC patients and Kaplan-Meier curve analysis suggested that HULC was a good predictor of the prognosis of GC^[7]. Circulating GACAT2 level was noteworthy up-regulated in preoperative GC patients than in the postoperative group. Moreover, the expression of GACAT2 were associated with the clinicopathologic features (lymphatic metastasis, distal metastasis, and perineural invasion) of patients with GC, which suggested that GACAT2 may be a potential tumor marker for the prediction of GC patient prognosis^[16]. It was reported that up-regulation of plasma MALAT1 was independently correlated with a poor prognosis in GC, demonstrating MALAT1 as a novel marker for the distant metastasis of GC patients^[38].

In conclusion, serum HOXA11-AS significantly increases in GC patients. HOXA11-AS promotes GC cell proliferation and invasion *via* SRSF1 and may function as a diagnostic and prognostic biomarker in GC.

Table 3 Univariate and multivariate survival analyses of 94 gastric cancer patients

Variable	Univariate analysis			Multivariate analysis		
	HR	95%CI	P-value	HR	95%CI	P-value
Gender (Male <i>vs</i> Female)	1.310	0.749-2.291	0.344	---	---	---
Age (yr) (<60 <i>vs</i> ≥60)	0.987	0.560-1.739	0.963	---	---	---
Tumor size (cm) (<5 <i>vs</i> ≥5)	0.527	0.301-0.924	0.025	0.816	0.442-1.508	0.516
TNM stage (I + II <i>vs</i> III + IV)	0.195	0.092-0.417	0.001	0.437	0.125-1.522	0.193
Differentiation (Well and moderate <i>vs</i> Poor)	0.923	0.529-1.61	0.779	---	---	---
Invasion depth (T1 + T2 <i>vs</i> T3 <i>vs</i> T4)	1.050	0.601-1.836	0.863	---	---	---
Lymph node metastasis (Positive <i>vs</i> Negative)	0.667	0.379-1.176	0.162	---	---	---
Serum HOXA11-AS (High <i>vs</i> Low)	0.201	0.104-0.385	0.001	0.338	0.115-0.989	0.048

**Figure 3** Kaplan-Meier survival curves demonstrating that gastric cancer patients with low HOXA11-AS expression has a better overall survival rate. The assays were repeated three times.




Figure 4 Transfection efficiencies and cell function assays. A and B: Transfection efficiencies of pcDNA-HOXA11-AS (A) and HOXA11-AS-siRNA (B) in gastric cancer cells; C and D: Migration of MKN-45 cells after HOXA11-AS overexpression (C) or knockdown (D); E and F: Invasion of MKN-45 cells after HOXA11-AS overexpression (E) or knockdown (F). The assays were repeated three times. ^a $P < 0.05$, ^b $P < 0.01$, ^c $P < 0.001$. OD: Optical density; NC: Normal control.




Figure 5 HOXA11-AS regulates expression of SRSF1 in gastric cancer cells. A: Protein expression of SRSF1 after overexpression of HOXA11-AS in gastric cancer (GC) cells. B: Protein expression of SRSF1 after knockdown of HOXA11-AS in GC cells. The assays were repeated three times. * $P < 0.05$, ^c $P < 0.001$. NC: Normal control.

ARTICLE HIGHLIGHTS

Research background

As one of the most frequent cancers, gastric cancer (GC) caused more than 700000 deaths in just 2012 worldwide. Many recent studies have demonstrated the molecular mechanisms involved in transcriptional regulation in GC, and long noncoding RNAs (lncRNAs) play an irreplaceable role in the initiation and progression of GC, such as maintaining cell growth, evasion of apoptosis, promotion of invasion and metastasis, stemness maintenance, and EMT.

Research motivation

To identify more biomarkers for the diagnosis and treatment of GC.

Research objectives

This study aimed to investigate the underlying mechanisms of HOXA11-AS in GC.

Research methods

HOXA11-AS expression was detected by qRT-PCR assay in GC tissues, cell lines, and serum samples. Clinicopathological characteristics were collected and expression analysis of HOXA11-AS was performed to evaluate the role of HOXA11-AS. Cell function assays were performed to explore the functions of HOXA11-AS in GC cell lines. Moreover, Western blot was performed to explore the target regulated by HOXA11-AS in GC cell lines.

Research results

We found that HOXA11-AS was upregulated in GC tissues, cell lines, and serum samples, and exhibited a significant negative correlation with tumor size, TNM stage, and lymph node metastasis. Cell experiments showed that HOXA11-AS promoted the proliferation and invasion capacity of GC cell lines, and SRSF1 may be the target regulated by HOXA11-AS in GC cells. Especially, GC patients with a lower HOXA11-AS level had a better overall survival rate.

Research conclusions

Our study demonstrated that HOXA11-AS can significantly promote GC cell growth, migration, and invasion. Furthermore, it can work through SRSF1. Therefore, our study provides the possible molecular mechanism and two new biomarkers for GC.

Research perspectives

In the future, research may reveal the important role of HOXA11-AS that enhances the sensitivity of GC detection and facilitate its application in anti-cancer treatments. The identification of the HOXA11-AS/SRSF1 molecular axis may further explain the underlying mechanism.

REFERENCES

- 1 **Van Cutsem E**, Sagaert X, Topal B, Haustermans K, Prenen H. Gastric cancer. *Lancet* 2016; **388**: 2654-2664 [PMID: 27156933 DOI: 10.1016/S0140-6736(16)30354-3]
- 2 **Siegel RL**, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin* 2019; **69**: 7-34 [PMID: 30620402 DOI: 10.3322/caac.21551]
- 3 **Thrumurthy SG**, Chaudry MA, Hochhauser D, Mughal M. The diagnosis and management of gastric cancer. *BMJ* 2013; **347**: f6367 [PMID: 24191271 DOI: 10.1136/bmj.f6367]
- 4 **Murad AM**, Santiago FF, Petroianu A, Rocha PR, Rodrigues MA, Rausch M. Modified therapy with 5-fluorouracil, doxorubicin, and methotrexate in advanced gastric cancer. *Cancer* 1993; **72**: 37-41 [PMID: 8508427 DOI: 10.1002/1097-0142(19930701)72:1<37::AID-CNCR2820720109>3.0.CO;2-P]
- 5 **Wagner AD**, Unverzagt S, Grothe W, Kleber G, Grothey A, Haerting J, Fleig WE. Chemotherapy for advanced gastric cancer. *Cochrane Database Syst Rev* 2010; CD004064 [PMID: 20238327 DOI: 10.1002/14651858.CD004064.pub3]
- 6 **Lee YR**, Kim G, Tak WY, Jang SY, Kweon YO, Park JG, Lee HW, Han YS, Chun JM, Park SY, Hur K. Circulating exosomal noncoding RNAs as prognostic biomarkers in human hepatocellular carcinoma. *Int J Cancer* 2019; **144**: 1444-1452 [PMID: 30338850 DOI: 10.1002/ijc.31931]
- 7 **Jin C**, Shi W, Wang F, Shen X, Qi J, Cong H, Yuan J, Shi L, Zhu B, Luo X, Zhang Y, Ju S. Long non-coding RNA HULC as a novel serum biomarker for diagnosis and prognosis prediction of gastric cancer. *Oncotarget* 2016; **7**: 51763-51772 [PMID: 27322075 DOI: 10.18632/oncotarget.10107]
- 8 **Zhao R**, Zhang Y, Zhang X, Yang Y, Zheng X, Li X, Liu Y, Zhang Y. Exosomal long noncoding RNA HOTTIP as potential novel diagnostic and prognostic biomarker test for gastric cancer. *Mol Cancer* 2018; **17**: 68 [PMID: 29486794 DOI: 10.1186/s12943-018-0817-x]
- 9 **Chen Q**, Su Y, He X, Zhao W, Wu C, Zhang W, Si X, Dong B, Zhao L, Gao Y, Yang X, Chen J, Lu J, Qiao X, Zhang Y. Plasma long non-coding RNA MALAT1 is associated with distant metastasis in patients with epithelial ovarian cancer. *Oncol Lett* 2016; **12**: 1361-1366 [PMID: 27446438 DOI: 10.3892/ol.2016.4800]
- 10 **Zhang X**, Mao L, Li L, He Z, Wang N, Song Y. Long noncoding RNA GIHCG functions as an oncogene and serves as a serum diagnostic biomarker for cervical cancer. *J Cancer* 2019; **10**: 672-681 [PMID: 30719165 DOI: 10.7150/jca.28525]
- 11 **Wang ZF**, Hu R, Pang JM, Zhang GZ, Yan W, Li ZN. Serum long noncoding RNA LRB1 as a potential biomarker for predicting the diagnosis and prognosis of human hepatocellular carcinoma. *Oncol Lett* 2018; **16**: 1593-1601 [PMID: 30008842 DOI: 10.3892/ol.2018.8825]
- 12 **Lu CW**, Zhou DD, Xie T, Hao JL, Pant OP, Lu CB, Liu XF. HOXA11 antisense long noncoding RNA (HOXA11-AS): A promising lncRNA in human cancers. *Cancer Med* 2018; **7**: 3792-3799 [PMID: 29992790 DOI: 10.1002/cam4.1571]
- 13 **Liu Z**, Chen Z, Fan R, Jiang B, Chen X, Chen Q, Nie F, Lu K, Sun M. Over-expressed long noncoding RNA HOXA11-AS promotes cell cycle progression and metastasis in gastric cancer. *Mol Cancer* 2017; **16**: 82 [PMID: 28441948 DOI: 10.1186/s12943-017-0651-6]
- 14 **Sun M**, Nie F, Wang Y, Zhang Z, Hou J, He D, Xie M, Xu L, De W, Wang Z, Wang J. LncRNA HOXA11-AS Promotes Proliferation and Invasion of Gastric Cancer by Scaffolding the Chromatin Modification Factors PRC2, LSD1, and DNMT1. *Cancer Res* 2016; **76**: 6299-6310 [PMID: 27651312 DOI: 10.1158/0008-5472.CAN-16-0356]
- 15 **Siegel RL**, Miller KD, Jemal A. Cancer statistics, 2015. *CA Cancer J Clin* 2015; **65**: 5-29 [PMID: 25559415 DOI: 10.3322/caac.21254]
- 16 **Tan L**, Yang Y, Shao Y, Zhang H, Guo J. Plasma lncRNA-GACAT2 is a valuable marker for the screening of gastric cancer. *Oncol Lett* 2016; **12**: 4845-4849 [PMID: 28105191 DOI: 10.3892/ol.2016.5297]
- 17 **Li W**, Jia G, Qu Y, Du Q, Liu B, Liu B. Long Non-Coding RNA (LncRNA) HOXA11-AS Promotes Breast Cancer Invasion and Metastasis by Regulating Epithelial-Mesenchymal Transition. *Med Sci Monit* 2017; **23**: 3393-3403 [PMID: 28701685]
- 18 **Yang FQ**, Zhang JQ, Jin JJ, Yang CY, Zhang WJ, Zhang HM, Zheng JH, Weng ZM. HOXA11-AS promotes the growth and invasion of renal cancer by sponging miR-146b-5p to upregulate MMP16 expression. *J Cell Physiol* 2018; **233**: 9611-9619 [PMID: 29953617 DOI: 10.1002/jcp.26864]
- 19 **Cui M**, Wang J, Li Q, Zhang J, Jia J, Zhan X. Long non-coding RNA HOXA11-AS functions as a competing endogenous RNA to regulate ROCK1 expression by sponging miR-124-3p in osteosarcoma. *Biomed Pharmacother* 2017; **92**: 437-444 [PMID: 28558357 DOI: 10.1016/j.biopha.2017.05.081]
- 20 **Yao Y**, Chen X, Lu S, Zhou C, Xu G, Yan Z, Yang J, Yu T, Chen W, Qian Y, Ding S, Tang J, Chen Y, Zhang Y. Circulating Long Noncoding RNAs as Biomarkers for Predicting Head and Neck Squamous Cell Carcinoma. *Cell Physiol Biochem* 2018; **50**: 1429-1440 [PMID: 30355951 DOI: 10.1159/000494605]
- 21 **Sun XY**, Wang XF, Cui YB, Cao XG, Zhao RH, Wei HY, Cao W, Wu W. [Expression level and clinical significance of LncRNA HOXA11-AS in esophageal squamous cell carcinoma patients]. *Zhonghua Zhong Liu Za Zhi* 2018; **40**: 186-190 [PMID: 29575836 DOI: 10.3760/cma.j.issn.0253-3766.2018.03.005]
- 22 **Mu S**, Ai L, Fan F, Sun C, Hu Y. Prognostic and clinicopathological significance of long noncoding RNA HOXA11-AS expression in human solid tumors: a meta-analysis. *Cancer Cell Int* 2018; **18**: 1 [PMID: 29308050 DOI: 10.1186/s12935-017-0498-3]
- 23 **Chen JH**, Zhou LY, Xu S, Zheng YL, Wan YF, Hu CP. Overexpression of lncRNA HOXA11-AS promotes cell epithelial-mesenchymal transition by repressing miR-200b in non-small cell lung cancer. *Cancer Cell Int* 2017; **17**: 64 [PMID: 28615992 DOI: 10.1186/s12935-017-0433-7]
- 24 **Liu B**, Li J, Liu X, Zheng M, Yang Y, Lyu Q, Jin L. Long non-coding RNA HOXA11-AS promotes the proliferation HCC cells by epigenetically silencing DUSP5. *Oncotarget* 2017; **8**: 109509-109521 [PMID: 29312625 DOI: 10.18632/oncotarget.22723]
- 25 **Erkelenz S**, Mueller WF, Evans MS, Busch A, Schöneweis K, Hertel KJ, Schaal H. Position-dependent splicing activation and repression by SR and hnRNP proteins rely on common mechanisms. *RNA* 2013; **19**: 96-102 [PMID: 23175589 DOI: 10.1261/ma.037044.112]
- 26 **Wang Y**, Xiao X, Zhang J, Choudhury R, Robertson A, Li K, Ma M, Burge CB, Wang Z. A complex network of factors with overlapping affinities represses splicing through intronic elements. *Nat Struct Mol Biol* 2013; **20**: 36-45 [PMID: 23241926 DOI: 10.1038/nsmb.2459]
- 27 **Das R**, Yu J, Zhang Z, Gygi MP, Krainer AR, Gygi SP, Reed R. SR proteins function in coupling RNAP II transcription to pre-mRNA splicing. *Mol Cell* 2007; **26**: 867-881 [PMID: 17588520 DOI: 10.1016/j.molcel.2007.05.036]

- 28 **Fregoso OI**, Das S, Akerman M, Krainer AR. Splicing-factor oncoprotein SRSF1 stabilizes p53 via RPL5 and induces cellular senescence. *Mol Cell* 2013; **50**: 56-66 [PMID: [23478443](#) DOI: [10.1016/j.molcel.2013.02.001](#)]
- 29 **Zhou X**, Wang R, Li X, Yu L, Hua D, Sun C, Shi C, Luo W, Rao C, Jiang Z, Feng Y, Wang Q, Yu S. Splicing factor SRSF1 promotes gliomagenesis via oncogenic splice-switching of MYO1B. *J Clin Invest* 2019; **129**: 676-693 [PMID: [30481162](#) DOI: [10.1172/jci120279](#)]
- 30 **Zhao X**, Li X, Zhou L, Ni J, Yan W, Ma R, Wu J, Feng J, Chen P. LncRNA HOXA11-AS drives cisplatin resistance of human LUAD cells via modulating miR-454-3p/Stat3. *Cancer Sci* 2018; **109**: 3068-3079 [PMID: [30099826](#) DOI: [10.1111/cas.13764](#)]
- 31 **Chen D**, Sun Q, Zhang L, Zhou X, Cheng X, Zhou D, Ye F, Lin J, Wang W. The lncRNA HOXA11-AS functions as a competing endogenous RNA to regulate PADI2 expression by sponging miR-125a-5p in liver metastasis of colorectal cancer. *Oncotarget* 2017; **8**: 70642-70652 [PMID: [29050308](#) DOI: [10.18632/oncotarget.19956](#)]
- 32 **Se YB**, Kim SH, Kim JY, Kim JE, Dho YS, Kim JW, Kim YH, Woo HG, Kim SH, Kang SH, Kim HJ, Kim TM, Lee ST, Choi SH, Park SH, Kim IH, Kim DG, Park CK. Underexpression of HOXA11 Is Associated with Treatment Resistance and Poor Prognosis in Glioblastoma. *Cancer Res Treat* 2017; **49**: 387-398 [PMID: [27456940](#) DOI: [10.4143/crt.2016.106](#)]
- 33 **Lu Q**, Zhao N, Zha G, Wang H, Tong Q, Xin S. LncRNA HOXA11-AS Exerts Oncogenic Functions by Repressing p21 and miR-124 in Uveal Melanoma. *DNA Cell Biol* 2017; **36**: 837-844 [PMID: [28749709](#) DOI: [10.1089/dna.2017.3808](#)]
- 34 **Kim HJ**, Eoh KJ, Kim LK, Nam EJ, Yoon SO, Kim KH, Lee JK, Kim SW, Kim YT. The long noncoding RNA HOXA11 antisense induces tumor progression and stemness maintenance in cervical cancer. *Oncotarget* 2016; **7**: 83001-83016 [PMID: [27792998](#) DOI: [10.18632/oncotarget.12863](#)]
- 35 **Zong W**, Feng W, Jiang Y, Ju S, Cui M, Jing R. Evaluating the diagnostic and prognostic value of serum long non-coding RNA CTC-497E21.4 in gastric cancer. *Clin Chem Lab Med* 2019; Epub ahead of print [PMID: [30763257](#) DOI: [10.1515/cclm-2018-0929](#)]
- 36 **Ji B**, Huang Y, Gu T, Zhang L, Li G, Zhang C. Potential diagnostic and prognostic value of plasma long noncoding RNA LINC00086 and miR-214 expression in gastric cancer. *Cancer Biomark* 2019; **24**: 249-255 [PMID: [30689553](#) DOI: [10.3233/cbm-181486](#)]
- 37 **Yörüker EE**, Keskin M, Kulle CB, Holdenrieder S, Gezer U. Diagnostic and prognostic value of circulating lncRNA H19 in gastric cancer. *Biomed Rep* 2018; **9**: 181-186 [PMID: [30083318](#) DOI: [10.3892/br.2018.1116](#)]
- 38 **Xia H**, Chen Q, Chen Y, Ge X, Leng W, Tang Q, Ren M, Chen L, Yuan D, Zhang Y, Liu M, Gong Q, Bi F. The lncRNA MALAT1 is a novel biomarker for gastric cancer metastasis. *Oncotarget* 2016; **7**: 56209-56218 [PMID: [27486823](#) DOI: [10.18632/oncotarget.10941](#)]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

