



国内版

国际版

Epstein-Barr Virus-Positive Posttransplant Lymphoproliferative Disorder Prese



All

Images

Videos

翻译成中文

关闭取词

64 Results

Any time ▾

Primary Intestinal Epstein-Barr Virus-associated Natural ...

https://www.researchgate.net/publication/324472911_Primary_Intestinal_Epstein-Barr...

Background and aims: Primary intestinal **Epstein-Barr** virus (EBV)-associated natural killer/T-cell **lymphoproliferative disorder** (PIEBV+ NK/T-LPD) is a rare clinical entity, which is difficult to ...

Abstracts and Case Studies From the College of American ...

<https://www.archivesofpathology.org/doi/10.5858/arpa.2018-0293-AB>

Abstract and **case** study poster sessions will be conducted during the 2018 College of American Pathologists Annual Meeting (CAP18), which is scheduled for October 20 to 24, 2018. The meeting will take place at the Hyatt Regency, Chicago, Illinois. The poster sessions will occur in the CAP18 Exhibit Hall. Specific dates and times for each poster session are listed below; "poster focus" times ...

EAHP2018 - eahp-sh2018.com

www.eahp-sh2018.com/wp-content/uploads/sites/81/Flip/epaper-EAHP2018 ▾

In addition, CAD was the **presenting** clinical manifestation. Cold agglutinin **disease** is often refractory to treatment and often linked to an underlying **lymphoproliferative disorder** mainly **lymphoplasmacytic lymphoma**. Only three cases of primary **bone marrow** DLBCL, **NOS** **presenting** with CAD have been reported in the literature, as separate case reports.

29 - Nutrition | Current Medical Diagnosis and Treatment ...

<https://flylib.com/books/en/2.857.1.38/1> ▾

EBV-induced lymphoproliferation gives rise to B cell lymphomas among immunodeficient patients, such as the HIV-infected individuals (HIV-infected patients, even on HAART, show higher EBV DNA loads than HIV-noninfected patients) or after transplantation (**posttransplant lymphoproliferative disorder**).

infectious complications treatment: Topics by Science.gov

<https://www.science.gov/topicpages/i/infectious+complications+treatment.html> ▾

The biopsy revealed an **Epstein-Barr virus**-positive **lymphoproliferative disorder**. Systemic disease was absent. Following discontinuation of 6-mercaptopurine, the patient was treated with **two** courses of intravenous cyclophosphamide. The lesion resolved completely and she remains disease free at 14 months following diagnosis.

Match Overview



1 Internet 15 words
crawled on 20-Jun-2019
f6publishing.blob.core.windows.net

1%

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 48114

Manuscript Type: CASE REPORT

Epstein-Barr **virus**-positive posttransplant lymphoproliferative disorder
presenting as hematochezia and enterobrosis in renal transplant recipients
in china: Two case reports

Sun ZJ *et al.* EBV-positive posttransplant lymphoproliferative disorder

Ze-Jia Sun, Xiao-Peng Hu, Bo-Han Fan, Wei Wang

Abstract

Epstein-Barr virus-positive posttransplant lymphoproliferative



ALL

IMAGES

VIDEOS

关闭取词

75 Results

Any time ▾

Primary Intestinal Epstein-Barr Virus-associated Natural ...

https://www.researchgate.net/publication/324472911_Primary_Intestinal_Epstein-Barr...

Epstein-Barr virus (EBV)-associated T-cell lymphoproliferative disease (LPD) is not uncommon in China, but gastrointestinal involvement is very rare. We report on an immunocompetent patient with ...

Epstein Barr virus induced lymphoma in a child with IPEX ...

https://www.researchgate.net/publication/6011390_Epstein_Barr_virus_induced_lymphoma...

Epstein Barr virus induced lymphoma in a child with IPEX syndrome ... correlated assay methods in vitro using a model of posttransplant lymphoproliferative disorder. ... kidney transplant ...

Abstracts and Case Studies From the College of American ...

<https://archivesofpathology.org/doi/full/10.5858/arpa.2016-0246-AB>

The final diagnosis was an EBV-related classic Hodgkin lymphoma-type posttransplant lymphoproliferative disorder, explaining the high EBV titers and the presence of hemophagocytic lymphohistiocytosis. HLH should be considered in patients presenting with ...

EAHP2018 - eahp-sh2018.com

www.eahp-sh2018.com/wp-content/uploads/sites/81/Flip/epaper-EAHP2018 ▾

Non-Hodgkin lymphoma secondary to Hodgkin lymphoma in an Adult Patient with Nijmegen Breakage Syndrome Clinico-histomorphological correlation of Epstein Barr Virus in Hodgkin's Lymphoma EBV-positive NK/T-cell lymphoma, associated with chronic active EBV infection: A case report A Case Report of Myeloproliferative Neoplasms Following Treatment of Myeloid Sarcoma Presenting as Spleen and Splenic lymph node ...

VIRAL CAUSES OF DIARRHEA - Gastroenterology Clinics

[https://www.gastro.theclinics.com/article/S0889-8553\(05\)70210-7/fulltext](https://www.gastro.theclinics.com/article/S0889-8553(05)70210-7/fulltext)

When the gastrointestinal tract is involved, patients present with diarrhea, abdominal pain, gastrointestinal bleeding, or an acute abdomen from perforation. 20 x 20 Cao, S., Cox, K., Esquivel, C. et al. Posttransplant lymphoproliferative disorders and gastrointestinal manifestations of Epstein-Barr virus infection in children following liver transplantation.

Cited by: 68

Author: Richard W. Goodgame

Publish Year: 2001

76 Results

Any time ▾

Primary Intestinal Epstein-Barr Virus-associated Natural ...

https://www.researchgate.net/publication/324472911_Primary_Intestinal_Epstein-Barr...

Epstein-Barr virus (EBV)-associated T-cell lymphoproliferative disease (LPD) is not uncommon in China, but gastrointestinal involvement is very rare. We report on an immunocompetent patient with ...

Epstein Barr virus induced lymphoma in a child with IPEX ...

https://www.researchgate.net/publication/6011390_Epstein_Barr_virus_induced_lymphoma...

Epstein Barr virus induced lymphoma in a child with IPEX syndrome ... correlated assay methods in vitro using a model of posttransplant lymphoproliferative disorder. ... kidney transplant ...

Abstracts and Case Studies From the College of American ...

<https://archivesofpathology.org/doi/full/10.5858/arpa.2016-0246-AB>

The final diagnosis was an EBV-related classic Hodgkin lymphoma-type posttransplant lymphoproliferative disorder, explaining the high EBV titers and the presence of hemophagocytic lymphohistiocytosis. HLH should be considered in patients presenting with ...

EAHP2018 - eahp-sh2018.com

www.eahp-sh2018.com/wp-content/uploads/sites/81/Flip/epaper-EAHP2018 ▾

Non-Hodgkin lymphoma secondary to Hodgkin lymphoma in an Adult Patient with Nijmegen Breakage Syndrome Clinico-histomorphological correlation of Epstein Barr Virus in Hodgkin's Lymphoma EBV-positive NK/T-cell lymphoma, associated with chronic active EBV infection: A case report A Case Report of Myeloproliferative Neoplasms Following Treatment of Myeloid Sarcoma Presenting as Spleen and Splenic lymph node ...

VIRAL CAUSES OF DIARRHEA - Gastroenterology Clinics

[https://www.gastro.theclinics.com/article/S0889-8553\(05\)70210-7/fulltext](https://www.gastro.theclinics.com/article/S0889-8553(05)70210-7/fulltext)

When the gastrointestinal tract is involved, patients present with diarrhea, abdominal pain, gastrointestinal bleeding, or an acute abdomen from perforation. 20 x 20 Cao, S., Cox, K., Esquivel, C. et al. Posttransplant lymphoproliferative disorders and gastrointestinal manifestations of Epstein-Barr virus infection in children following liver transplantation.

Cited by: 68

Author: Richard W. Goodgame

Publish Year: 2001