

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 3,850,000 条结果 (用时 0.68 秒)

So **hepatitis B virus related acute-on-chronic liver failure (HBV-ACLF)** caused by **HBV infection** is the most common type in China with various **complications** and a **mortality rate as high as 40%-90%** [1, 6, 7]. 2019年2月3日

[The Effect of Modified Sini Decoction on Survival Rates of Patients ...](#)

<https://www.hindawi.com/journals/ecam/2019/2501847/>

关于这条结果的详细信息 反馈

[Acute-on-chronic liver failure: an update | Gut](#)

<https://gut.bmj.com/content/66/3/541> - 翻译此页

作者: R Hernaez - 2017 - 被引用次数: 105 - 相关文章

The concept of **acute-on-chronic liver failure** (ACLF) has been widely used in critical ... The change in dietary patterns and lifestyle will likely **lead** to a shift on the ACLF this **complication**.40 **Mortality related** to bacterial sepsis is greater in **patients** would explain the **high risk** of nosocomial **infections** in **patients** with ACLF.

[Dexamethasone in outcome of patients with hepatitis B virus-related ...](#)

<https://www.ncbi.nlm.nih.gov/pubmed/24224656> ▾ 翻译此页

作者: JF Chen - 2014 - 被引用次数: 19 - 相关文章

BACKGROUND AND AIM: **Acute-on-chronic liver failure** (ACLF) caused by **hepatitis B virus (HBV)** is a **severe disease** with **high mortality**. ... differences in the **incidence** of **complications** (i.e. **infection**, gastrointestinal bleeding, encephalopathy, ...

[Potent antiviral therapy improves survival in acute on chronic liver ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4239490/> ▾ 翻译此页

作者: CA Philips - 2014 - 被引用次数: 26 - 相关文章

跳到**ACLF-HBV AND LIVER TRANSPLANTATION** - **Liver** transplantation in **patients** with ACLF has ... **Patients** who developed **infectious complications** ... of **acute** on **chronic liver failure** secondary ... The current concepts in management of **HBV-related** ACLF have ... of **patients** who have **high mortality rates** in ...

Name of journal: *World Journal of Clinical Cases*

Manuscript NO: 48175

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

1 Incidence of infection complication leads to high mortality in patients with hepatitis B virus-related acute-on-chronic liver failure

Wang C *et al.* Infections in patients with HBV-ACLF

Abstract

BACKGROUND

In China, hepatitis B virus-related acute-on-chronic liver failure (HBV-ACLF) is the most common liver failure characterized by serious clinical syndromes of liver decompensation with very high mortality. Bacterial and/or fungal infection is the most common complication that is associated with high short-term mortality. Bacterial translocation from the intestine, impaired hepatic clearance and immune paralysis of circulating immune cells are

Match Overview

1	Internet 133 words crawled on 28-Mar-2019 link.springer.com	3%
2	Internet 46 words crawled on 13-Mar-2019 www.glynns.co.uk	1%
3	Crossref 42 words Javier Fernández, Juan Acevedo, Reiner Weist, Thierry G ustot et al. "Bacterial and fungal infections in acute-on- ...	1%
4	Crossref 39 words Jacqueline G. O'Leary, K. Rajender Reddy, Florence Wong, Patrick S. Kamath et al. "Long-term Use of Antibiotics	1%
5	Internet 33 words crawled on 14-Jan-2019 www.frontiersin.org	1%
6	Internet 27 words crawled on 22-Oct-2017 www.impactjournals.com	1%
7	Internet 25 words crawled on 12-Aug-2014 www.ncbi.nlm.nih.gov	1%
8	Crossref 16 words Xia, Qi, Xiahong Dai, Yimin Zhang, Yongzheng Guo, Xiao wei Xu, Qian Yang, Weibo Du, Xiaoli Liu, Yuemei Chen, J	<1%
9	Internet 14 words crawled on 20-May-2018 www.spandidos-publications.com	<1%
10	Crossref 14 words "Abstracts of the 26th Annual Conference of APASL, February 15–19, 2017, Shanghai, China", Hepatology Interna	<1%
11	Internet 13 words crawled on 31-Aug-2011 www.pkuph.com.cn	<1%



All

Images

Videos

翻译成中文

关闭取词

333,000 Results

Any time ▾

Population-representative Incidence of Acute-On-Chronic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4978606>

Results: **Chronic hepatitis B virus (HBV) infection** was the **leading cause of chronic liver disease** and HBV reactivation was the most common cause of **acute hepatic event**. The 28-day **mortality** for the **ACLF patients** had a clear decline during the study period, from 50.39% in 2005 to 35.44% in 2014.

Cited by: 8

Author: Gang Qin, Jian-Guo Shao, Yong-Chang Z...

Publish Year: 2016

Infectious Complications After Liver Transplantation

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4849501>

The effect of pretransplant infections on the **incidence** and severity of posttransplant infections has been investigated and has shown conflicting results. 14, 15 It is well known that SBP is the most common **infection** among **patients** awaiting **liver** transplantation, followed by bloodstream infections, cellulitis, pneumonia, and UTIs. 16, 17 ...

Cited by: 15

Author: Maria Del Pilar Hernandez, Paul Martin, J...

Publish Year: 2015

Management of hepatitis B virus-related acute liver failure

<https://link.springer.com/article/10.1007/s12328-013-0447-1> ▾

Jan 24, 2014 · **Hepatitis B virus (HBV)** is the most important cause of **acute liver failure (ALF)** in Eastern countries [1, 2, 3]. HBV-related ALF may occur after **acute HBV infection (A-ALF)** or during an acute exacerbation (flare) of **chronic HBV infection (C-ALF)**.

Cited by: 4

Author: Makoto Oketani, Hirofumi Uto, Akio Ido, ...

Publish Year: 2014

Acute Liver Failure - Cleveland Clinic

www.clevelandclinicmeded.com/medicalpubs/diseasemanagement/hepatology/acute-liver-failure

INTRODUCTION

INCIDENCE

DEFINITION

CLASSIFICATION



Acute liver failure (ALF) is a rare but a life-threatening condition. ALF causes severe injury and massive necrosis of hepatocytes resulting in severe liver dysfunction that can lead to multi-organ failure and death. It can occur in patients without preexisting liver disease and cause rapid deterioration of liver function within days. Patients with ALF are almost always managed in an intensive care unit and in some

[国内版](#)[国际版](#)

Incidence of infection complication is associated with high mortality in patients wi

[All](#)[Images](#)[Videos](#)[翻译成中文](#)[开启取词](#)

71,400 Results

Any time ▾

Epidemiology of Hepatitis B in the United States

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3290915>

Hepatitis B virus (HBV) remains an important cause of acute and chronic liver disease globally and in the United States. An encouraging trend is that the incidence of acute hepatitis B in the United States declined as much as 80% between 1987 and 2004, attributable to effective vaccination programs as well as universal precautions in needle use and in healthcare in general.

Cited by: 201

Author: W. Ray Kim

Publish Year: 2009

Characteristics, Diagnosis and Prognosis of Acute-on ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4857102>

Abstract. Given that infection with HBV may be associated with disease flares, which can lead to rapid deterioration in liver function despite introduction of effective antiviral therapy, it is unclear whether criteria developed for non-HBV induced ACLF can also be applied to patients with HBV related ACLF.

Cited by: 34

Author: H Li, Ly Chen, Nn Zhang, St Li, B Zeng, ...

Publish Year: 2016

Is HBV viral load at admission associated with development ...

<https://bmcinfectdis.biomedcentral.com/articles/10.1186/s12879-019-3988-1> ▾

Apr 30, 2019 · Hepatitis B virus (HBV) reactivation is one of the most common precipitating events associated with acute decompensation (AD) or acute-on-chronic liver failure (ACLF) in chronic hepatitis B (CHB)-related cirrhotic patients.

Author: Jian-Hua Lei, Feng Peng, Zi Chen, Xi...

Publish Year: 2019

Hepatitis B - Complications | BMJ Best Practice

<https://bestpractice.bmj.com/topics/en-gb/127/complications> ▾

Most people are asymptomatic, although some will present with complications such as cirrhosis, hepatocellular carcinoma, or liver failure. People from endemic areas (Asia, Africa), or injection drug users and those with high-risk sexual behaviours, are at an increased risk for infection. Serologi...

Association of hepatitis B infection with high-risk ...

<https://bmcmusculoskeletdisord.biomedcentral.com/articles/10.1186/s12891-019-2535-y> ▾

Apr 11, 2019 · A total number of 196 patients who underwent primary TJAs were involved in the study, including 49 patients with hepatitis B and 147 non-hepatitis B subjects. Among all the patients with TJAs, 5.5% of patients were infected with HBV for the first time. The incidence rate of complications in patients after arthroplasty with hepatitis B infection ...

Author: Jin Wang, Guanglei Zhao, Jie Chen, J...

Publish Year: 2019

Hepatitis B virus-related decompensated liver cirrhosis ...